

VIDYAPEETH



BATCH CODE: 19-PJ301EA 2025

SUBJECT NAME: CHEMISTRY

CHAPTER NAME:
Atomic Structure

ATDB.uno



Lecture No. 04

By – Swapnil Sir



Today's Goal



Subtopic

- # Dual nature
 - # Spectrum of Hydrogen
-

ATDB.uno

Dual nature $\rightarrow$

de-Broglie relation

Plank's eq. $E = \frac{hc}{\lambda}$ — ①

Einstein's eq. $E = mc^2$

$\Rightarrow \frac{hc}{\lambda} = mc^2$

$\Rightarrow \frac{h}{\lambda} = m \cdot c$

$\frac{h}{\lambda} = (m \cdot v) = p$

$p \cdot \lambda = h$

$\downarrow$

particle nature

$\downarrow$

wave nature

any moving object associated with dual nature
wave & particle nature & product
always constant.wave associated $\rightarrow$ de-Broglie's wavelength ✓ $\lambda = \frac{h}{p}$

$$\lambda = \frac{h}{p}$$

$$\lambda = \frac{h}{m \cdot v}$$

#1

#2

$$K.E. = \frac{1}{2} m v^2$$

$$\Rightarrow 2 \cdot m \cdot (K.E.) = m^2 v^2$$

$$\Rightarrow p^2 = 2 m \cdot (K.E.)$$

$$p = \sqrt{2 m (K.E.)}$$

$$\lambda = \frac{h}{\sqrt{2 m (K.E.)}}$$

#2

#3 a charge particle
accelerated through a potential
V volt.

$$(K.E.) = \underline{\underline{q \cdot V}}$$

$$\lambda = \frac{h}{\sqrt{2 \cdot m \cdot q \cdot V}}$$

#3

V = Potential (Volt)



Spectrum $\Rightarrow$

1

Emmision

Absorptive

2


Line

Band

3

Continuous

discontinuous



Spectrum of Hydrogen (Line) $n_2 > n_1$

$$E_{n_1} = -K \frac{Z^2}{n_1^2}$$

$$E_{n_2} = -K \cdot \frac{Z^2}{n_2^2}$$

ATDB.uno

$$\Rightarrow E_{n_2} - E_{n_1} = \Delta E = K Z^2 \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] \quad \text{--- ①}$$

Planks Theory Emitted radiation

$$\Delta E = \frac{hc}{\lambda} \quad \text{--- ②}$$

$$\frac{hc}{\lambda} = KZ^2 \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$$

$$\Rightarrow \frac{1}{\lambda} = \left(\frac{K}{hc} \right) Z^2 \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$$


$$\Rightarrow \boxed{\frac{1}{\lambda} = R_H \cdot Z^2 \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]}$$

R_H = Rydberg's constant (cm^{-1})

n_1 = Returning level

n_2 = from where returning

ATDB.uno



$$E \propto \frac{1}{\lambda}$$

$\rightarrow \lambda_{\text{max.}} \rightarrow 4^{\text{th}} \text{ S.L.}$
 $\rightarrow \lambda_{\text{min.}} \rightarrow \text{from } \infty$

$n_1 = 1$

Lyman series

$n_1 = 2$

Balmer series
(Series of colorful lines)

$n_1 = 3$

Paschen series

$n_1 = 4$

Brackett-

$n_1 = 5$

Pfund

$n_1 = 6$

Humphrey

$\frac{1}{\lambda} = R_H \left[1 - \frac{1}{n^2} \right]$

$n > 1$

$\frac{1}{\lambda} = R_H \left[\frac{1}{4} - \frac{1}{n^2} \right]$

$n > 2$

$\frac{1}{\lambda} = R_H \left[\frac{1}{9} - \frac{1}{n^2} \right]$

$n > 3$

$\lambda = \text{wavelength}$


$\frac{1}{\lambda} = \text{wave number}$



ATDB.uno

	I st	II nd	III rd	IV th	λ_{min}
Lyman	$2 \rightarrow 1$	$3 \rightarrow 1$	$4 \rightarrow 1$	$5 \rightarrow 1$	$\infty \rightarrow 1$
Balmer	$3 \rightarrow 2$	$4 \rightarrow 2$	$5 \rightarrow 2$	$6 \rightarrow 2$	$\infty \rightarrow 2$
Paschen	$4 \rightarrow 3$	$5 \rightarrow 3$	$6 \rightarrow 3$	$7 \rightarrow 3$	$\infty \rightarrow 3$

Heisenberg's Uncertainty principle.



Uncertainty in momentum = Δp

Uncertainty in position = Δx

$$\Delta p \cdot \Delta x \geq \frac{h}{4\pi}$$

ATDB.uno

The product of uncertainty in momentum & uncertainty in position is always greater than or equal to $\frac{h}{4\pi}$

$$\Delta p \propto \frac{1}{\Delta x}$$



$$\Delta p \cdot \Delta x = \frac{h}{4\pi}$$



*28 *

$$\lambda = \frac{h}{mv}$$

$$mv\lambda = \frac{n \cdot h}{2\pi}$$

$$\lambda = \frac{\cancel{h} \cdot 2\pi r_n}{n\cancel{h}}$$

$$2\pi \times r_0 \left(\frac{\cancel{16}^4}{\cancel{1}} \right) = \underline{8\pi r_0}$$

ATDB.uno

$$\lambda = \frac{2\pi r_n}{n}$$



$$\begin{aligned}
 K.E. &= \frac{1}{2} m v^2 \\
 &= \frac{1}{2} \left[\frac{1}{4\pi\epsilon_0} \cdot \frac{Ze^2}{r} \right] \\
 &= \frac{1}{2} \cdot \frac{1}{4\pi\epsilon_0} \cdot \frac{1 \times e^2}{r_0 \cdot 4} \\
 &= \frac{1}{32\pi\epsilon_0} \frac{e^2}{r_0} \\
 &= \frac{2 \times 4 \times \pi^2 \times 4 \times 10}{320 \times \pi^2}
 \end{aligned}$$

$$\begin{aligned}
 m v r &= \frac{n h}{2\pi} \\
 K.E. &= \frac{1}{2} m \left(\frac{h^2 h^2}{4\pi^2 \times m^2 \times r^2} \right) \\
 &= \frac{1}{2} \frac{h^2 h^2 \cdot 2^2}{4\pi^2 \times m \times r_0^2 \times h^4}
 \end{aligned}$$


$$\begin{aligned}
 &= \frac{1}{2} \frac{h^2}{4\pi^2 \times m \cdot r_0^2 \times h^2} = \frac{h^2}{m a_0^2} \\
 &= \frac{h^2}{2 \times 4 \times \pi^2 \times 4} (m a_0^2)
 \end{aligned}$$



Rate your today's class in the Batch



Solve the DPP and check Solution



"SCAN" to join our "TELEGRAM" channel

VIDYAPEETH

WORK, POWER AND ENERGY

DPP-1 (JAP/046)

[Introduction, Definition of work, work done by constant force, Area under force-displacement curve]

1. A particle moves from position $\vec{r}_1 = 3\hat{i} + 2\hat{j} - 6\hat{k}$ to position $\vec{r}_2 = 14\hat{i} + 13\hat{j} + 9\hat{k}$ under the action of force $-4\hat{i} + \hat{j} + 3\hat{k}$ N. The work done by this force will be

(A) 100 J
(B) 50 J

(A) 8×10^{-2} joules
(B) 16×10^{-2} joules
(C) $A = 10 \times 10^{-4}$ joules

Thank You!!!!

Check your performance through Student Dashboard