

# 01

## UNCERTAINTY PRINCIPLE

$$\Delta x \cdot \Delta p \geq \frac{h}{4\pi}$$

$$\Delta x \cdot m\Delta v \geq \frac{h}{4\pi}$$

Q. According to Heisenberg's uncertainty principle,  $\Delta x \cdot \Delta p \geq \frac{h}{4\pi}$  which of the following is correct ?

- If  $\Delta x = 0$  then  $\Delta p = \infty$
- If  $\Delta v = 0$  then  $\Delta p = 0$
- If  $\Delta p = 0$  then  $\Delta x = \infty$
- All are correct

Q. Find uncertainty in velocity, if uncertainty position is equal to uncertainty in momentum.

- $\frac{h}{2\sqrt{m\pi}}$
- $\frac{1}{2m} \sqrt{\frac{h}{\pi}}$
- $\frac{1}{m} \sqrt{\frac{h}{\pi}}$
- $\frac{1}{2} \sqrt{\frac{h}{m\pi}}$

Q. The uncertainty involved in the measurement of velocity within a distance of  $0.1 \text{ \AA}$  is:

- $5.79 \times 10^5 \text{ m/s}$
- $5.79 \times 10^7 \text{ m/s}$
- $5.79 \times 10^8 \text{ m/s}$
- $5.79 \times 10^9 \text{ m/s}$

## PRINCIPLE QUANTUM NUMBER

In  $n^{\text{th}}$  Shell ,  
 Number of subshells =  $n$   
 Number of orbitals =  $n^2$   
 Max. number of electrons =  $2n^2$

Q. Find angular momentum of  
 (i) 2s orbital (ii) 3d orbital  
 (iii) 4p orbital (iv)  $e^-$  in 4<sup>th</sup> orbit

## AZIMUTHAL QUANTUM NUMBER

- It describes shell or orbit  
 $n = 1, 2, 3, 4, \dots$   
 K, L, M, N,  $\dots$
- It describes size & energy of shell.  
 $r \propto n^2$      $E \propto \frac{1}{n^2}$
- It defines the angular momentum  
 $mvr = \frac{nh}{2\pi}$

Q. Find maximum no. of  $e^-$  having  
 (i)  $n=4, s = -1/2$  (ii)  $n=3, l=1, m=0$   
 (iii)  $n=2, l=0$  (iv)  $n=3, l=1$

## MAGNETIC QUANTUM NUMBER

- It describes subshell value from 0 to  $n-1$   
 $l=0 \rightarrow s$      $l=2 \rightarrow d$   
 $l=1 \rightarrow p$      $l=3 \rightarrow f$
- Orbital angular momentum  
 $= \sqrt{l(l+1)} \hbar, \hbar = \frac{h}{2\pi}$
- Maximum no. of orbital in a subshell =  $2l + 1$   
 Maximum no. of electrons in a subshell =  $4l + 2$

If  $l=2$   
 1) Orbital = d  
 2) No. of orbitals =  $2(2+1)=5$   
 ( $d_{xy}, d_{xz}, d_{yz}, d_{x^2-y^2}, d_{z^2}$ )  
 3) Total  $e^-s = 2(2l+1) = 10 e^-s$   
 4) Orbital angular momentum =  $\sqrt{2(2+1)} \hbar = \sqrt{6} \hbar$

## SPIN QUANTUM NUMBER

Value of  $m = -l \leq m \leq l$   
 Total values of  $m = 2l + 1$

$n = 4$   
 $l = 0 \quad m = 0$   
 $l = 1 \quad m = -1, 0, +1$   
 $l = 2 \quad m = -2, -1, 0, +1, +2$   
 $l = 3 \quad m = -3, -2, -1, 0, +1, +2, +3$

SPIN  
 — CLOCKWISE (+ 1/2)  
 — ANTICLOCKWISE (- 1/2)

# ATDB.uno

## STRUCTURE OF ATOM

## ENERGY OF ORBITALS

- Mono electronic species  
 Energy defined upon  $n$   
 $1s < 2s = 2p < 3s = 3p = 3d$
- Multi electronic species  
 $3s < 3p < 4s < 3d$

**(n+l) rule**

$\rightarrow As (n + l) \uparrow, E \uparrow$   
 $\rightarrow If (n + l) \text{ is same, then } n \uparrow E \uparrow$

Orbital	2s	3d
(n+l) value	$n = 2$ $l = 0$ $n+l = 2$	$n = 3$ $l = 2$ $n+l = 5$

## SHAPE OF ORBITALS

- s orbital - Spherical shape
- p orbital - dumb bell shape
- d orbital - double dumb bell shape

## NODES

$\Psi \rightarrow e^-$  wave function  
 $\Psi^2 \rightarrow$  probability of finding the electrons

- \* Node  $\rightarrow$  Probability of finding the electron is zero.
- \* Node plane  $\rightarrow$  Plane; where  $\Psi^2 = 0$
- \* Radial nodes  $\rightarrow n - l - 1$
- \* Angular nodes =  $l$
- \* Total nodes =  $n - 1$

## FILLING OF ATOMIC ORBITAL

- Aufbau principle**  
 Electrons are filled in the increasing order of energy  
 $1s < 2s < 2p < 3s < 3p < 4s < 3d \dots$
- Pauli's exclusion principle**  
 No two electrons can have same four quantum numbers  
 $1s^3$  - against Pauli's exclusion principle
- Hund's rule**  
 Pairing is only takes place after each orbital is singly occupied.  
 $\uparrow \downarrow \uparrow \uparrow$  - Against Hund's rule

