



CHAPTER

1

Some Basic Concepts of Chemistry



Topicwise Weightage of JEE Main 6 Years Paper (124 Sets)



“How’s the Josh?” for these Topics: *Mark your confidence level in the last six months of the topic (Low-L, Medium-M, High-H)*

MATTER

A substance which occupies space, possesses mass and can be felt by any one or more of the five senses is called matter.

Physical Classification of Matter

It is based on physical state under ordinary conditions of temperature and pressure.

- Solid:** A substance is said to be solid if it possesses a definite volume and a definite shape, e.g. sugar, iron, gold, wood etc.
- Liquid:** A substance is said to be liquid if it possesses a definite volume but not definite shape. They take up the shape of the container, e.g. water, milk, oil, mercury, alcohol etc.
- Gas:** A substance is said to be gas if it neither possesses a definite volume nor a definite shape. This is because they fill up the whole container, e.g. Hydrogen (H_2), Oxygen (O_2), Carbon dioxide (CO_2) etc.

Chemical Classification of Matter

- Pure Substance:** A material containing only one type of substance. Pure Substance can not be separated into simpler substance by physical method.

e.g.: Element = Na, Mg, Ca etc.
Compound = HCl , H_2O , CO_2 , HNO_3 etc.

Pure substance is classified into two types:

- Element** (II) **Compound**
 - Element:** The pure substance containing only one kind of atoms. It is classified into 3 types
 - Metal $\rightarrow$ Zn, Cu, Hg, Ag, Sn, Pb etc.
 - Non-metal $\rightarrow$ N_2 , O_2 , Cl_2 , Br_2 , F_2 , P_4 , S_8 etc.
 - Metalloids $\rightarrow$ B, Si, As, Te etc.
 - Compound:** It is defined as pure substance containing more than one kind of elements or atoms which are combined together in a fixed proportion by weight and which can be decomposed into simpler substance by the suitable chemical method. The properties of a compound are completely different from those of its constituent element, e.g. HCl , H_2O , H_2SO_4 , HClO_4 , HNO_3 etc.
- Mixture:** A material which contain more than one type of substances and which are mixed in any ratio by weight is called as mixture. The property of the mixture is the property of its components. The mixture can be separated by simple physical method.

Classification of Mixture

- Homogeneous mixture:** The mixture, in which all the components are present uniformly is called as

homogeneous mixture. Components of mixture are present in asinglephase, e.g. Water+ Salt, Water+ Sugar, Water+alcohol. Homogeneous substances are of two types:

(a) **Pure substances:** Substances which have definite and constant chemical composition are known as pure substances. For example, all elements and compounds are pure substances.

(b) **Solutions:** A homogenous mixture of two or more pure substances is known as a solution. For example, air, a mixture of NaCl and water, alcohol and water, etc. A solution does not have a definite composition.

(ii) **Heterogeneous mixture:** The mixture in which all the components are present non-uniformly, e.g. Water + Sand, Water + Oil, blood, petrol etc.

ATOMS AND MOLECULES

Atom: The term atom was introduced by Dalton. Atom is the smallest particle of matter that takes part in a chemical reaction. Atom is also defined as the smallest particle of an element that retains all the properties of an element.

Molecule: The term molecule was introduced by Avogadro. Molecule is the smallest particle of matter that exists independently and is formed by the combination of atoms. Molecule is also defined as the smallest particle of matter that can exist and retains all the properties of that substance.

Physical and Chemical Properties

(a) **Physical Property:** The property which can be measured without changing the chemical composition of the substance is known as physical property like mass, volume, density, refractive index etc.

(b) **Chemical Property:** The property which can be evaluated at the cost of matter itself is known as chemical property. For example combustible nature of hydrogen gas can be verified by burning of hydrogen. The sweet taste of sugar by consuming it.

Physical Quantity and their Measurements in Chemistry

Physical Quantities

All quantities that can be measured are called physical quantities. eg. time, length, mass, force, work done, etc.

Fundamental Quantities

A set of physical quantities which are completely independent of each other but all other physical quantities can be expressed in terms of these physical quantities is called set of Fundamental Quantities. Fundamental units are those units which can neither be derived from one another nor they can be further resolved into any other units. The Fundamental Quantities that are currently being accepted by the scientific community are mass, time, length, current, temperature, luminous intensity and amount of substance.

International System (SI) of Units

Table: SI base quantities and their units			
S. No.	Physical quantity	unit	Symbol
1	Length	metre	m
2	Mass	kilogram	kg
3	Time	second	s
4	Temperature	kelvin	K
5	Electric current	ampere	A
6	Luminous Intensity	candela	cd
7	Amount of substance	mole	mol

SCIENTIFIC NOTATION

If a number *P* can be expressed as

$$P = A \times 10^x$$

where $1 \leq A < 10$, this is called Scientific Notation and *x* is called order of magnitude of the number.

SI Prefixes: The magnitudes of physical quantities vary over a wide range. The mass of an electron is 9.1×10^{-31} kg and that of our earth is about 6×10^{24} kg. Standard prefixes for certain power of 10. Table shows these prefixes:

Power of 10	Prefix	Symbol
12	tera	T
9	giga	G
6	mega	M
3	kilo	k
2	hecto	h
1	deca	da
−1	deci	d
−2	centi	c
−3	milli	m
−6	micro	µ
−9	nano	n
−12	pico	p
−15	femto	f

SIGNIFICANT FIGURES OR DIGITS

The significant figures (SF) in a measurement are the figures or digits that are known with certainty plus one that is uncertain. Larger the number of significant figures obtained in a measurement, greater is its accuracy and vice versa.

1. Rules to find out the number of significant figures:

- I Rule:** All the non-zero digits are significant e.g. 1984 has 4 SF.
- II Rule:** All the zeros between two non-zero digits are significant, e.g. 10806 has 5 SF.
- III Rule:** All the zeros to the left of first non-zero digit are not significant, e.g.00108 has 3 SF.
- IV Rule:** If the number is less than 1, zeros on the right of the decimal point but to the left of the first non-zero digit are not significant, e.g. 0.002308 has 4 SF.

V Rule: The trailing zeros (zeros to the right of the last non-zero digit) in a number with a decimal point are significant, e.g. 01.080 has 4 SF.

VI Rule: The trailing zeros in a number without a decimal point are not significant e.g. 010100 has 3 SF.

VII Rule: When the number is expressed in exponential form, the exponential term does not affect the number of S.F. For example in $x = 12.3 = 1.23 \times 10^1 = .123 \times 10^2 = 0.0123 \times 10^3 = 123 \times 10^{-1}$, each term has 3 SF only. (Note: It has 3 significant figure in each expression.)

2. Rules for arithmetical operations with significant figures:

I Rule: In addition or subtraction the number of decimal places in the result should be equal to the number of decimal places of that term in the operation which contain lesser number of decimal places, e.g. $12.587 - 12.5 = 0.087 = 0.1$ ($\because$ second term contain lesser i.e. one decimal place)

II Rule: In multiplication or division, the number of SF in the product or quotient is same as the smallest number of SF in any of the factors, e.g. $5.0 \times 0.125 = 0.625 = 0.62$

To avoid the confusion regarding the trailing zeros of the numbers without the decimal point the best way is to report every measurement in scientific notation (in the power of 10). In this notation every number is expressed in the form $a \times 10^b$, where a is the base number between 1 and 10 and b is any positive or negative exponent of 10. The base number a is written in decimal form with the decimal after the first digit. While counting the number of SF only base number is considered (Rule VII).

❖ The change in the unit of measurement of a quantity does not affect the number of SF. For example $23.08 \text{ cm} = 23.08 \text{ mm} = 0.02308 \text{ m} = 23080 \mu\text{m}$ each term has 4 SF.

ACCURACY AND PRECISION

The accuracy of a measurement is a measure of how close the measured value is to the true value of the quantity. Precision tells us to what resolution or limit the quantity is measured.

Illustration: Assertion: If the true value for a result is 2.00 g and a student ‘A’ takes two measurements and reports the results as 1.95 g and 1.93 g. These values are precise as they are close to each other but are not accurate.

Reason: Precision refers to the closeness of various measurements for the same quantity. Whereas, accuracy is the agreement of a particular value to the true value of the result.

In the light of the above statement, choose the correct answer from the options given below:

- (a) Both Assertion and Reason are true and Reason is correct explanation of Assertion.
- (b) Both Assertion and Reason are true but Reason is not a correct explanation of Assertion.
- (c) Assertion is true but Reason is false.
- (d) Assertion is false but Reason is true.

Sol. (a)

Illustration: Assertion: 100 has only one significant figure, but 100.0 has three significant figures and 100.0 has four significant figures.

Some Basic Concepts of Chemistry

Reason: Zeros at the end or right of a number are significant provided they are on the right side of the decimal point.

In the light of the above statement, choose the correct answer from the options given below:

- (a) Both Assertion and Reason are true and Reason is correct explanation of Assertion.
- (b) Both Assertion and Reason are true but Reason is not a correct explanation of Assertion.
- (c) Assertion is true but Reason is false.
- (d) Assertion is false but Reason is true.

Sol. (a)

Illustration: Assertion: All the zeros to the left of first non-zero digit are not significant.

Reason: Zeros at the end or right of a non-zero digit are significant provided they are on the right side of the decimal point.


In the light of the above statement, choose the correct answer from the options given below:

- (a) Both Assertion and Reason are true and Reason is correct explanation of Assertion.
- (b) Both Assertion and Reason are true but Reason is not a correct explanation of Assertion.
- (c) Assertion is true but Reason is false.
- (d) Assertion is false but Reason is true.

Sol. (b)

LAWS OF CHEMICAL COMBINATION

The law of conservation of mass (Law of Indestructibility of matter) According to this law, the mass can neither be created nor be destroyed in a balanced chemical reaction but one form is changed into another form. In a chemical change total mass remains conserved i.e., total mass before the reaction is always equal to total mass after the reaction.



Mass before the reaction = $1 \times 2 + \frac{1}{2} \times 32 = 18 \text{ gm}$

Mass after the reaction = $1 \times 18 = 18 \text{ gm}$

[Total mass of reactants = Total mass of products + Mass of unreacted reactants]

Law of constant composition (definite proportions): All chemical compounds are found to have constant composition of elements irrespective of their method of preparation or sources. E.g. In H_2O , hydrogen & oxygen combine in 2 : 1 molar ratio, this ratio remains constant whether it is tap water, river water or seawater or produced by any chemical reaction.

Law of Multiple Proportions: When one element combines with the other element to form two or more different compounds, the mass of one element, which combines with a constant mass of the other bear a simple ratio to one another.

Nitrogen and oxygen combine to form five oxides: weights of oxygen which combine with the fixed weight of nitrogen in these oxides are calculated as under:


Oxide	Ratio of weights of nitrogen and oxygen
N ₂ O	28 : 16
NO	28 : 32
N ₂ O ₃	28 : 48
N ₂ O ₄	28 : 64
N ₂ O ₅	28 : 80

Number of parts by weight of oxygen which combine with 14 parts by weight of nitrogen from the above are 8, 16, 24, 32 and 40 respectively. Their ratio is 1 : 2 : 3 : 4 : 5, which is a simple whole number ratio. Hence, the law of multiple proportion is illustrated.


ADVANCED LEARNING

Law of Reciprocal Proportion: The ratio of the masses of two elements A and B which combine separately with a fixed mass of the third element C is either the same or some simple multiple of the ratio of the masses in which A and B combine directly with each other.

Like CH₄, CO₂ and H₂O
 CH₄ → C : H = 12 : 4
 CO₂ → C : O = 12 : 32
 H₂O → H : O = 2 : 16



Gay-Lussac's Law of Combining Volume: Gases combine in a simple whole number ratio of their volumes provided all measurements should be done at the same temperature and pressure.



Avogadro's hypothesis: Equal volume of all gases have equal number of molecules (not atoms) at same temperature and pressure condition. **S.T.P.** (Standard Temperature and Pressure) Temperature = 0°C or 273 K, Pressure = 1 atm = 760 mm of Hg. Volume of one mole of gas at STP is found to be experimentally equal to 22.4 litres which is known as molar volume.

- ❖ Measuring the volume is equivalent to counting the number of molecules of the gas.

Train Your Brain

Example 1: A 15.9 g sample of sodium carbonate is added to a solution of acetic acid weighing 20.0 g. The two substances react, releasing carbon dioxide gas to the atmosphere. After reaction, the contents of the reaction vessel weigh 29.3 g. What is the mass of carbon dioxide given off during the reaction?

Sol. The total mass of reactants taken = 15.9 + 20.0 = 35.9 gm. From the conservation of mass, the final mass of the contents of the vessel should also be 35.9 gm. But it is only 29.3 gm. The difference is due to the mass of released carbon dioxide gas. Hence, the mass of carbon dioxide gas released = 35.9 – 29.3 = 6.6 gm

Example 2: The following are the results of analysis of two samples of the same or two different compounds of phosphorus and chlorine. From these results, decide whether the two samples are from the same or different compounds. Also state the law, which will be obeyed by the given samples.

	Amount of P	Amount of Cl
Compound A	1.156 gm	3.971 gm
Compound B	1.542 gm	5.297 gm

Sol. The mass ratio of phosphorus and chlorine in compound A, $m_P : m_{Cl} = 1.156 : 3.971 = 0.2911 : 1.000$

The mass ratio of phosphorus and chlorine in compound B, $m_P : m_{Cl} = 1.542 : 5.297 = 0.2911 : 1.000$

As the mass ratio is same, both the compounds are **same** and the samples obey the **law of definite proportion**.

Example 3: 2.5 ml of a gaseous hydrocarbon exactly requires 12.5 ml oxygen for complete combustion and produces 7.5 ml carbon dioxide and 10.0 ml water vapour. All the volumes are measured at the same pressure and temperature. Show that the data illustrates Gay Lussac's law of volume combination.

Sol. $V_{\text{hydrocarbon}} : V_{\text{oxygen}} : V_{\text{carbon dioxide}} : V_{\text{water vapour}}$
 = 2.5 : 12.5 : 7.5 : 10.0
 = 1 : 5 : 3 : 4 (simple whole no. ratio)
 Hence, the data is according to the **law of volume combination**.

Concept Application

- A sample of pure carbon dioxide, irrespective of its source contains 27.27% carbon and 72.73% oxygen. The data support:
 - Law of constant composition.
 - Law of conservation of mass.
 - Law of reciprocal proportions.
 - Law of multiple proportions.
- The percentage of hydrogen in water and hydrogen peroxide is 11.1 and 5.9 respectively. These figures illustrate:
 - Law of multiple proportions.
 - Law of conservation of mass.
 - Law of constant proportions.
 - Law of combining volumes.
- 1.0 g of an oxide of A contained 0.5 g of A. 4.0 g of another oxide of A contained 1.6 g of A. The data indicate the law of:
 - Reciprocal proportions.
 - Constant proportions.
 - Conservation of energy.
 - Multiple proportions.
- Carbon is found to form two oxides which contain 42.9% & 27.3% of carbon respectively show that these figures shows the
 - Law of multiple proportion
 - Law of definite proportion
 - Law of mass conservation
 - All of these

ATOMIC MASS & MOLECULAR MASS

Relative Atomic Mass: One of the most important concept from Dalton's atomic theory was that of relative atomic mass or relative atomic weight. This is done by expressing mass of one atom with respect to a fixed standard. Dalton used hydrogen as the standard ($H = 1$). Later on oxygen ($O = 16$) replaced hydrogen as the reference.

- ❖ The present standard unit which was adopted internationally in 1961, is based on the mass of one carbon-12 isotopic atom, taken as exactly 12.000 u (amu).

Relative atomic mass (R.A.M)

$$= \frac{\text{Mass of one atom of an element}}{\frac{1}{12} \times \text{Mass of one } C^{12} \text{ atom}}$$

Atomic Mass Unit (or amu): The atomic mass unit (amu) is equal to $\left(\frac{1}{12}\right)^{\text{th}}$ of mass of one atom of carbon-12 isotope.

$$\therefore 1 \text{ amu} = \frac{1}{12} \times \text{mass of one } C^{12} \text{ isotopic atom}$$

$$\simeq \text{mass of one nucleon in } C^{12} \text{ atom.}$$

$$1 \text{ amu} = 1.66 \times 10^{-24} \text{ gm or } 1.66 \times 10^{-27} \text{ kg}$$

- ❖ Today, amu has been replaced by 'u' which is known as unified atomic mass
- ❖ **One amu is also called One Dalton (Da).**
- ❖ Atomic mass = R.A.M $\times$ 1 amu

- ❖ Relative atomic mass indicates the number of nucleons present in the atom.

Average / Mean Atomic Mass

The weighted average of the isotopic masses of the element's naturally occurring isotopes.

Mathematically, average atomic mass of X (A_x)

$$= \frac{a_1x_1 + a_2x_2 + \dots + a_nx_n}{100}$$

Where:

a_1, a_2, a_3 atomic mass of isotopes.

and x_1, x_2, x_3 mole% or % of natural abundance of isotopes.

Key Note

- Atomic weights of many elements are not whole numbers due to the presence of stable isotopes.
- The number of atoms of a particular isotope present in 100 atoms of a natural sample of that element is called its relative abundance which always remains constant for a given element.
- Shortcut for % determination if average atomic weight is given for X having isotopes XA & XB .

$$\% \text{ of } XA = \left| \frac{\text{Average atomic weight} - \text{wt of } X^B}{\text{difference in weight of } X^A \text{ \& } X^B} \right| \times 100$$

Illustration: Naturally occurring chlorine is 75% Cl^{35} which has an atomic mass of 35 amu and 25% Cl^{37} which has a mass of 37 amu. Calculate the average atomic mass of chlorine:

(a) 35.5amu (b) 36.5amu (c) 71amu (d) 72amu

Sol. (a) Average atomic mass

$$= \frac{(\% \text{ of I isotope}) \times (\text{its A.M}) + (\% \text{ of II isotope}) \times (\text{its A.M.})}{100}$$

$$= \frac{75 \times 35 + 25 \times 37}{100} = 35.5 \text{ amu}$$

Relative molecular mass:

$$= \frac{\text{mass of one molecule of the substance}}{\frac{1}{12} \times \text{mass of one } C^{12} - \text{atom}}$$

- ❖ Molecular mass = Relative molecular mass $\times$ 1 amu

Mean Molar Mass or Molecular Mass

The average molar mass of the different substance present in the container = $\frac{n_1M_1 + n_2M_2 + \dots n_nM_n}{n_1 + n_2 + \dots n_n}$.

Where:

M_1, M_2, M_3 are molar masses.

n_1, n_2, n_3 are moles of substances.

Formula Mass

The formula mass of a substance is defined as the sum of the atomic masses of constituent atoms in an ionic compound. This is generally used for ionic compounds which do not contain discrete molecules, but ions as their constituent units.

For example: Formula mass of NaCl is:

Formula mass = mass of sodium atom + mass of chlorine atom
= (23 + 35.5) u = 58.5 u

Illustration: The molar composition of polluted air is as follows:

Gas	At. wt.	mole percentage composition
Oxygen	16	16%
Nitrogen	14	80%
Carbon dioxide	-	03%
Sulphur dioxide	-	01%

What is the average molecular weight of the given polluted air?
(Given, atomic weights of C and S are 12 and 32 respectively.

$$\text{Sol. } M_{\text{avg}} = \frac{\sum_{j=1}^{j=n} n_j M_j}{\sum_{j=1}^{j=n} n_j} \quad \text{Here } \sum_{j=1}^{j=n} n_j = 100$$

$$\therefore M_{\text{avg}} = \frac{16 \times 32 + 80 \times 28 + 44 \times 3 + 64 \times 1}{100}$$

$$= \frac{512 + 2240 + 132 + 64}{100} = \frac{2948}{100} = 29.48 \text{ Ans.}$$

Train Your Brain

Example 4: Find the relative atomic mass of ‘O’ atom and its atomic mass.

Sol. The number of nucleons present in ‘O’ atom is 16.

∴ Relative atomic mass of ‘O’ atom = 16.

Atomic mass = R.A.M × 1 amu = 16 × 1 amu = 16 amu

Example 5: The weight of one atom of uranium is 235 amu.

Its actual weight in g is:

Sol. $235 \times 1.67 \times 10^{-24}$ g
 $= 3.95 \times 10^{-22}$ g

Concept Application

- The Relative molecular mass of ammonia is:
 (a) 17 (b) 22 (c) 28 (d) 44
- The mass of an atom of sodium is:
 (a) 23 amu (b) 23 gm
 (c) 46 amu (d) 12 amu
- The atomic mass & molecular mass of hydrogen is:
 (a) 1amu & 2amu (b) 2 amu & 4 amu
 (c) 3amu & 6amu (d) 4 amu & 8 amu
- One ‘u’ stands for the
 (a) mass of an atom of carbon-12 atom.
 (b) $1/12^{\text{th}}$ of mass of carbon-12.
 (c) $1/12^{\text{th}}$ of mass of hydrogen atom.
 (d) mass of one atom of any of the element.
- Mass of 1 amu in g is equal to:
 (a) 1.66×10^{24} g (b) 1.66×10^{-24} g
 (c) 1.008 g (d) 9.1×10^{-28} g

MOLE CONCEPT

Mole

Mole is a counting unit mostly used for microscopic particles and is defined as follows:

A mole is the amount of a substance that contains as many entities (atoms, molecules or other particles) as there are atoms in exactly 0.012 kg (or 12 gm) of the carbon-12 isotope.

From mass spectrometer we found that there are 6.022×10^{23} atoms present in 12 gm of C^{12} isotope.

The number of entities in 1 mol is so important that it is given a separate name and symbol known as Avogadro constant denoted by N_A . i.e., on the whole we can say that 1 mole is the collection of 6.022×10^{23} entities. Here entities may represent atoms, ions, molecules or even pens, chair, paper etc but as this number (N_A) is very large therefore it is significant only for very micro-particles.

Illustration: Chlorophyll, the green colouring material of plants contains 3.68 % of magnesium by mass. Calculate the number of magnesium atom in 5.00 g of the complex.

Sol. Mass of magnesium in 5.0 g of complex

$$= \frac{3.68}{100} \times 5.00 = 0.184 \text{ g}$$

Atomic mass of magnesium = 24

24 g of magnesium contain = 6.022×10^{23} atoms

$$0.184 \text{ g of magnesium would contain} = \frac{6.022 \times 10^{23}}{24} \times 0.184$$

$$= 4.617 \times 10^{21}$$

Therefore, 5.00 g of the given complex would contain 4.617×10^{21} atoms of magnesium.

How Big is a Mole?

Amount of water in world's oceans (litres)	Age of earth (seconds)
Population of earth	
Avogadro's number	602,200,000,000,000,000,000
Distance from earth to sun (centimeters)	

❖ In modern practice gram-atom and gram-molecule are termed as mole.

Gram Atomic Mass: The atomic mass of an element expressed in gram is called gram atomic mass of the element.

or

It is also defined as mass of 6.022×10^{23} atoms.

or

It is also defined as the mass of one mole atoms.

Eg:

Element	R.A.M. (Relative Atomic Mass)	Atomic mass (mass of one atom)	Gram Atomic mass/weight
N	14	14 amu	14 gm
He	4	4 amu	4 gm
C	12	12 amu	12 gm

For example, for oxygen atom: Atomic mass of ‘O’ atom = mass of one ‘O’ atom = 16 amu

$$\text{Gram atomic mass} = \text{mass of } 6.022 \times 10^{23} \text{ ‘O’ atoms}$$

$$= 16 \text{ amu} \times 6.022 \times 10^{23}$$

$$= 16 \times 1.66 \times 10^{-24} \text{ g} \times 6.022 \times 10^{23} = 16 \text{ g}$$

(∵ $1.66 \times 10^{-24} \times 6.022 \times 10^{23} \simeq 1$)

Gram Molecular Mass: The molecular mass of a compound expressed in gram is called the gram-molecular mass of the compound.

or

It is also defined as mass of 6.022×10^{23} molecules.

or

It is also defined as the mass of one mole molecules.

For example, for ‘O₂’ molecule: Molecular mass of ‘O₂’ molecule
 = mass of one ‘O₂’ molecule
 = 2 × mass of one ‘O’ atom = 2 × 16 amu = 32 amu
 Gram molecular mass = mass of 6.022×10^{23} ‘O₂’ molecules
 = 32 amu × 6.022×10^{23}
 = $32 \times 1.66 \times 10^{-24} \text{ gm} \times 6.022 \times 10^{23} = 32 \text{ gm}$

RELATIONSHIP BETWEEN GRAM AND AMU

$$1 \text{ amu} = \frac{1}{12} \text{ of wt. of one } \text{C}^{12} \text{ atom.}$$

For carbon (C^{12}) atom, 1 mole C = 12 gm = 6.022×10^{23} atoms
 wt. of 6.022×10^{23} C^{12} atoms = 12 gm

$$\text{wt. of 1 atom of } \text{C}^{12} = \frac{12}{N_A} \text{ gm}$$

($N_A \rightarrow$ Avogadro's number = 6.022×10^{23})

$$1 \text{ amu} = \frac{1}{12} \text{ of wt. of one } C^{12} \text{ atom} = \frac{1}{12} \times \frac{12}{N_A} \text{ gm}$$

$$1 \text{ amu} = \frac{1}{N_A} \text{ gm}$$

METHODS OF CALCULATIONS OF MOLE

(a) If no. of particles of some species is given, then no. of moles

$$= \frac{\text{Given no. of particles}}{N_A}$$


(b) If weight of a given species is given, then no. of moles =

$$\frac{\text{Given wt.}}{\text{Atomic wt.}} \text{ (for atoms),}$$

 or
$$\frac{\text{Given wt.}}{\text{Molecular wt.}} \text{ (for molecules)}$$

(c) If volume of a gas is given along with its temperature (T) and pressure (P) use $n = \frac{PV}{RT}$ (assuming gas to be ideal) where $R = 0.0821 \text{ lit-atm/mol-K}$ (when P is in atmosphere and V is in litre.)

Y-map: Interconversion of mole - volume, mass and number of particles:



ATOMICITY

It is equal to number of atoms present in one molecule. For example atomicity of H_2 , CO_2 , O_3 , CCl_4 , respectively.

DENSITY

For Liquids and Solids,

(a) Absolute density

$$\text{Absolute density} = \frac{\text{Mass}}{\text{Volume}}$$

(b) Relative density

$$\text{Relative density} = \frac{\text{Density of substance}}{\text{Density of standard substance}}$$

$$\text{e.g., Specific gravity} = \frac{\text{Density of substance}}{\text{Density of } H_2O \text{ at } 4^\circ C}$$

For Gases:

$$\text{Absolute density (mass / volume)} = \frac{PM}{RT}$$

where P is pressure of gas, M = mol. wt. of gas, R is the gas constant, T is the temperature.

Vapour density: It is defined only for gas.

It is a density of gas with respect to H_2 gas at same temperature & pressure.

$$V.D. = \frac{d_{\text{gas}}}{d_{H_2}} = \frac{PM_{\text{gas}} / RT}{PM_{H_2} / RT} = \frac{M_{\text{gas}}}{M_{H_2}} = \frac{M}{2} \quad \boxed{V.D. = \frac{M}{2}}$$

Key Note

- 1 g O , 1 g O_2 , 1 g O_3 each have same number of oxygen atoms.
- Density of liquid water at $4^\circ C$ is 1 g/mL = 1 g/cc = 10^3 kg/m^3

Train Your Brain

Example 6: Total number of atoms of all elements present in 1 mole of ammonium dichromate is?

- (a) 14 (b) 19
 (c) 6×10^{23} (d) 114×10^{23}

Sol. (d) $(NH_4)_2Cr_2O_7 = 19 \times 6.022 \times 10^{23} \approx 114 \times 10^{23}$ atoms.

Example 7: How many atoms of oxygen are there in 16 g of oxygen?

- (a) $2 N_A$ (b) N_A (c) $1.5 N_A$ (d) $4 N_A$

Sol. (b) Let x atoms of oxygen are present

$$\text{So, } 16 \times 1.66 \times 10^{-24} \times x = 16 \text{ g} \Rightarrow x = \frac{1}{1.66 \times 10^{-24}} = N_A$$

Example 8: Calculate the volume in litres of 20 g hydrogen gas at STP.

- (a) 2.24 L (b) 22.4 L
 (c) 224 L (d) 4.48 L

$$\begin{aligned} \text{Sol. (b) No. of moles of hydrogen gas} &= \frac{\text{Mass}}{\text{Molecular mass}} \\ &= \frac{20 \text{ gm}}{2 \text{ gm}} = 10 \text{ mol} \end{aligned}$$

$$\text{Volume of hydrogen gas at STP} = 10 \times 22.4 \text{ L}$$

Example 9: The number of atoms contained in 11.2 L of SO_2 at S.T.P. are:

- (a) $3/2 \times 6.022 \times 10^{23}$ (b) $2 \times 6.022 \times 10^{23}$
 (c) 6.022×10^{23} (d) $4 \times 6.022 \times 10^{23}$

Sol. (a) 22.4 litre gas has = 1 mole

$$\begin{aligned} 1 \text{ litre gas has} &= \frac{1}{22.4} \times 11.2 = \frac{1}{2} \text{ mole of molecules} \\ &= 3 \times \frac{1}{2} \text{ mole of atoms} = \frac{3}{2} \times 6.022 \times 10^{23} \text{ atoms} \end{aligned}$$

Example 10: 7.5 litre of the particular gas at S.T.P. weighs 16 gram. What is the V.D. of gas?

$$\text{Sol. Moles at S.T.P.} = \frac{7.5}{22.4} = \frac{16}{M}$$

$$M = 48 \text{ gram; V.D.} = \frac{48}{2} = 24.$$

Example 11: Find the vapour density of $CO_2(g)$ with respect to $N_2O(g)$.

$$\text{Sol. V.D.} = \frac{\text{M.wt. of } CO_2}{\text{M.wt. of } N_2O} = \frac{44}{44} = 1$$

Example 12: Find the vapour density of N_2O_5 .

$$\text{Sol. V.D.} = \frac{\text{Mol. wt. of } N_2O_5}{2} = \frac{108}{2} = 54.$$

Some Basic Concepts of Chemistry

Concept Application

- The number of electrons present in 1 mol of methane molecule are:
 (a) 6.022×10^{25} (b) 6.022×10^{24}
 (c) 6.022×10^{23} (d) 6.022×10^{22}
- The mass of one molecule of water is approximately:
 (a) 3×10^{-23} g (b) 18 g
 (c) 1.5×10^{-23} g (d) 4.5×10^{-23} g
- The molar mass of ferrous sulphate ($\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$) is:
 (a) 152 gm (b) 278 gm
 (c) 137 gm (d) None of these
- The vapour density of carbon dioxide is:
 (a) 44 (b) 32 (c) 22 (d) 12
- The density of air is 0.001293 g/cm^3 at STP. Identify which of the following statement is correct?
 (a) Vapour density is 12.72.
 (b) Molecular weight is 28.96.
 (c) Vapour density is 0.001293 g/cm^3 .
 (d) Vapour density and molecular weight cannot be determined.

ELEMENTAL ANALYSIS

For n mole of a compound ($\text{C}_3\text{H}_7\text{O}_2$);

Moles of C = 3n
 Moles of H = 7n
 Moles of O = 2n

PERCENTAGE FORMULAE COMPOSITION

% of element in a compound

$$= \frac{\text{Atomic weight of element} \times \text{Number of atom of that element in one molecule} \times 100}{\text{Total molecular weight of compound}}$$

Here we are going to find out the percentage of each element in the compound by knowing the molecular formula of compound.

EMPIRICAL AND MOLECULAR FORMULA

We have just seen that knowing the molecular formula of the compound, we can calculate percentage composition of the elements. Conversely, if we know the percentage composition of the elements initially, we can calculate the relative number of atoms of each element in the molecules of the compound. This gives us the empirical formula of the compound. Further if the molecular mass is known then the molecular formula can easily be determined.

An empirical formula represents the simplest whole number ratio of various atoms present in a compound.

The molecular formula gives the actual number of atoms of each element in a molecule. The molecular formula shows the exact number of different types of atoms present in a molecule of a compound.

The molecular formula is an integral multiple of the empirical formula.

i.e. Molecular formula = Empirical formula $\times$ n

$$\text{where } n = \frac{\text{Molecular Formula Mass}}{\text{Empirical Formula Mass}}$$

❖ If sum of mass percent of all elements is less than 100 then difference is due to oxygen.

$$\text{❖ Mass \% of C} = \frac{12}{44} \times \frac{m_{\text{CO}_2}}{m_{\text{compound}}} \times 100$$

$$\text{❖ Mass \% of H} = \frac{2}{18} \times \frac{m_{\text{H}_2\text{O}}}{m_{\text{compound}}} \times 100$$

Train Your Brain

Example 13: Every molecule of ammonia always has formula NH_3 irrespective of method of preparation or sources. i.e. 1 mole of ammonia always contains 1 mol of N and 3 mole of H. In other words 17 gm of NH_3 always contains 14 gm of N and 3 gm of H. Now find out % of each element in the compound.

Sol. Mass % of N in NH_3 =

$$\frac{\text{Mass of N in 1 mole } \text{NH}_3}{\text{Mass of 1 mole of } \text{NH}_3} \times 100 = \frac{14}{17} \times 100 = 82.35\%$$

Mass % of H in NH_3 =

$$\begin{aligned} & \frac{3 \times \text{Mass of H in 1 mole } \text{NH}_3}{\text{Mass of 1 mole of } \text{NH}_3} \times 100 \\ &= \frac{3}{17} \times 100 = 17.65\% \end{aligned}$$

Example 14: Acetylene and benzene both have the empirical formula CH. The molecular masses of acetylene and benzene are 26 and 78 respectively. Deduce their molecular formulae.

Sol. $\therefore$ Empirical Formula is CH

Step-1: The empirical formula of the compound is CH

$\therefore$ Empirical formula mass = $(1 \times 12) + 1 = 13$.

Molecular mass = 26

Step-2: To calculate the value of 'n'

$$n = \frac{\text{Molecular mass}}{\text{Empirical formula mass}} = \frac{26}{13} = 2$$

Step-3: To calculate the molecular formula of the Compound.

Molecular formula = $n \times$ (Empirical formula of the compound) = $2 \times \text{CH} = \text{C}_2\text{H}_2$

Thus the molecular formula is C_2H_2

Similarly for benzene

To calculate the value of 'n'

$$n = \frac{\text{Molecular mass}}{\text{Empirical formula mass}} = \frac{78}{13} = 6.$$

Thus the molecular formula is $6 \times \text{CH} = \text{C}_6\text{H}_6$

Example 15: An organic substance containing carbon, hydrogen and oxygen gave the following percentage composition.

C = 40.684% ; H = 5.085% and O = 54.228%

The molecular weight of the compound is 118. Calculate the molecular formula of the compound.

- (a) C₄H₆O₄ (b) C₂H₃O₂
(c) C₂H₄O₂ (d) C₄H₆O₈

Sol. (a) Step-1: To calculate the empirical formula of the compound.

Element	Symbol	Percentage of element	At. mass of element	Relative no. of atoms = $\frac{\text{Percentage}}{\text{At. mass}}$	Simplest atomic ratio	Simplest whole no. atomic ratio
Carbon	C	40.684	12	$\frac{40.684}{12} = 3.390$	$\frac{3.390}{3.389} = 1$	2
Hydrogen	H	5.085	1	$\frac{5.085}{1} = 5.085$	$\frac{5.085}{3.389} = 1.5$	3
Oxygen	O	54.228	16	$\frac{54.228}{16} = 3.389$	$\frac{3.389}{3.389} = 1$	2

∴ Empirical Formula is C₂ H₃ O₂

Step-2: To calculate the empirical formula mass.

The empirical formula mass of the compound is 12 × 2 + 3 × 1 + 16 × 2 = 59.

Step-3: To calculate the value of ‘n’

$$n = \frac{\text{Molecular mass}}{\text{Empirical formula mass}} = \frac{118}{59} = 2$$

Step-4: To calculate the molecular formula of the salt

Molecular formula = n × (Empirical formula)

$$= 2 \times \text{C}_2\text{H}_3\text{O}_2 = \text{C}_4\text{H}_6\text{O}_4$$

Thus the molecular formula is C₄ H₆ O₄.

Example 16: Acetylene & butene have empirical formula CH & CH₂ respectively. The molecular mass of acetylene and butene are 26 & 56 respectively. Deduce their molecular formula.

Sol.

$$n = \frac{\text{Molecular mass}}{\text{Empirical formula mass}}$$

$$\text{For Acetylene : } n = \frac{26}{13} = 2$$

$$\therefore \text{Molecular formula} = \text{C}_2\text{H}_2$$

$$\text{For Butene: } n = \frac{56}{14} = 4$$

$$\therefore \text{Molecular formula} = \text{C}_4\text{H}_8$$

Example 17: An oxide of nitrogen gave the following percentage composition by mass:

N = 25.94 and O = 74.06

Calculate the empirical formula of the compound.

- (a) NO₂ (b) N₂O₄ (c) N₂O₅ (d) N₂O

Sol. (c)

Element	%/Atomic mass	divide by smallest ratio	Simple integer ratio
N	$\frac{25.94}{14} = 1.85$	$\frac{1.85}{1.85} = 1$	2
O	$\frac{74.06}{16} = 4.63$	$\frac{4.63}{1.85} = 2.5$	5

So empirical formula is N₂O₅.

Concept Application

15. A compound contains 25% hydrogen and 75% carbon by mass. Determine the empirical formula of the compound.

- (a) CH₄ (b) C₂H₆
(c) C₃H₈ (d) C₂H₂

16. The empirical formula of a compound of molecular mass 120 u is CH₂O. The molecular formula of the compound is:

- (a) C₂H₄O₂ (b) C₄H₈O₄
(c) C₃H₅C₃ (d) All of these

17. Calculate the molecular formula of compound which contains 20% Ca and 80% Br (by wt.) if molecular weight of compound is 200 u. (Atomic wt. Ca = 40, Br = 80)

- (a) Ca_{1/2}Br (b) CaBr₂
(c) CaBr (d) Ca₂Br

CONCENTRATION OF SOLUTION

Concentration of a solution can be expressed in any of the following ways.

(i) **% by wt.:** Amount of solute (in g) dissolved in 100 gm of solution.

4.9% H₂SO₄ by wt.

⇒ 100 gm of solution contains 4.9 gm of H₂SO₄.

(ii) **% by volume:** Volume of solute (in ml) dissolved in 100 ml of solution.

x% H₂SO₄ by volume

⇒ 100 ml of solution contains x ml H₂SO₄.

(iii) **% wt. by volume:** wt. of solute (in g) present in 100 ml of solution.

CONCENTRATION TERMS

Molarity (M): No. of moles of solute present in 1000 ml of solution.

$$\text{Molarity (M)} = \frac{\text{Moles of solute}}{\text{Volume of solution (L)}}$$

Molality (m): No. of moles of solute present in 1000 gm of solvent.

$$m = \frac{\text{Moles of solute}}{\text{wt. of solvent in kg}}$$

Mole fraction: The mole fraction of a particular component in a solution is defined as the number of moles of that component per mole of solution.

If a solution has n_A mole of A & n_B mole of B.

mole fraction of A (X_A) = $\frac{n_A}{n_A + n_B}$

mole fraction of B (X_B) = $\frac{n_B}{n_A + n_B}$
 $X_A + X_B = 1$

Parts per million (ppm):

$$= \frac{\text{Mass of solute}}{\text{Mass of solvent}} \times 10^6 \cong \frac{\text{Mass of solute}}{\text{Mass of solution}} \times 10^6$$

CONVERSION OF CONCENTRATION TERMS

1. Molarity and % solute by mass: Let d = density of solution in g/mL and let it contains x% (w/w) solute by mass.

$$M = \frac{x \times d \times 10}{m_A}$$

❖ Specific gravity has no units and its numerical value equals density in g/mL.

2. Molality and mole fraction: Consider a binary solution consisting of two components A (Solute) and B (Solvent). Let X_A & X_B are the mole fraction of A & B respectively.

$$X_A = \frac{n_A}{n_A + n_B}, \quad X_B = \frac{n_B}{n_A + n_B}$$

If molality of solution be m then:

$$m = \frac{n_A}{\text{mass of solvent}} \times 1000 = \frac{n_A}{n_B \times M_B} \times 1000$$

where M_B is the molecular wt. of the solvent B.

$$m = \frac{X_A}{X_B} \times \frac{1000}{M_B} \Rightarrow m = \frac{\text{mole fraction of A}}{\text{mole fraction of B}} \times \frac{1000}{M_B}$$

$$m = \frac{\text{mole fraction of solute}}{\text{mole fraction of solvent}} \times \frac{1000}{\text{molecular wt. of solvent}}$$

3. Mole fraction of solute into molarity of solution

$$M = \frac{X_2 \times d \times 1000}{X_1 M_1 + X_2 M_2}$$

Mole fraction of solvent and solute are X_1 and X_2 so $X_1 + X_2 = 1$

Suppose total mole of solution is = 1 then mole of solute and solvent are X_2 & X_1 respectively

weight of solute = $X_2 M_2$, weight of solvent = $X_1 M_1$

& total wt of solution = $X_1 M_1 + X_2 M_2$

volume of solution = $\frac{X_1 M_1 + X_2 M_2}{d}$ ml = $\frac{X_1 M_1 + X_2 M_2}{d \times 1000}$ L

molarity (M) = $\frac{X_2 \times d \times 1000}{X_1 M_1 + X_2 M_2}$

4. Molarity into mole fraction $X_2 = \frac{1000M}{1000d - MM_2}$

Molarity = M moles solute in 1000 ml of solution

So, moles of solute = M & mass of solution = $d \times 1000$

wt. of solute = MM_2 & wt. of solvent = $1000d - MM_2$

Where M_2 is molar mass of solute

mole fraction of solute = $\frac{1000M}{1000d - MM_2}$

5. Molality into mole fraction $X_2 = \frac{mM_1}{1000 + mM_1}$

Molality = moles of solute in 1000 gm of solvent = m

moles of solvent = $\frac{1000}{M_1}$ where M_1 is molar mass of solvent

mole fraction $X_2 = \frac{m}{\frac{1000}{M_1} + m} = \frac{mM_1}{1000 + mM_1}$

6. Molality into molarity $M = \frac{md \times 1000}{1000 + mM_2}$

Molality = m moles of solute in 1000 gm of solvent

mole of solute = m & weight of solute = mM_2

Weight of solution = $1000 + mM_2$

volume of solution = $\frac{1000 + mM_2}{d}$ mL = $\frac{1000 + mM_2}{d \times 1000}$ L

molarity = $\frac{m \times d \times 1000}{1000 + mM_2}$

7. Molarity into Molality $m = \frac{M \times 1000}{1000d - MM_2}$

M_1 and M_2 are molar masses of solvent and solute.

Molarity = M mole of solute in 1000 ml of solution

mole of solute = M & weight of solute = MM_2

weight of solvent = $1000d - MM_2$

mass of solvent = $1000d - MM_2$

molality = $\frac{M \times 1000}{1000d - MM_2}$

on simplifying $d = M \left[\frac{1}{m} + \frac{M_2}{1000} \right]$

DILUTION & MIXING OF TWO LIQUIDS

❖ Upon dilution no. of moles of solute remains constant. If a particular solution having volume V_1 mL and molarity M_1 is diluted upto volume V_2 mL.

$$M_1 V_1 = M_2 V_2$$

M_2 : final molarity

❖ If a solution having volume V_1 and molarity M_1 is mixed with another solution of same solute having volume V_2 & molarity M_2 then $M_1 V_1 + M_2 V_2 = MR (V_1 + V_2)$

MR = Resultant molarity = $\frac{M_1 V_1 + M_2 V_2}{V_1 + V_2}$

Key Note

○ Molality is the most convenient method to express the concentration because it involves the mass of liquids rather than their volumes. It is also independent of the variation in temperature.

- No. of moles of solute = Molarity $\times$ volume of solution (in L)
- No. of millimoles of solute = Molarity $\times$ volume of solution (in mL)
- 1 mole = 1000 millimole
- All those concentration terms which does not involve volume terms e.g., ppm, mass %, molality, mole fraction are independent of temperature of the solution.
- For dilue aqueous solution, molarity = molality
- Concentration of solids or pure liquids is constant. ($C = n/V = d/M$)

Train Your Brain

Example 18: 0.2 mole of HCl and 0.1 mole of barium chloride were dissolved in 0.5 L of water to produce solution. The molarity of the Cl^- ions is:

- (a) 0.06 M (b) 0.09 M
(c) 0.12 M (d) 0.80 M

Sol. (d) $\text{HCl} \longrightarrow \text{Cl}^-$

0.2 mole

$\text{BaCl}_2 \longrightarrow 2 \text{Cl}^-$

$$2 \times 0.1 = 0.2$$

$$\text{Total moles of } \text{Cl}^- = 0.4 \Rightarrow M = \frac{w \times 1000}{m \times v}$$

$$\text{Molarity} = \frac{0.4 \times 1000}{500} = 0.8 \therefore \left(\frac{w}{1} = 0.4 \right)$$

Example 19: 149 g of potassium chloride (KCl) is dissolved in 10 L of an aqueous solution. Determine the molarity of the solution. (K = 39, Cl = 35.5)

- (a) 0.2 M (b) 0.4 M (c) 0.5 M (d) 2.2 M

Sol. (a) Molecular mass of KCl = 39 + 35.5 = 74.5 g

$$\therefore \text{Moles of KCl} = \frac{149 \text{ g}}{74.5 \text{ g}} = 2$$

$$\therefore \text{Molarity of the solution} = \frac{2}{10} = 0.2 \text{ M}$$

Example 20: 255 g of an aqueous solution contains 5 g of urea. What is the concentration of the solution in terms of molality? (Mol. wt. of urea = 60)

- (a) 0.222 m (b) 0.333 m
(c) 2.22 m (d) 3.33 m

Sol. (b) Mass of urea = 5 g

Molecular mass of urea = 60

$$\text{Number of moles of urea} = \frac{5}{60} = 0.083$$

Mass of solvent = (255 – 5) = 250 g

$\therefore$ Molality of the solution

$$= \frac{\text{Number of moles of solute}}{\text{Mass of solvent in gram}} \times 1000$$

$$= \frac{0.083}{250} \times 1000 = 0.332 \text{ m}$$

Example 21: 117 g NaCl is dissolved in 500 ml aqueous solution. Find the molarity of the solution.

$$\text{Sol. Molarity} = \frac{117 / 58.5}{500 / 1000} = 4 \text{ M.}$$

Example 22: Calculate the resultant molarity of following:

- (a) 200 ml 1M HCl + 300 ml water
(b) 1500 ml 1M HCl + 18.25 g HCl
(c) 200 ml 1M HCl + 100 ml 0.5 M H_2SO_4
(d) 200 ml 1M HCl + 100 ml 0.5 M HCl

$$\text{Sol. (a) Final molarity} = \frac{200 \times 1 + 0}{200 + 300} = 0.4 \text{ M.}$$

$$(b) \text{ Final molarity} = \frac{1500 \times 1 + \frac{18.25 \times 1000}{36.5}}{1500} = 1.33 \text{ M}$$

$$(c) \text{ Final molarity of } \text{H}^+ = \frac{200 \times 1 + 100 \times 0.5 \times 2}{200 + 100} = 1 \text{ M}$$

$$(d) \text{ Final molarity} = \frac{200 \times 1 + 100 \times 0.5}{200 + 100} = 0.83 \text{ M.}$$

Example 23: 518 g of an aqueous solution contains 18 g of glucose (mol.wt. = 180). What is the molality of the solution?

Sol. wt. of solvent = 518 – 18 = 500 g.

$$\Rightarrow \text{So molality} = \frac{18 / 180}{500 / 1000} = 0.2 \text{ M}$$

Example 24: 0.25 g of a substance is dissolved in 6.25 g of a solvent. Calculate the percentage amount of the substance in the solution.

Sol. Wt. of solution = 0.25 + 6.25 = 6.50

$$\text{So \% (w/w)} = \frac{0.25}{6.50} \times 100 = 3.8\%$$

Example 25: An aqueous solution is 1.33 molal in methanol. Determine the mole fraction of methanol & H_2O .

Sol. Molality

$$= \frac{\text{Mole fraction of solute}}{\text{Mole fraction of solvent} \times \text{mol.wt. of solvent}} \times 1000$$

$$1.33 = \frac{X_A}{X_B \times M_B} \times 1000; \frac{1.33 \times 18}{1000} = \frac{X_A}{X_B}; \frac{23.94}{1000} = \frac{X_A}{X_B}$$

$$\Rightarrow X_A = 0.02394 X_B, X_A + X_B = 1 \Rightarrow 1.02394 X_B = 1$$

$$X_B = \frac{1}{1.02394} = 0.98, X_A = 0.02$$

2nd Method: Let wt. of solvent = 1000 gm, molality = 1.33 means 1.33 moles of solute are present in 1000 g solvent

$$\begin{aligned} \text{mole fraction of solute} &= \frac{\text{moles of solute}}{\text{moles of solute} + \text{moles of solvent}} \\ &= \frac{m}{m + \frac{1000}{18}} = \frac{1.33}{1.33 + (1000/18)} \end{aligned}$$

Mole fraction of solute = 0.02
Mole fraction of solvent = 1 – 0.02 = 0.98

Concept Application


18. If 500 ml of 1 M solution of glucose is mixed with 500 ml of 1 M solution of glucose, final molarity of solution will be:
(a) 1 M (b) 0.5 M (c) 2 M (d) 1.5 M
19. The volume of water that must be added to a mixture of 250 ml of 0.6 M HCl and 750 ml of 0.2 M HCl to obtain 0.25 M solution of HCl is:
(a) 750 ml (b) 100 ml (c) 200 ml (d) 300ml
20. The molarity of Cl^- in an aqueous solution which was (w/v) 2% NaCl, 4% CaCl_2 and 6% NH_4Cl will be:
(a) 0.342 (b) 0.721 (c) 1.12 (d) 2.18
21. 2M of 100 ml Na_2SO_4 is mixed with 3M of 100 ml NaCl solution and 1 M of 200 ml CaCl_2 solution. Then the ratio of the concentration of Na^+ ion and Cl^- ion.
(a) 1/2 (b) 2 (c) 1 (d) 4
22. Equal moles of H_2O and NaCl are present in a solution. Hence, molality of NaCl solution is :
(a) 0.55 (b) 55.5 (c) 1.00 (d) 0.18
23. Mole fraction of A in H_2O is 0.2. The molality of A in H_2O is:
(a) 13.9 (b) 15.5 (c) 14.5 (d) 16.8
24. What is the molarity of H_2SO_4 solution that has a density of 1.84 g/cc and contains 98% by mass of H_2SO_4 ? (Given atomic mass of S = 32)
(a) 4.18 M (b) 8.14M (c) 18.4 M (d) 18 M
25. The molarity of the solution containing 2.8% (mass/volume) solution of KOH is : (Given atomic mass of K = 39) is :
(a) 0.1 M (b) 0.5 M (c) 0.2 M (d) 1 M

STOICHIOMETRY BASED CONCEPT (PROBLEMS BASED ON CHEMICAL REACTION)

All chemical reaction are represented by chemical equations by using chemical formulae of reactants and products. Qualitatively a chemical equation simply describes what the reactants and products are. However, a balanced chemical equation gives us a lot of quantitative information mainly the molar ratio in which reactants combine and the molar ratio in which products are formed.

Example:

When potassium chlorate (KClO_3) is heated it gives potassium chloride (KCl) and oxygen (O_2).



Attributes of a balanced chemical equation:

- (a) It contains an equal number of atoms of each element on both sides of equation.
- (b) It should follow law of charge conservation on either side.
- (c) Physical states of all the reagents should be included in brackets.
- (d) All reagents should be written in their standard molecular forms (not as atoms)
- (e) The coefficients give the relative molar ratios of each reagent.

Balancing a Chemical Equation

Many chemical equations can be balanced by trial and error. Let us take the reactions of a few metals and non-metals with oxygen to give oxides

For Example:

Combustion reaction of C_2H_6 : $\text{C}_2\text{H}_6 + \text{O}_2 \rightarrow \text{CO}_2 + \text{H}_2\text{O}$
(skeleton equation)

Balance carbon atoms: $\text{C}_2\text{H}_6 + \text{O}_2 \rightarrow 2\text{CO}_2 + \text{H}_2\text{O}$

Now balance hydrogen atoms: $\text{C}_2\text{H}_6 + \text{O}_2 \rightarrow 2\text{CO}_2 + 3\text{H}_2\text{O}$

Now balance oxygen atoms: $\text{C}_2\text{H}_6 + \frac{7}{2}\text{O}_2 \rightarrow 2\text{CO}_2 + 3\text{H}_2\text{O}$


Always remember that subscripts in formula of reactants and products cannot be changed to balance an equation.

One of the most important aspects of a chemical equation is that when it is written in the balanced form, it gives quantitative relationships between the various reactants and products in terms of moles, masses, molecules and volumes.

Mole - Mole Analysis

This analysis is very much important for quantitative analysis point of view.

Consider the decomposition of KClO_3 .



In very first step of mole-mole analysis you should read the balanced chemical equation like **2 moles KClO_3 on decomposition gives you 2 moles KCl and 3 moles O_2** and from the stoichiometry of reaction we can write

$$\frac{\text{Moles of KClO}_3}{2} = \frac{\text{Moles of KCl}}{2} = \frac{\text{Moles of O}_2}{3}$$


Now for any general balanced chemical equation like



you can write.

$$\begin{aligned} \frac{\text{Moles of A reacted}}{a} &= \frac{\text{Moles of B reacted}}{b} \\ &= \frac{\text{Moles of C produced}}{c} = \frac{\text{Moles of D produced}}{d} \end{aligned}$$


Further, a balanced chemical equation along with the quantitative information conveyed by it is given below:



Thus,

- 1 mole of calcium carbonate reacts with 2 moles of hydrochloric acid to give 1 mole of calcium chloride, 1 mole of water and 1 mole of carbon dioxide.
- 100 g of calcium carbonate react with 73 g hydrochloric acid to give 111 g of calcium chloride, 18 g of water and 44 g (or 22.4 litres at STP) of carbon dioxide.

Stoichiometry:



(law of conservation of mass is followed).

- Mass can not be represented by stoichiometry.
- The quantitative information conveyed by a chemical equation helps in a number of calculations. The problems involving these calculations may be classified into the following different types:

Type (I) Mass - Mass Relationships i.e. mass of one of the reactants or products is given and the mass of some other reactant or product is to be calculated.

Mass - Mass Analysis

Consider the reaction $2 \text{KClO}_3 \rightarrow 2\text{KCl} + 3\text{O}_2$

According to stoichiometry of the reaction

mass-mass ratio = $2 \times 122.5 : 2 \times 74.5 : 3 \times 32$

$$\text{or } \frac{\text{Mass of KClO}_3}{\text{Mass of KCl}} = \frac{2 \times 122.5}{2 \times 74.5}$$

$$\frac{\text{Mass of KClO}_3}{\text{Mass of O}_2} = \frac{2 \times 122.5}{3 \times 32}$$

Illustration: Calculate the weight of iron which will be converted into its oxide by the action of 36 g of steam.



Sol. Mole ratio of reaction suggests,


$$\frac{\text{Mole of Fe}}{\text{Mole of H}_2\text{O}} = \frac{3}{4}$$

$$\therefore \text{Mole of Fe} = \frac{3}{4} \times \text{mol of H}_2\text{O} = \frac{3}{4} \times \frac{36}{18} = \frac{3}{2}$$

$$\text{wt. of Fe} = \frac{3}{2} \times 56 = 84 \text{ g}$$

Illustration: What amount of silver chloride is formed by the action of 5.850 g of sodium chloride on an excess of silver nitrate?

Sol. Writing the balanced equation for the reaction



$$n_{\text{NaCl}} = \frac{\text{Weight}}{M_w} = \frac{5.85}{58.5} = 0.1 \text{ mol}$$


$$\frac{\text{mole}(\text{NaCl})}{\text{mole}(\text{AgCl})} = \frac{1}{1}$$

$$\Rightarrow \text{mole}(\text{AgCl}) = 0.1 = \frac{\text{Weight}}{M_w} = \frac{\text{Weight}}{143.5}$$

$$\Rightarrow \text{weight} = 0.1 \times 143.5 \text{ g} = 14.35 \text{ g.}$$

Illustration: How much iron can be theoretically obtained in the reduction of 1 kg of Fe_2O_3 ?

Sol. Writing the balanced equation for the decomposition



$$n_{\text{Fe}_2\text{O}_3} = \frac{\text{Weight}}{M_w} = \frac{1000}{160} \text{ mol}$$


$$\frac{\text{mole}(\text{Fe}_2\text{O}_3)}{\text{mole}(\text{Fe})} = \frac{1}{2}$$

$$\Rightarrow \text{moles of Fe} = \frac{2 \times 1000}{160} = 12.5 \text{ mol}$$

$$= \frac{\text{Weight}}{\text{Atomic weight}} = \frac{\text{Weight}}{56}$$

$$\text{Weight of iron obtained} = 12.5 \times 56 \text{ g} = 700 \text{ g}$$

Type (II) Mass - Volume Relationships i.e. mass/volume of one of the reactants or products is given and the volume/mass of the other is to be calculated.



Mass volume ratio =


$2 \times 122.5 \text{ g} : 2 \times 74.5 \text{ g} : 3 \times 22.4 \text{ L at STP}$

we can use two relation for volume of oxygen

$$\frac{\text{Mass of KClO}_3}{\text{Volume of O}_2 \text{ at STP}} = \frac{2 \times 122.5 \text{ g}}{3 \times 22.4 \text{ L}} \quad \dots(\text{i})$$

$$\text{and } \frac{\text{Mass of KCl}}{\text{Volume of O}_2 \text{ at STP}} = \frac{2 \times 74.5 \text{ g}}{3 \times 22.4 \text{ L}} \quad \dots(\text{ii})$$

Illustration: How much marble of 90.5% purity would be required to prepare 10 litres of CO_2 at STP when the marble is acted upon by dilute HCl?



100 g 22.4 litre

22.4 L of CO_2 at STP will be obtained from 100 g of CaCO_3

$\therefore$ 10 L of CO_2 at STP will be obtained from pure


$$\text{CaCO}_3 = \frac{100}{22.4} \times 10 = 44.64 \text{ g}$$

$$\therefore \text{Impure marble required} = \frac{100}{90.5} \times 44.64 = 49.326 \text{ g}$$



Concept Application

26. If 1.5 moles of dioxygen combine with Al to form Al_2O_3 , the weight of Al used in the reaction is:
 (a) 27 g (b) 40.5 g (c) 54 g (d) 81 g
27. How many litres of CO_2 at STP will be formed when 0.01 mol of H_2SO_4 reacts with excess of Na_2CO_3 ?
 $\text{Na}_2\text{CO}_3 + \text{H}_2\text{SO}_4 \longrightarrow \text{Na}_2\text{SO}_4 + \text{CO}_2 + \text{H}_2\text{O}$
 (a) 22.4 L (b) 2.24 L (c) 0.224 L (d) 1.12 L
28. How many moles of potassium chlorate need to be heated to produce 11.2 litre oxygen at S.T.P.?



- (a) $\frac{1}{2}$ mol (b) $\frac{1}{3}$ mol
 (c) $\frac{1}{4}$ mol (d) $\frac{2}{3}$ mol


LIMITING REAGENT (L.R.) CONCEPT

Quite often one of the reactants is present in larger amount than the other as required according to the balanced equation. The amount of the product formed then depends upon the reactant which has reacted completely. This reactant is called the limiting reactant. The excess of the other is left unreacted.

Limiting Reagent (L.R.): The reactant which is completely consumed in a reaction is called as L.R.

Calculation of Limiting Reagent:

- (a) By calculating the required amount by the equation and comparing it with given amount. [Useful when only two reactant are there]
- (b) By calculating amount of any one product obtained taking each reactant one by one irrespective of other reactants. The one giving least product is limiting reagent.
- (c) Divide given moles of each reactant by their stoichiometric coefficient, the one with least ratio is limiting reagent. [Useful when number of reactants are more than two.]




A is L.R.

Formula for checking L.R. =

$$\frac{\text{Given value (moles, volume, or molecules)}}{\text{Stoichiometry Coefficient}}$$

Least value indicate the L.R.



Some Basic Concepts of Chemistry

PERCENTAGE YIELD

The percentage yield of product

$$= \frac{\text{Actual yield}}{\text{theoretical maximum yield}} \times 100$$

The actual amount of any limiting reagent consumed is given by
 (% yield $\times$ given moles of limiting reagent).

Train Your Brain

Example 29: $\text{A} + 5\text{B} \longrightarrow \text{C} + 3\text{D}$, in this reaction which is a L.R.?

Sol. For A For B


$$\frac{10}{1} = 10 \quad \frac{10}{5} = 2$$

2 < 10 So, B is L.R.

Example 30: $\text{H}_2(\text{g}) + \frac{1}{2}\text{O}_2(\text{g}) \longrightarrow \text{H}_2\text{O}(\text{g})$; in the above

reaction, what is the volume of water vapour produced at STP?

- (a) 4.48 L (b) 44.8 L
 (c) 2.24 L (d) 11.2 L



For H_2 For O_2

$$n = \frac{4}{2} = 2 \text{ mol} \quad n = \frac{32}{32} = 1 \text{ mol}$$

$$\text{For L.R. } \frac{2}{1} = 2 \quad \frac{1}{\frac{1}{2}} = 2 \text{ mol}$$

$$\text{Moles of } \text{H}_2\text{O}(\text{g}) \text{ produced} = 2 \text{ mol} = \frac{V}{22.4}$$

Both H_2 & O_2 are L.R.

$$\text{Volume of } \text{H}_2\text{O}(\text{g}) \text{ produced at STP} = 22.4 \times 2 = 44.8 \text{ litre}$$

Example 31: At STP, In a container 100 mL N_2 and 100 mL of H_2 are mixed together. Then find out the produced volume of NH_3 .

- (a) 6.66 mL (b) 66.6 mL
 (c) 3.33 mL (d) 5.55 mL

Sol. (b) Balanced equation will be $\text{N}_2 + 3\text{H}_2 \rightarrow 2\text{NH}_3$.

Given 100mL 100mL

For determination of Limiting reagent. Divide the given quantities by stoichiometry coefficients

$$\frac{100}{1} = 100 \quad \frac{100}{3} = 33.3 \text{ (Limiting reagent)}$$

In this reaction H_2 is limiting reagent so reaction will proceed according to H_2 .


According to stoichiometry from 3 mL of H_2 produced volume of $\text{NH}_3 = 2$ ml

That is from 100 ml of H_2 produced volume of

$$\text{NH}_3 = \frac{2}{3} \times 100 = 66.6 \text{ mL}$$

Example 32: Number of moles of NH_3 produced if 140 gm of N_2 reacts with 40 gm of di-hydrogen. (Given % yield of reaction is 50%)

- (a) 12 (b) 10 (c) 5 (d) 6



or 5 mol 20 mol

$\therefore$ Number of moles of NH_3 produced = $5 \times 2 \times 0.5$
= 5 mole

Concept Application

29. 4 mole of MgCO_3 is reacted with 6 moles of HCl solution. Find the volume of CO_2 gas (in litres) produced at STP, the reaction is:




- (a) 11.2 (b) 22.4
(c) 67.2 (d) 44.8

30. For a reaction, $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightarrow 2\text{NH}_3(\text{g})$; which of the following reaction mixtures has dihydrogen (H_2) as a limiting reagent, :

- (a) 14g of N_2 + 4g of H_2 (b) 35g of N_2 + 8g of H_2
(c) 28g of N_2 + 6g of H_2 (d) 56g of N_2 + 10g of H_2


31. The percent yield for the following reaction carried out in carbon tetrachloride (CCl_4) solution is 80%.



How many moles of BrCl is formed from the reaction of 0.025 mol Br_2 and 0.025 mol Cl_2 ?

- (a) 0.04 (b) 0.08]
(c) 0.02 (d) 0.01

32. If 240 g of carbon is taken in a container to convert it completely to CO_2 but in industry it has been found that 280 g of CO was also formed along with CO_2 . Find the mole percentage yield of CO_2 . The reactions occurring are:



- (a) 25 (b) 50 (c) 75 (d) 35




ADVANCED LEARNING

SEQUENTIAL REACTION

Here, we solve problems in which the products of one reaction are used up in one or more subsequent reactions. In order to attempt such problems the following solving strategy has to be used.

- The balanced and molecular equations are written for all reactions involved separately.
- Later the equations are multiplied, as a whole, by suitable factors, so that products of one reaction which are utilized in subsequent reactions are cancelled out.
- The final reaction obtained is used to find out the required quantities.

Illustration: How many kilograms of pure H_2SO_4 could be obtained from one kilogram of pure iron pyrites (FeS_2) according to the following reactions?






- (a) 0.184kg (b) 1.633kg
(c) 2.643kg (d) 3.234kg

Sol. Gram molecular weight of $\text{FeS}_2 = 120\text{g}$

Gram molecular weight of $\text{H}_2\text{SO}_4 = 98\text{ g}$

Let us multiply the equations with suitable factors.



From the above, it is clear that, 4 moles of FeS_2 produces 8 moles of H_2SO_4 (or)

1 mole of $\text{FeS}_2 \xrightarrow{\text{produces}} 2 \text{ moles of } \text{H}_2\text{SO}_4$

$\therefore$ 120g (1 mole) of $\text{FeS}_2 \xrightarrow{\text{produces}} 2 \times 98 \text{ g (2 moles) of } \text{H}_2\text{SO}_4$

$\therefore$ 1000g (1 kg) of $\text{FeS}_2 \xrightarrow{\text{produces}} \frac{1000 \times 2 \times 98}{120} = 1633.33 \text{ g}$

or 1.633 kg of H_2SO_4

Therefore, 1.633 kg of H_2SO_4 is produced from 1 kg of iron pyrites.

Illustration: 20 g of KClO_3 on heating give enough oxygen to react completely with hydrogen produced by the action of dil. H_2SO_4 on zinc. Find the weight of zinc required for the purpose. The reactions are as follows: ($\text{K} = 39$, $\text{Zn} = 65$, $\text{Cl} = 35.5$)

Aarambh (Solved Examples)

- 1.80 g of a certain metal burnt in oxygen gave 3.0 g of its oxide. 1.50 g of the same metal heated in steam gave 2.50 g of its oxide. The law shown by above data is:
(a) Law of constant proportion
(b) Law of multiple proportion
(c) Law of reciprocal proportion
(d) All of the above

(Prayas JEE Chemistry M-1)

Sol. In the first sample of oxide,

Weight of metal = 1.80 g;

Weight of oxygen = (3.0 – 1.80) g = 1.2 g

$$\therefore \frac{\text{wt of metal}}{\text{wt of oxygen}} = \frac{1.80\text{g}}{1.2\text{g}} = 1.5$$

In the second sample of the oxide,

Weight of metal = 1.50 g;

Weight of oxygen = (2.50 – 1.50) g = 1 g

$$\frac{\text{wt of metal}}{\text{wt of oxygen}} = 1.5$$

Thus, in both samples of the oxide, the proportions of the weights of the metal and oxygen are fixed. Hence the results follow the law of constant proportion.

Therefore, option (a) is the correct answer.

- Calculate the total charge present on 4.2 gm of N^{3-} .
(a) $8.67 \times 10^4 \text{ C}$ (b) $9.65 \times 10^4 \text{ C}$
(c) $8.67 \times 10^3 \text{ C}$ (d) $7.67 \times 10^4 \text{ C}$

(Prayas JEE Chemistry M-1)

Sol. Mole = $\frac{\text{wt.in gm}}{\text{Ionic wt.}} = \frac{4.2}{14} = 0.3$

Total no. of ions = $0.3 \times N_A$ ions.

Total charge = $0.3 N_A \times 3 \times 1.6 \times 10^{-19} \text{ C}$

$$= 0.3 \times 6.022 \times 10^{23} \times 3 \times 1.6 \times 10^{-19} = 8.67 \times 10^4 \text{ C}$$

Therefore, option (a) is the correct answer.

- How many carbon atoms are present in 0.35 mol of $\text{C}_6\text{H}_{12}\text{O}_6$?
(a) 6.022×10^{23} carbon atoms.
(b) 1.26×10^{23} carbon atoms.
(c) 1.26×10^{24} carbon atoms.
(d) 6.022×10^{24} carbon atoms.

(Prayas JEE Chemistry M-1)

Sol. $\therefore$ 1 mol of $\text{C}_6\text{H}_{12}\text{O}_6$ has $6 N_A$ atoms of C

$\therefore$ 0.35 mol of $\text{C}_6\text{H}_{12}\text{O}_6$ has $6 \times 0.35 N_A$ atoms of C

$$= 2.1 N_A \text{ atoms} = 2.1 \times 6.022 \times 10^{23} = 1.26 \times 10^{24} \text{ carbon atoms.}$$

Therefore, option (c) is the correct answer.

- How many molecules are present in 5.23 gm of glucose ($\text{C}_6\text{H}_{12}\text{O}_6$)?

- 1.65×10^{22}
- 1.75×10^{22}
- 1.75×10^{21}
- None of these

(Prayas JEE Chemistry M-1)

Sol. $\therefore$ 180 gm glucose has = N_A molecules

$$\therefore 5.23 \text{ gm glucose has} = \frac{5.23 \times 6.022 \times 10^{23}}{180}$$

$$= 1.75 \times 10^{22} \text{ molecules}$$

Therefore, option (b) is the correct answer.

- A sample of (C_2H_6) ethane has the same mass as 10^7 molecules of methane. How many C_2H_6 molecules does the sample contain?
(a) 5.34×10^6 (b) 1.26×10^8
(c) 4.26×10^6 (d) 6.022×10^6

(Prayas JEE Chemistry M-1)

Sol. Moles of $\text{CH}_4 = \frac{10^7}{N_A}$

$$\text{Mass of } \text{CH}_4 = \frac{10^7}{N_A} \times 16 = \text{mass of } \text{C}_2\text{H}_6$$

$$\text{So Moles of } \text{C}_2\text{H}_6 = \frac{10^7 \times 16}{N_A \times 30}$$

$$\text{So no. of molecules of } \text{C}_2\text{H}_6 = \frac{10^7 \times 16}{N_A \times 30} \times N_A = 5.34 \times 10^6.$$

Therefore, option (a) is the correct answer.

- From 160 g of SO_2 (g) sample, 1.2046×10^{24} molecules of SO_2 are removed then find out the volume of left over SO_2 (g) at STP.
(a) 11.2 L (b) 12.5 L (c) 9.5 L (d) 10.8 L

(Prayas JEE Chemistry M-1)

Sol. Given moles = $\frac{160}{64} = 2.5$.

$$\text{Removed moles} = \frac{1.2046 \times 10^{24}}{6.022 \times 10^{23}} = 2.$$

So left moles = 0.5.

Volume left at STP = $0.5 \times 22.4 = 11.2 \text{ L}$.

Therefore, option (a) is the correct answer.

- 14 g of Nitrogen gas and 22 g of CO_2 gas are mixed together. Find the volume of gaseous mixture at STP.
(a) 10.2 L (b) 12.2 L (c) 15.5 L (d) 22.4 L

(Prayas JEE Chemistry M-1)

Sol. Moles of $\text{N}_2 = \frac{14}{28} = 0.5$.

$$\text{Moles of } \text{CO}_2 = \frac{22}{44} = 0.5.$$

So total moles = $0.5 + 0.5 = 1$.

So vol. at STP = $1 \times 22.4 = 22.4$ L.
Therefore, option (d) is the correct answer.

8. How many years it would take to spend Avogadro's number of rupees at the rate of 1 million rupees per second?
(a) 19.098×10^{19} years (b) 19.098 years
(c) 19.098×10^9 years (d) None of these

(Prayas JEE Chemistry M-1)

Sol. $\therefore$ 10^6 rupees are spent in 1 sec.
 $\therefore$ 6.022×10^{23} rupees are spent in
$$\frac{1 \times 6.022 \times 10^{23}}{10^6} \text{ sec}$$
or
$$\frac{1 \times 6.022 \times 10^{23}}{10^6 \times 60 \times 60 \times 24 \times 365} \text{ years} = 19.098 \times 10^9 \text{ years}$$
Therefore, option (c) is the correct answer.

9. The density of O_2 at STP is 1.429g/litre. Calculate the standard molar volume of gas.
(a) 22.4 lit. (b) 11.2 lit (c) 33.6 lit (d) 5.6 lit.

(Prayas JEE Chemistry M-1)

Sol. $\therefore$ 1.429 gm of O_2 gas occupies volume = 1 litre.
 $\therefore$ 32 gm of O_2 gas occupies = $\frac{32}{1.429} = 22.4$ litre/mol.
Therefore, option (a) is the correct answer.

10. Calculate the weight of lime (CaO) obtained by heating 200 kg of 95% pure lime stone ($CaCO_3$).
(a) 104.4 kg (b) 105.4 kg
(c) 212.8 kg (d) 106.4 kg

(Prayas JEE Chemistry M-1)

Sol. $\therefore$ 100 kg impure sample has pure
 $CaCO_3 = 95$ kg
 $\therefore$ 200 kg impure sample has pure $CaCO_3$
$$= \frac{95 \times 200}{100} = 190 \text{ kg.}$$
$$CaCO_3 \rightarrow CaO + CO_2$$
$$\therefore 100 \text{ kg } CaCO_3 \text{ gives } CaO = 56 \text{ kg.}$$
$$\therefore 190 \text{ kg } CaCO_3 \text{ gives } CaO = \frac{56 \times 190}{100} = 106.4 \text{ kg.}$$
Therefore, option (d) is the correct answer.

11. A compound containing beryllium has the following composition, Be = 6.1%, N = 37.8%, Cl=48%, H = 8.1%. One mole of the compound has mass of 148g and average atomic mass of beryllium is 9. The molecular formula of the compound is:

(a) $BeN_4H_{12}Cl_2$ (b) $BeN_2H_{10}Cl$
(c) $BeN_4H_2Cl_3$ (d) $Be_2N_4H_{10}Cl_2$

(Prayas JEE Chemistry M-1)

Sol.	Element	%	%/A	Simplest ratio
	Be	6.1	$6.1/9 = 0.677$	1
	N	37.8	$37.8/14 = 2.7$	4
	Cl	48	$48/35.5 = 1.35$	2

H 8.1 $8.1/1 = 8.1$ 12
Empirical formula = $BeN_4Cl_2H_{12}$
 $= 9 + 56 + 71 + 12$
 $= 148 \Rightarrow n = 1$
Molecular formula = $BeN_4Cl_2H_{12}$
Therefore, option (a) is the correct answer.

12. One litre of a mixture of CO and CO_2 is passed through red hot charcoal in tube. The new volume becomes 1.4 litre. Find out % composition of mixture by volume. All measurements are made at same P and T.
(a) CO_2 40%, CO 60% (b) CO_2 60%, CO 40%
(c) CO_2 25%, CO 75% (d) CO_2 30%, CO 70%

(Prayas JEE Chemistry M-1)

Sol. On passing through charcoal only CO_2 reduces to CO.
 $CO + C \longrightarrow$ No reaction
Volume a
 $CO_2 + C \longrightarrow 2 CO$
Volume before reaction b 0
Volume after reaction 0 2b
As given a + b = 1 and a + 2b = 1.4
 $\therefore b = 0.4$ litre $\therefore \% \text{ of } b = \frac{0.4}{1} \times 100 = 40 \%$
 $\therefore a = 0.6$ litre $\therefore \% \text{ of } a = \frac{0.6}{1} \times 100 = 60 \%$
Therefore, option (a) is the correct answer.

13. Calculate the molarity of H^+ ion in the resulting solution when 200 ml 0.5M HCl is mixed with 200 ml 0.5M H_2SO_4

(Prayas JEE Chemistry M-1)

Sol. $n_{H^+} = 0.1$ (from HCl) & $n_{H^+} = 0.2$ (from H_2SO_4)
Total $H^+ = n_{H^+}$ (from HCl) + n_{H^+} (from H_2SO_4) = $0.1 + 0.2 = 0.3$
Total volume = $200 + 200 = 400$ mL = 0.4 L
$$MR = \text{Resultant molarity} = \frac{n_{H^+}}{V_{\text{solution}}} = \frac{0.3}{0.4} = 0.75 \text{ M Ans.}$$
Therefore, [0.75] is the correct answer.

14. What are the final concentration of all the ions when following are mixed 50 ml of 0.12 M $Fe(NO_3)_3$ + 100 ml of 0.1 M $FeCl_3$ + 100 ml of 0.26 M $Mg(NO_3)_2$.

(Prayas JEE Chemistry M-1)

Sol. $[NO_3^-] = \frac{50 \times 0.12 \times 3 + 100 \times 0.26 \times 2}{250} = \frac{18 + 52}{250} = \frac{70}{250} = 0.28$
 $[Cl^-] = 0.12$ M; $[Mg^{++}] = 0.104$ M; $[Fe^{3+}] = 0.064$ M
Therefore, [0.064] is the correct answer.

15. A sample of 3 g containing Na_2CO_3 and $NaHCO_3$ loses 0.248 g when heated to $300^\circ C$, the temperature at which $NaHCO_3$ decomposes to Na_2CO_3 , CO_2 and H_2O . What is the percentage of Na_2CO_3 in the given mixture?

(Prayas JEE Chemistry M-1)

Sol. The loss in weight is due to removal of CO_2 and H_2O which escape out on heating.
wt. of Na_2CO_3 in the product = $3.00 - 0.248 = 2.752$ g
Let wt. of Na_2CO_3 in the mixture be x g

$\therefore$ wt. of $\text{NaHCO}_3 = (3.00 - x)$ g
 Since Na_2CO_3 in the products contains x g of unchanged reactant Na_2CO_3 and rest produced from NaHCO_3 .
 The wt. of Na_2CO_3 produced by $\text{NaHCO}_3 = (2.752 - x)$ g

$$\text{NaHCO}_3 \longrightarrow \text{Na}_2\text{CO}_3 + (\text{H}_2\text{O} + \text{CO}_2)\uparrow$$

$$(3.0 - x) \qquad (2.752 - x)$$
 Applying POAC for Na atom

$1 \times \text{moles of NaHCO}_3 = 2 \times \text{moles of Na}_2\text{CO}_3$

$$\Rightarrow \frac{(3-x)}{84} = 2 \times \frac{(2.752-x)}{106}$$
 $\therefore x = 2.328 \text{ g}$

$$\therefore \% \text{ of Na}_2\text{CO}_3 = \frac{2.328}{3} \times 100 = 77.6 \%$$
 Therefore, [77.6] is the correct answer.

Prarambh (Topicwise)

FUNDAMENTAL QUANTITIES, LAWS OF CHEMICAL COMBINATION

1. Express the result of $(0.582 + 324.65)$ to the appropriate number of significant figures:

(a) 325.24 (b) 325.23 (c) 325.2 (d) 325.232

(Prayas JEE Chemistry M-1)

2. The correctly reported answer of the area of rectangle which is 12.34 cm long and 1.23 cm wide is :

(a) 15.2 m² (b) 15.2 cm² (c) 15.1 cm² (d) 15.17 cm²

(Prayas JEE Chemistry M-1)

3. Two elements X and Y combine in gaseous state to form XY in the ratio 1:35.5 by mass. The mass of Y that will be required to react with 2 g of X is:

(a) 7.1 g (b) 3.55 g (c) 71 g (d) 35.5 g

(Prayas JEE Chemistry M-1)

4. 4.4 g of an oxide of nitrogen gives 2.24 L of nitrogen and 60 g of another oxide of nitrogen gives 22.4 L of nitrogen at S.T.P. The data illustrates:

(a) Law of conservation of mass
 (b) Law of constant proportions
 (c) Law of multiple proportions
 (d) Law of reciprocal proportions

(Prayas JEE Chemistry M-1)

5. Two elements X and Y combine to form compounds A, B and C. The ratio of different masses of Y which combine with a fixed mass of X in A, B and C is 1 : 3 : 5. If 32 parts by mass of X combines with 84 parts by mass of Y in B, then in C, 16 parts by mass of X will combine with;

(a) 14 parts by mass of Y (b) 42 parts by mass of Y
 (c) 70 parts by mass of Y (d) 84 parts by mass of Y

(Prayas JEE Chemistry M-1)

- (b) 4 g-atom of Hydrogen
 (c) 1.81×10^{23} molecules of CH_4
 (d) 3.0 g of carbon

(Prayas JEE Chemistry M-1)

7. 7.5 grams of a gas occupy 5.6 litres of volume at STP, the gas is:

(a) NO (b) N_2O (c) CO (d) CO_2

(Prayas JEE Chemistry M-1)

8. The number of atoms in 4.25 g of NH_3 is approximately:

(a) 1×10^{23} (b) 2×10^{23}
 (c) 4×10^{23} (d) 6×10^{23}

(Prayas JEE Chemistry M-1)

9. One litre of gas at STP weighs 1.16 g. The possible gas is:

(a) C_2H_2 (b) CO (c) O_2 (d) CH_4

(Prayas JEE Chemistry M-1)

10. If N_A is Avogadro's number, then number of valence electrons in 4.2 g of nitride ions (N^{3-}) is:

(a) $2.4 N_A$ (b) $4.2 N_A$ (c) $1.6 N_A$ (d) $3.2 N_A$

(Prayas JEE Chemistry M-1)

11. The number of molecules at STP in 1 ml of an ideal gas will be:

(a) 6×10^{23} (b) 2.69×10^{19}
 (c) 2.69×10^{23} (d) None of these

(Prayas JEE Chemistry M-1)

12. Volume of a gas at STP is 1.12×10^{-7} cc. The number of molecules in it are:

(a) 3.01×10^{20} (b) 3.01×10^{12}
 (c) 3.01×10^{23} (d) 3.01×10^{24}

(Prayas JEE Chemistry M-1)

13. 4.4 g of an unknown gas occupies 2.24 L of volume at standard temperature and pressure. The gas may be:

(a) Carbon dioxide (b) Carbon monoxide
 (c) Oxygen (d) Sulphur dioxide

(Prayas JEE Chemistry M-1)

14. The number of oxygen atoms in 4.4 g of CO_2 is approx.:

(a) 1.2×10^{23} (b) 6×10^{22}

ATOMIC MASS & MOLECULAR MASS, MOLE CONCEPT AND APPLICATIONS

6. 1 mol of CH_4 contains:

(a) 6.02×10^{23} atoms of H

- (c) 6×10^{23} (d) 12×10^{23}
(Prayas JEE Chemistry M-1)
15. The total number of protons in 10 g of calcium carbonate is: ($N_A = 6.022 \times 10^{23}$)
 (a) 1.5057×10^{24} (b) 2.0478×10^{24}
 (c) 3.0115×10^{24} (d) 4.0956×10^{24}
(Prayas JEE Chemistry M-1)
16. Number of molecules in 100 ml each of O_2 , NH_3 and CO_2 at STP are:
 (a) In the order: $CO_2 < O_2 < NH_3$
 (b) In the order: $NH_3 < O_2 < CO_2$
 (c) The same in all
 (d) In the order: $NH_3 < CO_2 < O_2$
(Prayas JEE Chemistry M-1)
17. The number of water molecules in 1 litre of water is:
 (a) 18 (b) 18×1000
 (c) N_A (d) $55.55 N_A$
(Prayas JEE Chemistry M-1)
18. 2 g of oxygen contains number of atoms equal to that in:
 (a) 0.5 g of hydrogen (b) 4 g of sulphur
 (c) 7 g of nitrogen (d) 2.3 g of sodium
(Prayas JEE Chemistry M-1)

PERCENTAGE COMPOSITION, EMPIRICAL FORMULA & MOLECULAR FORMULA

19. Caffeine has a molecular weight of 194. If it contains 28.9% by mass of nitrogen, number of atoms of nitrogen in one molecule of caffeine is:
 (a) 4 (b) 6 (c) 2 (d) 3
(Prayas JEE Chemistry M-1)
20. A compound (60 g) on analysis gave C = 24 g, H = 4 g, O = 32 g. Its empirical formula is:
 (a) $C_2H_2O_2$ (b) C_2H_2O (c) CH_2O_2 (d) CH_2O
(Prayas JEE Chemistry M-1)
21. What is the % of H_2O in $Fe(CNS)_3 \cdot 3H_2O$?
 (a) 45 (b) 30 (c) 19 (d) 25
(Prayas JEE Chemistry M-1)
22. A hydrocarbon contains 86% carbon, Then, the hydrocarbon is an:
 (a) Alkane (b) Alkene (c) Alkyne (d) Arene
(Prayas JEE Chemistry M-1)
23. The simplest formula of a compound containing 50% of element X (atomic mass 10) and 50% of element Y (atomic mass 20) is:
 (a) XY (b) X_2Y (c) XY_3 (d) X_2Y_3
(Prayas JEE Chemistry M-1)
24. Haemoglobin contains 0.33% of iron by weight. The molecular weight of haemoglobin is approximately 67,200.

The number of iron atoms (At. wt. of Fe = 56) present in one molecule of haemoglobin is:

- (a) 6 (b) 1 (c) 4 (d) 2

(Prayas JEE Chemistry M-1)

CONCENTRATION TERMS

25. What is the molarity of NaOH solution if 250 mL of it contains 1 mg of NaOH?
 (a) 10^{-1} M (b) 10^{-2} M (c) 10^{-4} M (d) 10^{-3} M
(Prayas JEE Chemistry M-1)
26. Molarity of H_2SO_4 (density 1.8 g/mL) is 18 M. The molality of this H_2SO_4 is:
 (a) 36 (b) 200 (c) 500 (d) 18
(Prayas JEE Chemistry M-1)
27. The percentage of sodium in a breakfast cereal be labeled as 110 mg of sodium per 100 g of cereal is
 (a) 11% (b) 1.10% (c) 0.11% (d) 1.10%
(Prayas JEE Chemistry M-1)
28. 200 ml, 3M NaOH is mixed with 300 ml 2 M NaOH forming a solution of density 1200 Kg/m³. Then molality of final solution is:
 (a) 2.2 m (b) 3 m (c) 2.4 m (d) 2 m
(Prayas JEE Chemistry M-1)
29. Calculate molality of 0.2 M urea solution having density 1.2 gm/mL.
 (a) $\approx \frac{1}{5}$ m (b) $\approx \frac{1}{3}$ m
 (c) $\approx \frac{1}{9}$ m (d) $\approx \frac{1}{12}$ m
(Prayas JEE Chemistry M-1)
30. Commercial HNO_3 (aq) has density 1.41 g/mL and is 10M. Then mass percentage of solution is :
 (a) $\frac{100}{3}$ % (b) 50% (c) 75% (d) 25%
(Prayas JEE Chemistry M-1)
31. 0.02 gm of an unknown substance is dissolved in 3.9 gm of benzene. The molality of solution is 0.08 m. Calculate the molecular mass of unknown substance:
 (a) 60 g/mol (b) 164 g/mol
 (c) 264 g/mol (d) 64.1 g/mol
(Prayas JEE Chemistry M-1)
32. A 500 g solution has 0.1 g fluoride concentration. Then fluoride concentration in term of ppm will be:
 (a) 200 ppm (b) 100 ppm
 (c) 400 ppm (d) 50 ppm
(Prayas JEE Chemistry M-1)

STOICHIOMETRY, EQUATION BASED CALCULATIONS

33. In the reaction;

- $4\text{NH}_3(\text{g}) + 5\text{O}_2(\text{g}) \rightarrow 4\text{NO}(\text{g}) + 6\text{H}_2\text{O}(\text{g})$, when 1 mole of ammonia and 1 mole of O_2 are made to react to completion:
 - 1.0 mole of H_2O is produced.
 - 1.0 mole of NO will be produced.
 - All the oxygen will be consumed.
 - All the ammonia will be consumed.**(Prayas JEE Chemistry M-1)**
- 34.** H_2 evolved at STP on complete reaction of 27 g of aluminium with excess of aqueous NaOH would be:

$$\text{Al} + \text{H}_2\text{O} + \text{NaOH} \rightarrow \text{NaAlO}_2 + \frac{3}{2}\text{H}_2$$
 - 22.4 litres
 - 44.8 litres
 - 67.2 litres
 - 33.6 litres**(Prayas JEE Chemistry M-1)**
- 35.** 12 g of Mg (at. mass 24) will react completely with acid to give:
 - One mole of H_2
 - $\frac{1}{2}$ mole of H_2
 - $\frac{2}{3}$ mole of O_2
 - Both $\frac{1}{2}$ mol of H_2 and $\frac{1}{2}$ mol of O_2**(Prayas JEE Chemistry M-1)**
- 36.** 100 g CaCO_3 reacts with 1 litre 1 N HCl . On completion of reaction, how much weight of CO_2 will be obtained?
 - 5.5 g
 - 11 g
 - 22 g
 - 33 g**(Prayas JEE Chemistry M-1)**
- 37.** What weight of HNO_3 is needed to convert 5 g of iodine into iodic acid according to the reaction,

$$\text{I}_2 + 10\text{HNO}_3 \rightarrow 2\text{HIO}_3 + 10\text{NO}_2 + 4\text{H}_2\text{O} ?$$
 - 12.4 g
 - 24.8 g
 - 0.248 g
 - 49.6 g**(Prayas JEE Chemistry M-1)**
- 38.** How much Cl_2 at STP is liberated when 1 mole KMnO_4 reacts with HCl ?
 - 11.2 L
 - 22.4 L
 - 44.8 L
 - 56 L**(Prayas JEE Chemistry M-1)**
- 39.** If 0.50 mole of BaCl_2 is mixed with 0.20 mol of Na_3PO_4 , the maximum number of moles of $\text{Ba}_3(\text{PO}_4)_2$ that can be formed is:
 - 0.70
 - 0.50
 - 0.20
 - 0.10**(Prayas JEE Chemistry M-1)**
- 40.** 0.5 mole of H_2SO_4 is mixed with 0.2 mole of $\text{Ca}(\text{OH})_2$. The maximum number of moles of CaSO_4 formed is:
 - 0.2
 - 0.5
 - 0.4
 - 1.5**(Prayas JEE Chemistry M-1)**

Prabal (JEE Main Level)

- 1.** A sample of calcium phosphate $\text{Ca}_3(\text{PO}_4)_2$ contains 8 mol of O atoms. The number of mole of Ca atoms in the sample is:
 - 4
 - 1.5
 - 3
 - 8**(Prayas JEE Chemistry M-1)**
- 2.** Ratio of masses of H_2SO_4 and $\text{Al}_2(\text{SO}_4)_3$ each containing 32 grams of S is _____.
 - 0.86
 - 1.72
 - 0.43
 - 2.15**(Prayas JEE Chemistry M-1)**
- 3.** Which has maximum number of atoms of oxygen?
 - 10 ml $\text{H}_2\text{O}(\text{l})$
 - 0.1 mole of V_2O_5
 - 12 gm $\text{O}_3(\text{g})$
 - 12.044×10^{22} molecules of CO_2**(Prayas JEE Chemistry M-1)**
- 4.** Mass of one atom of the element A is $3.9854 \times 10^{-23}\text{g}$. How many atoms are contained in 1g of the element A?
 - 2.509×10^{22}
 - 6.022×10^{23}
 - 12.044×10^{23}
 - None of these**(Prayas JEE Chemistry M-1)**
- 5.** The number of atoms present in 0.5 g-atoms of nitrogen is same as the atoms in:
 - 12 g of C
 - 32 g of S
 - 8 g of oxygen
 - 24 g of Mg**(Prayas JEE Chemistry M-1)**
- 6.** How many moles of magnesium phosphate $\text{Mg}_3(\text{PO}_4)_2$ will contain 0.25 mole of oxygen atoms?
 - 0.02
 - 3.125×10^{-2}
 - 1.25×10^{-2}
 - 2.5×10^{-2}**(Prayas JEE Chemistry M-1)**
- 7.** 64 g of an organic compound has 24 g carbon and 8 g hydrogen and the rest is oxygen. The empirical formula of the compound is:
 - CH_4O
 - CH_2O
 - $\text{C}_2\text{H}_4\text{O}$
 - None of these**(Prayas JEE Chemistry M-1)**
- 8.** Two elements X (atomic mass = 75) and Y (atomic mass = 16) combine to give a compound having 75.8% of X. The formula of the compound is:
 - X_2Y_3
 - X_2Y
 - X_2Y_2
 - XY**(Prayas JEE Chemistry M-1)**
- 9.** A definite amount of gaseous hydrocarbon was burnt with just sufficient amount of O_2 . The volume of all reactants was 600 ml, after the explosion the volume

of the products $[\text{CO}_2(\text{g})$ and $\text{H}_2\text{O}(\text{g})]$ was found to be 700 ml under the similar conditions. The possible molecular formula of the compound is:

- (a) C_3H_8 (b) C_3H_6 (c) C_3H_4 (d) C_4H_{10}

(Prayas JEE Chemistry M-1)

10. Mole fraction of ethyl alcohol in aqueous ethyl alcohol ($\text{C}_2\text{H}_5\text{OH}$) solution is 0.25. Hence, percentage of ethyl alcohol by weight is:

- (a) 54% (b) 25% (c) 75% (d) 46%

(Prayas JEE Chemistry M-1)

11. Weight of oxygen in Fe_2O_3 and FeO in the simple ratio for the same amount of iron, is:

- (a) 3 : 2 (b) 1 : 2 (c) 2 : 1 (d) 3 : 1

(Prayas JEE Chemistry M-1)

12. A person needs on average of 2.0 mg of riboflavin (vitamin B_2) per day. How many gm of butter should be taken by the person per day if it is the only source of riboflavin? (Butter contains 5.5 microgram riboflavin per gm.)

- (a) 363.6 gm (b) 2.75 mg
(c) 11 gm (d) 19.8 gm

(Prayas JEE Chemistry M-1)

13. The oxide of a metal contains 30% oxygen by weight. If the atomic ratio of metal and oxygen is 2 : 3, determine the atomic weight of metal.

- (a) 12 u (b) 56 u (c) 27 u (d) 52 u

(Prayas JEE Chemistry M-1)

14. When a mixture of 10 mole of SO_2 and 15 mole of O_2 was passed over catalyst, 8 mole of SO_3 was formed. How many mole of SO_2 and O_2 did not enter into combination?

- (a) 2 moles of SO_2 , 11 moles of O_2
(b) 3 moles of SO_2 , 11.5 moles of O_2
(c) 2 moles of SO_2 , 4 moles of O_2
(d) 8 moles of SO_2 , 4 moles of O_2


(Prayas JEE Chemistry M-1)

15. $\text{C}_6\text{H}_5\text{OH}(\text{g}) + \text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$
Magnitude of volume change if 30 ml of $\text{C}_6\text{H}_5\text{OH}(\text{g})$ is burnt with excess amount of oxygen, is:

- (a) 30 ml (b) 60 ml (c) 20 ml (d) 10 ml

(Prayas JEE Chemistry M-1)

16. Mass of sucrose $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ produced by mixing 84 gm of carbon, 12 gm of dihydrogen and 56 lit. O_2 at 1 atm & 273 K according to given reaction, is:



- (a) 138.5 (b) 155.5 (c) 172.5 (d) 199.5

(Prayas JEE Chemistry M-1)

17. What volume (in ml) of 0.2 M H_2SO_4 solution should be mixed with the 40 ml of 0.1 M NaOH solution such that the

resulting solution has the concentration of H_2SO_4 as $\frac{6}{55}$ M.

- (a) 70 (b) 45 (c) 30 (d) 58

(Prayas JEE Chemistry M-1)

18. For the reaction; $2x + 3y + 4z \rightarrow 5w$, initially 1 mol of x, 3 mol of y and 4 mol of z is taken. If 1.25 mol of w is obtained then % yield of this reaction is:

- (a) 50% (b) 60% (c) 70% (d) 40%

(Prayas JEE Chemistry M-1)

19. If 10 g of Ag reacts with 1 g of sulphur, the amount of Ag_2S formed will be:

- (a) 7.75 g (b) 0.775 g (c) 11 g (d) 10 g

(Prayas JEE Chemistry M-1)

20. A solution of A (MM = 20) and B (MM = 10), [Mole fraction $X_B = 0.6$] has density 0.7 gm/ml then molarity and molality of B in this solution will be _____ and _____ respectively.

- (a) 30 M, 75 m (b) 40 M, 75 m
(c) 30 M, 65 m (d) 50 M, 55 m

(Prayas JEE Chemistry M-1)

21. 125 ml of 8% w/w NaOH solution (sp. gravity = 1) is added to 125 ml of 10 % w/v HCl solution. The nature of resultant solution would be _____.

- (a) Acidic (b) Basic
(c) Neutral (d) None of these

(Prayas JEE Chemistry M-1)

22. 36.5 % (w/w) HCl has density equal to 1.20 g mL^{-1} . The molarity (M) and molality (m), respectively, are:

- (a) 15.7, 15.7 (b) 12, 12
(c) 15.7, 12 (d) 12, 15.7

(Prayas JEE Chemistry M-1)

23. 500 mL of a glucose solution contains 6.02×10^{22} molecules of glucose. The concentration of the solution is:

- (a) 0.1 M (b) 1.0 M (c) 0.2 M (d) 2.0 M

(Prayas JEE Chemistry M-1)

24. Equal moles of H_2O and NaCl are present in a solution. Hence, molality of NaCl solution is:

- (a) 0.55 (b) 55.5 (c) 1.00 (d) 0.18

(Prayas JEE Chemistry M-1)

25. Decreasing order of mass of pure NaOH in each of the aqueous solution.

- I. 50 g of 40% (w/w) NaOH
II. 50 ml of 50% (w/v) NaOH ($d_{\text{sol}} = 1.2 \text{ g/ml}$).
III. 50 g of 15 M NaOH ($d_{\text{sol}} = 1 \text{ g/ml}$).

- (a) I, II, III (b) III, II, I
(c) II, III, I (d) III = II = I

(Prayas JEE Chemistry M-1)

26. A solution of FeCl_3 is $\frac{M}{30}$, its molarity for Cl^- ion will be:
- (a) $\frac{M}{90}$ (b) $\frac{M}{30}$ (c) $\frac{M}{10}$ (d) $\frac{M}{5}$

(Prayas JEE Chemistry M-1)

27. What volume of a 0.8 M solution contains 100 milli moles of the solute?
- (a) 100 mL (b) 125 mL (c) 500 mL (d) 62.5 mL

(Prayas JEE Chemistry M-1)

28. 4.4 g of CO_2 and 2.24 litre of H_2 at STP are mixed in a container. The total number of molecules present in the container will be:
- (a) 6.022×10^{23} (b) 1.2046×10^{23}
 (c) 6.022×10^{22} (d) 6.022×10^{24}

(Prayas JEE Chemistry M-1)

29. 10 g of CaCO_3 on heating gives 5 g of the residue (as CaO). The percent yield of the reaction is approximately:
- (a) 50% (b) 72% (c) 89% (d) 100%

(Prayas JEE Chemistry M-1)

30. 33.6 g of an impure sample of sodium bicarbonate when heated strongly gave 4.4 g of CO_2 . The percentage purity of NaHCO_3 would be:
- (a) 25% (b) 50% (c) 75% (d) 100%

(Prayas JEE Chemistry M-1)

31. A gaseous mixture contains oxygen and nitrogen in the ratio of 1 : 4 by weight. Therefore the ratio of their number of molecules is:
- (a) 1 : 4 (b) 1 : 8 (c) 7 : 32 (d) 3 : 16

(Prayas JEE Chemistry M-1)

32. 2.76 g of silver carbonate on being strongly heated yields a residue weighing:
- (a) 2.16 g (b) 2.48 g (c) 2.32 g (d) 2.64 g

(Prayas JEE Chemistry M-1)

33. 12 g of alkaline earth metal gives 14.8 g of its nitride. Atomic weight of metal is:
- (a) 12 (b) 20 (c) 40 (d) 14.8

(Prayas JEE Chemistry M-1)

34. How many liters of CO_2 at STP will be formed when 0.01 mol of H_2SO_4 reacts with excess of Na_2CO_3 .
- $\text{Na}_2\text{CO}_3 + \text{H}_2\text{SO}_4 \rightarrow \text{Na}_2\text{SO}_4 + \text{CO}_2 + \text{H}_2\text{O}$
- (a) 22.4 L (b) 2.24 L (c) 0.224 L (d) 1.12 L

(Prayas JEE Chemistry M-1)

35. Equal weight of 'X' (At. wt. = 36) and 'Y' (At. wt. = 24) are reacted to form the compound X_2Y_3 . Then
- (a) X is the limiting reagent

- (b) Y is the limiting reagent

- (c) No reactant is left over and mass of X_2Y_3 formed is double the mass of 'X' taken

- (d) None of these

(Prayas JEE Chemistry M-1)

36. A person adds 1.71 gram of sugar ($\text{C}_{12}\text{H}_{22}\text{O}_{11}$) in order to sweeten his tea. The number of carbon atoms added are (mol. mass of sugar = 342)

- (a) 3.6×10^{22} (b) 7.2×10^{21}
 (c) 0.05×10^{20} (d) 6.6×10^{22}

(Prayas JEE Chemistry M-1)

37. When 100 g of ethylene polymerises entirely to polyethene, the weight of polyethene formed as per the equation $n(\text{C}_2\text{H}_4) \longrightarrow (-\text{CH}_2-\text{CH}_2-)_n$ is:
- (a) $(n/2)\text{g}$ (b) 100g
 (c) $(100/n)\text{g}$ (d) 100ng

(Prayas JEE Chemistry M-1)

38. 0.16 g of dibasic acid required 25 ml of M/10 NaOH for complete neutralization. Molecular weight of acid is:
- (a) 32 (b) 64 (c) 128 (d) 256

(Prayas JEE Chemistry M-1)

39. An aqueous solution of 6.3 g oxalic acid dihydrate is made up to 250 ml. The volume of 0.1 M NaOH required to completely neutralize 10 ml of this solution is:
- (a) 4 ml (b) 20 ml (c) 10 ml (d) 4 ml

(Prayas JEE Chemistry M-1)

40. Which of the following contains the greatest number of oxygen atoms?
- (a) 1g of O (b) 1g of O_2
 (c) 1g of O_3 (d) All same

(Prayas JEE Chemistry M-1)

41. For sequential reaction :



If % yield of (i) and (ii) reactions are 90% and 80% respectively then the overall % yield is expected to be:

- (a) 90% (b) 80% (c) 72% (d) 10%

(Prayas JEE Chemistry M-1)

42. If 32 g of O_2 contains 6.022×10^{23} molecules at STP then 32 g of S, under the same conditions, will contain,
- (a) 6.022×10^{23} S atoms (b) 3.011×10^{23} S atoms
 (c) 12.044×10^{23} S atoms (d) 1×10^{23} S atoms

(Prayas JEE Chemistry M-1)


INTEGER TYPE QUESTIONS

43. The specific gravity of a solution is 1.8, having 62% by weight of acid. It is to be diluted to specific gravity of 1.2.

What volume of water (in mL) should be added to 100 mL of this solution?

(Prayas JEE Chemistry M-1)


44. Carbon disulphide, CS₂, can be made from by-product SO₂. The overall reaction is



How much CS₂ (in kg) can be produced from 440 kg of waste SO₂ with 60 kg of coke if the SO₂ conversion is 80%?

(Prayas JEE Chemistry M-1)


45. Pure iron pyrite, FeS₂, is burnt with 50% excess air than required for complete oxidation of FeS₂, in a closed vessel.



Air contains 20% O₂ and 80% N₂, by volume. The mole percent of N₂ gas in the gaseous mixture, after complete reaction, is.

(Prayas JEE Chemistry M-1)

46. Mixture of 10 moles of Fe₂S₃, 20 moles of H₂O and 30 mole of O₂ react with 30% yield of given reaction:



Calculate moles of Fe(OH)₃ that can be produced in above reaction.

(Prayas JEE Chemistry M-1)

47. Sample of an element "X" consist of its three isotopes A₁, A₂

& A₃ and population of A₂ is three times the population of A₃. If the average molar mass of sample is 1.25, determine percentage population of A₁ (Molar masses of isotopes A₁, A₂ & A₃ are 1, 2 and 3 gm/mol respectively.)

(Prayas JEE Chemistry M-1)

48. A solution of A (MM=20) and B (MM= 10), [Mole fraction X_B = 0.6] has density 0.7 gm/ml then molarity and molality of B in this solution will be ____ M and ____ m respectively.

(Prayas JEE Chemistry M-1)

49. A piece of aluminium weighing 2.7 g is heated with 75.0 ml of H₂SO₄ (sp. gr. 1.2 containing 25% H₂SO₄ by mass). After the metal is completely dissolved, the solution is diluted to 400ml. What is the molarity of the free H₂SO₄ in the resulting solution (Multiply final answer by 10)?

(Prayas JEE Chemistry M-1)

50. The odour of skunk is caused by chemical compounds called thiols (C₄H₁₀S). These can be deodorized by reaction with household bleach (NaOCl) according to following unbalanced reaction:



How many gram of thiol can be deodorized by 74.5 gm of NaOCl?

(Prayas JEE Chemistry M-1)

ATDB.uno

Parikshit (JEE Advanced Level)

SINGLE CORRECT TYPE QUESTIONS

- (i) $2Al + 6HCl \longrightarrow 2AlCl_3 + 3H_2$
- (ii) $AlCl_3 + 3NaOH \longrightarrow Al(OH)_3 + 3NaCl$
- (iii) $Al(OH)_3 + NaOH \longrightarrow NaAlO_2 + 2H_2O$

Above series of reactions are carried out starting with 18 g of Al and 109.5 g of HCl in first step and further 100 g of NaOH is added for step (ii) and (iii). Find out limiting reagent in each step and calculate the maximum amount of NaAlO₂ that can be produced in step (iii). (Assume reactions are taken in sequence and also that each reaction goes to 100% completion)



	L.R. in step (I)	L.R. in step (II)	L.R. in step (III)	Moles of NaAlO ₂
(a)	Al	AlCl ₃	Al(OH) ₃	0.66
(b)	Al	Na(OH)	Al(OH) ₃	0.5
(c)	Al	AlCl ₃	NaOH	0.5
(d)	HCl	AlCl ₃	NaOH	0.5

(Prayas JEE Chemistry M-1)

2. A mixture of CH₄ and C₂H₂ was completely burnt in an excess of oxygen yielding equal volumes of CO₂ and steam, measured at the same temperature and pressure. The mole percent of CH₄ in the original mixture is
(a) 25% (b) 30% (c) 75% (d) 50%

(Prayas JEE Chemistry M-1)

3. 10 moles of X, 12 mole of Y and 20 moles of Z are mixed to produce a final product P, according to the given balanced reactions:



then the maximum no. of moles of P, which can be produced assuming that the products formed can also be reused in the reaction?

- (a) 6 mole (b) 9 mole (c) 10 mole (d) 12 mole

(Prayas JEE Chemistry M-1)

4. 56 gm of N₂ and 9 gm of H₂ are made to react completely to produce a mixture of NH₃ and N₂H₄. The ratio of moles

(d) 1/2 mole of A₃B₂ is left finally.

(Prayas JEE Chemistry M-1)

17. The **incorrect** statement(s) regarding 2 M MgCl₂ aqueous solution is/are: (d_{solution} = 1.09 gm/ml)
- (a) Molality of Cl⁻ is 4.44 m.
 - (b) Mole fraction of MgCl₂ is approximately 0.035.
 - (c) The conc. of MgCl₂ is 19% w/v. (approx)
 - (d) The conc. of MgCl₂ is 19 × 10⁴ ppm.


(Prayas JEE Chemistry M-1)

18. Solution containing 23 g HCOOH is/are:

- (a) 46 g of 70% $\left(\frac{w}{v}\right)$ HCOOH (d_{solution} = 1.40 g/mL)
- (b) 50 g of 10 M HCOOH (d_{solution} = 1 g/mL)
- (c) 50 g of 25% $\left(\frac{w}{w}\right)$ HCOOH
- (d) 46 g of 5 M HCOOH (d_{solution} = 1 g/mL)

(Prayas JEE Chemistry M-1)

19. A student was carrying out the following chemical reaction in Lab



He used 30 mole of A and 30 mole of B and at the end of reaction he found 40 moles of C were formed. Identify the correct statement(s).

- (a) Total mass before and after the reaction will remain same.
- (b) B is limiting reagent
- (c) % yield is 60.
- (d) At 50% yield 30 moles of C will be formed.

(Prayas JEE Chemistry M-1)

20. The non - stoichiometric compound, titanium monoxide, has a continuous range of composition from Ti_{0.75}O to TiO_{0.69}. Which of the following is/are correct regarding the possible composition of the compound? [Ti = 48]
- (a) The maximum percentage by mass of oxygen in the compound is 30.8.
 - (b) The minimum percentage by mass of titanium in the compound is 69.2.
 - (c) The minimum percentage by mass of oxygen in the compound is 18.7
 - (d) The minimum percentage by mass of titanium in the compound is 81.3.

(Prayas JEE Chemistry M-1)

21. For the following reaction;



Identify the compositions which will produce same amount of NH₃.

- (a) 140 g N₂ & 35 g H₂
- (b) 18 g H₂ & 52 g N₂
- (c) Total 20 moles of mixture having N₂ & H₂ present in stoichiometric ratio (No limiting reagent).

Some Basic Concepts of Chemistry

- (d) 136 gm of mixture having mass fraction of H₂= $\frac{6}{34}$.

(Prayas JEE Chemistry M-1)

22. Which of the following solution contains same molar concentration?
- (a) 166 g KI per liter solution
 - (b) 33.0 g (NH₄)₂SO₄ in 200 mL solution
 - (c) 25.0 g CuSO₄.5H₂O in 100 mL solution
 - (d) 27.0 mg Al³⁺ per mL solution

(Prayas JEE Chemistry M-1)

23. If 100 ml of 1M H₂SO₄ solution is mixed with 100 ml of 9.8%(w/w) H₂SO₄ solution (d = 1 g/mL), then:
- (a) Concentration of solution remains same.
 - (b) Volume of solution become 200 mL.
 - (c) Mass of H₂SO₄ in the final solution is 98 g.
 - (d) Mass of H₂SO₄ in the final solution is 19.6 g.

(Prayas JEE Chemistry M-1)

24. Aspartame is an artificial sweetener that is 160 times sweeter than sucrose (table sugar). It is marketed as Nutra sweet. Its molecular formulae is C₁₄H₁₈N₂O₅. If its 5 gm molecule are taken then which of the following is/are correct?
- (a) Mass of hydrogen is 90 gm.
 - (b) Ratio of mass of carbon to oxygen = 2.1
 - (c) Contain 5 gm atom of oxygen.
 - (d) Contain 159 mole of total atoms.

(Prayas JEE Chemistry M-1)

25. $xX + yY \longrightarrow \nu_3Y_4$ - This reaction is carried out by taking 6 moles each of X and Y respectively then
- (a) X is the limiting reagent
 - (b) 1.5 moles of X₃Y₄ is formed
 - (c) 1.5 moles of excess reagent is left behind
 - (d) 75% of excess reagent reacted

(Prayas JEE Chemistry M-1)

COMPREHENSION BASED QUESTIONS

Comprehension (Q. 26 to 28): A chemist decided to determine the molecular formula of an unknown compound. He collects following informations:

- I. Compound contain 2 : 1 ‘H’ to ‘O’ atoms (number of atoms).
- II. Compound has 40% C by mass.
- III. Approximate molecular mass of the compound is 178 g.
- IV. Compound contains C, H and O only.

26. What is the % by mass of oxygen in the compound?
- (a) 53.33%
 - (b) 88.88%
 - (c) 33.33%
 - (d) None of these
27. What is the empirical formula of the compound?
- (a) CH₃O
 - (b) CH₂O
 - (c) C₂H₂O
 - (d) CH₃O₂

(Prayas JEE Chemistry M-1)

28. Which of the following could be molecular formula of


compound?

- (a) C₆H₆O₆
- (b) C₆H₁₂O₆
- (c) C₆H₁₄O₁₂
- (d) C₆H₁₄O₆

(Prayas JEE Chemistry M-1)

Comprehension (Q. 29 to 31): Oleum is considered as a solution of SO₃ in H₂SO₄, which is obtained by passing SO₃ in solution of H₂SO₄. When 100 g sample of oleum is diluted with desired weight of H₂O then the total mass of H₂SO₄ obtained after dilution is known as % labelling in oleum.


For example, a oleum bottle labelled as 109% H₂SO₄ means the 109g total mass of pure H₂SO₄ will be formed when 100 g of oleum is diluted by 9 g of H₂O which combines with all the free SO₃ to form H₂SO₄ as



29. What is the % of free SO₃ in an oleum that is labelled as ‘104.5% H₂SO₄’?
- (a) 10
- (b) 20
- (c) 40
- (d) None of these

(Prayas JEE Chemistry M-1)

30. If excess water is added into a 100 g bottle sample labelled as 112% H₂SO₄ and is reacted with 5.3g Na₂CO₃, then find the volume of CO₂ evolved at 1 atm pressure and 300 k temperature after the completion of the reaction: [R = 0.0821 L atom mol⁻¹ K⁻¹]



- (a) 2.46 L
- (b) 24.6 L
- (c) 1.23 L
- (d) 123

(Prayas JEE Chemistry M-1)

31. 1g of oleum sample is diluted with water. The solution required 54 ml of 0.4 N NaOH for complete neutralisation. The % of free SO₃ in the sample is:
- (a) 74
- (b) 26
- (c) 20
- (d) None of these

(Prayas JEE Chemistry M-1)

Comprehension (Q. 32 to 34): According to the Avogadro’s law, equal number of moles of gases occupy the same volume at identical condition of temperature and pressure.

Now let us assume air to consist of 80% by volume of nitrogen (N₂) and 20% by volume of oxygen (O₂). If air is taken at STP, then its 1 mol would occupy 22.4 L. 1 mol of air would contain 0.8 mol of N₂ and 0.2 mol of O₂, hence the mole fractions of N₂ and O₂ are given by X_{N₂} = 0.8 and X_{O₂} = 0.2.

32. Volume occupied by air at STP containing exactly 11.2 g of nitrogen, is:
- (a) 22.4 L
- (b) 8.96 L
- (c) 11.2 L
- (d) 2.24 L

(Prayas JEE Chemistry M-1)

33. If air is treated as a solution of O₂ and N₂, then % w/w of oxygen is:
- (a) $\frac{10}{9}$
- (b) $\frac{200}{9}$
- (c) $\frac{700}{9}$
- (d) $\frac{350}{9}$

(Prayas JEE Chemistry M-1)

34. Density of air at STP is:

- (a) 1 g/L
- (b) $\frac{9}{7}$ g/L
- (c) $\frac{2}{7}$ g/L
- (d) Can’t be determined

(Prayas JEE Chemistry M-1)

MATCH THE COLUMN TYPE QUESTIONS

35.

Column-I		Column-II	
A.	100 mL of 0.2 M AlCl ₃ solution + 400 ml of 0.1 M HCl solution	p.	Total concentration of cation(s) = 0.12 M
B.	50 mL of 0.4 M KCl + 50 ml H ₂ O	q.	[SO ₄ ²⁻] = 0.06 M
C.	30 mL of 0.2 M K ₂ SO ₄ + 70 ml H ₂ O	r.	[SO ₄ ²⁻] = 2.5 M
D.	200 mL 24.5% (w/v) H ₂ SO ₄	s.	[Cl ⁻] = 0.2 M

- (a) A → r,s; B → s; C → p,q; D → r
- (b) A → p,s; B → s; C → p,q; D → r
- (c) A → p,s; B → s; C → s,q; D → r
- (d) A → p, B → p,s; C → p,q; D → r

(Prayas JEE Chemistry M-1)

36.

Column-I		Column-II	
A.	A gaseous organic compound containing C = 52.17%, H = 13.04% & O = 34.78% (by weight) having molar mass 46 g/mol.	p.	One mole of compound contains 4N _A atoms of hydrogen.
B.	0.3 g of an organic compound containing C, H and O on combustion yields 0.44 g of CO ₂ and 0.18 g of H ₂ O, and two O per molecule	q.	The empirical formula of the compound is same as its molecular formula.
C.	A hydrocarbon containing C = 42.857% and H = 57.143% (by mole) and 3C atoms per molecule.	r.	Combustion products of one mole of compound contains larger number of moles of CO ₂ than that of H ₂ O.



PYQ's (Past Year Questions)

PROPERTIES OF MATTER AND THEIR MEASUREMENT

1. The candela is the luminous intensity, in a given direction, of a source that emits monochromatic radiation of frequency ' $A \times 10^{12}$ hertz and that has a radiant intensity in that direction of $\frac{1}{B}$ watt per steradian. ' A ' and ' B ' are respectively.

[09 April, 2024 (Shift-II)]

- (a) 540 and $\frac{1}{683}$ (b) 540 and 683
(c) 450 and $\frac{1}{683}$ (d) 450 and 683

(Prayas JEE Chemistry M-1)

UNCERTAINTY IN MEASUREMENT AND LAWS OF CHEMICAL COMBINATIONS

2. Which of the following have same number of significant figures?

[8 April, 2023 (Shift-II)]

- A. 0.00253 B. 1.0003
C. 15.0 D. 163

(Prayas JEE Chemistry M-1)

Choose the correct answer from the options given below:

- (a) A, B and C only (b) C and D only
(c) A, C and D only (d) B and C only

(Prayas JEE Chemistry M-1)

3. Using the rules for significant figures, the correct answer for the expression $\frac{0.02858 \times 0.112}{0.5702}$

[29 June, 2022 (Shift-II)]

- (a) 0.005613 (b) 0.00561
(c) 0.0056 (d) 0.006

(Prayas JEE Chemistry M-1)

4. The number of significant figures in 50000.020×10^{-3} is _____.

[26 Feb, 2021 (Shift-I)]

(Prayas JEE Chemistry M-1)

5. To check the principle of multiple proportions, a series of pure binary compounds (P_mQ_n) were analyzed and their composition is tabulated below. The correct option(s) is(are)

[JEE Adv 2022]

Compound	Weight % of P	Weight % of Q
1	50	50
2	44.4	55.6
3	40	60

- (a) If empirical formula of compound 3 is P_3Q_4 , then the empirical formula of compound 2 is P_3Q_5 .
(b) If empirical formula of compound 3 is P_3Q_2 and atomic weight of element P is 20, then the atomic weight of Q is 45.
(c) If empirical formula of compound 2 is PQ, then the empirical formula of the compound 1 is P_5Q_4 .
(d) If atomic weight of P and Q are 70 and 35, respectively,

then the empirical formula of compound 1 is P_2Q .

(Prayas JEE Chemistry M-1)

DALTON'S ATOMIC THEORY

6. Choose the Incorrect Statement about Dalton's Atomic Theory

[04 April, 2024 (Shift-II)]

- (a) Compounds are formed when atoms of different elements combine in any ratio
(b) All the atoms of a given element have identical properties including identical mass
(c) Matter consists of indivisible atoms
(d) Chemical reactions involve reorganization of atoms

(Prayas JEE Chemistry M-1)

7. The incorrect postulates of the Dalton's atomic theory are:

- (A) Atoms of different elements differ in mass.
(B) Matter consists of divisible atoms.
(C) Compounds are formed when atoms of different element combine in a fixed ratio.
(D) All the atoms of given element have different properties including mass.
(E) Chemical reactions involve reorganisation of atoms.

Choose the correct answer from the options given below:

[05 April, 2024 (Shift-I)]

- (a) (B), (D), (E) only (b) (A), (B), (D) only
(c) (C), (D), (E) only (d) (B), (D) only

(Prayas JEE Chemistry M-1)

ATOMIC & MOLECULAR MASSES

8. The average molar mass of chlorine is 35.5 g mol^{-1} . The ratio of ^{35}Cl to ^{37}Cl in naturally occurring chlorine is close to:

[6 Sept, 2020 (Shift-II)]

- (a) 4 : 1 (b) 3 : 1 (c) 2 : 1 (d) 1 : 1

(Prayas JEE Chemistry M-1)

MOLE CONCEPT AND MOLAR MASSES

9. 0.05 cm thick coating of silver is deposited on a plate of 0.05 m^2 area. The number of silver atoms deposited on plate are _____ $\times 10^{23}$. (At mass Ag = 108, $d = 7.9 \text{ g cm}^{-3}$)

[30 Jan, 2024 (Shift-I)]

(Prayas JEE Chemistry M-1)

10. Match List-I with List-II. [10 April, 2023 (Shift-II)]

List-I		List-II	
A.	16g of $\text{CH}_4(\text{g})$	p.	Weighs 28 g
B.	1g of $\text{H}_2(\text{g})$	q.	60.2×10^{23} electrons



- (a) CH_3COOH (b) HCHO
(c) CH_3OOCH_3 (d) CH_3CHO

(Prayas JEE Chemistry M-1)

24. A 2.0 g sample containing MnO_2 is treated with HCl liberating Cl_2 . The Cl_2 gas is passed into a solution of KI and 60.0 mL of 0.1 M $\text{Na}_2\text{S}_2\text{O}_3$ is required to titrate the liberated iodine. The percentage of MnO_2 in the sample is _____. (Nearest integer)

[Atomic masses (in u) $\text{Mn} = 55$; $\text{Cl} = 35.5$; $\text{O} = 16$, $\text{I} = 127$, $\text{Na} = 23$, $\text{K} = 39$, $\text{S} = 32$] [28 June, 2022 (Shift-I)]

(Prayas JEE Chemistry M-1)

25. Complete combustion of 750g of an organic compound provides 420 g of CO_2 and 210 g of H_2O . The percentage composition of carbon and hydrogen in organic compound is 15.3 and _____ respectively. (Round off to the nearest Integer). [16 March, 2021 (Shift-I)]

(Prayas JEE Chemistry M-1)

26. A 10 mg effervescent tablet containing sodium bicarbonate and oxalic acid releases 0.25 ml of CO_2 at $T = 298.15\text{ K}$ and $p = 1\text{ bar}$. If molar volume of CO_2 is 25.0 L under such condition, what is the percentage of sodium bicarbonate in each tablet? [Molar mass of $\text{NaHCO}_3 = 84\text{ g mol}^{-1}$]

[11 Jan, 2019 (Shift-I)]

- (a) 0.84 (b) 33.6 (c) 16.1 (d) 8.4

(Praya JEE Chemistry M-1)

STOICHIOMETRY & STOICHIOMETRIC CALCULATIONS

27. Mass of methane required to produce 22 g of CO_2 after complete combustion is _____ g.

(Given Molar mass in g mol^{-1}

$\text{C} = 12.0$

$\text{H} = 1.0$

$\text{O} = 16.0$)


[27 Jan, 2024 (Shift-I)]

(Prayas JEE Chemistry M-1)

28. Number of moles of methane required to produce 22g $\text{CO}_{2(\text{g})}$ after combustion is $x \times 10^{-2}$ moles. The value of x is [31 Jan, 2024 (Shift-I)]

(Prayas JEE Chemistry M-1)

29. Consider the following reaction:



If 72 mmol of PbCl_2 is mixed with 50 mmol of $(\text{NH}_4)_3\text{PO}_4$, then amount of $\text{Pb}_3(\text{PO}_4)_2$ formed is _____ mmol. (nearest integer) [01 Feb, 2024 (Shift-I)]

(Prayas JEE Chemistry M-1)

30. Combustion of glucose ($\text{C}_6\text{H}_{12}\text{O}_6$) produces CO_2 and water. The amount of oxygen (in g) required for the complete

combustion of 900 g of glucose is: (Molar mass of glucose in g mol^{-1}) = 180 [08 April, 2024 (Shift-I)]

- (a) 480 (b) 960 (c) 800 (d) 32

(Prayas JEE Chemistry M-1)

31. The number of moles of methane required to produce 11g $\text{CO}_{2(\text{g})}$ after complete combustion is: (Given molar mass of methane in g mol^{-1} : 16)

[05 April, 2024 (Shift-II)]

- (a) 0.75 (b) 0.25 (c) 0.35 (d) 0.5

(Prayas JEE Chemistry M-1)

32. 1 g of a carbonate (M_2CO_3) on treatment with excess HCl produces 0.01 mol of CO_2 . The molar mass of M_2CO_3 is _____ g mol^{-1} . (Nearest integer) [13 April, 2023 (Shift-II)]

(Prayas JEE Chemistry M-1)

33. If a rocket runs on a fuel ($\text{C}_{15}\text{H}_{30}$) and liquid oxygen, the weight of oxygen required and CO_2 released for every litre of fuel respectively are: [24 June, 2022 (Shift-I)]

(Given: density of the fuel is 0.756 g / mL)

- (a) 1188 g and 1296 g (b) 2376 g and 2592 g


- (c) 2592 g and 2376 g (d) 3429 g and 3142 g

(Prayas JEE Chemistry M-1)

34. A 0.166 g sample of an organic compound was digested with conc. H_2SO_4 and then distilled with NaOH . The ammonia gas evolved was passed through 50.0 mL of 0.5 N H_2SO_4 . The used acid required 30.0 mL of 0.25 N NaOH for complete neutralization. The mass percentage of nitrogen in the organic compound is _____.

[24 June, 2022 (Shift-I)]

(Prayas JEE Chemistry M-1)



In the above reaction, 3.9 g of benzene on nitration gives 4.92 g of nitrobenzene. The percentage yield of nitrobenzene in the above reaction is _____ %


(Round off to the nearest integer).

(Given atomic mass : $\text{C} : 12.0\text{ u}$, $\text{H} : 1.0\text{ u}$, $\text{O} : 16.0\text{ u}$, $\text{N} : 14.0\text{ u}$)

[17 March, 2021 (Shift-I)]

(Prayas JEE Chemistry M-1)

36. For a reaction,



Identify dihydrogen (H_2) as a limiting reagent in the following reaction mixtures. [9 April, 2019 (Shift-I)]

- (a) 14 g of N_2 + 4 g of H_2 (b) 28 g of N_2 + 6 g of H_2

- (c) 56 g of N_2 + 10 g of H_2 (d) 35 g of N_2 + 8 g of H_2

(Prayas JEE Chemistry M-1)

PW Challengers

NUMERICAL TYPE QUESTIONS (ANSWER UPTO TWO DECIMAL PLACE)

- Excess of calcium orthophosphate is reacted with magnesium to form Ca_3P_2 along with MgO . Ca_3P_2 on reaction with water liberated PH_3 along with $\text{Ca}(\text{OH})_2$. PH_3 is burnt in excess of oxygen to form P_2O_5 along with water. Oxides of magnesium and phosphorous react to give magnesium metaphosphate. Calculate grams of magnesium metaphosphate obtained if 1.92 gm of Mg is taken (Mg = 24, P = 31, O = 16)

(Prayas JEE Chemistry M-1)

- In one process for waterproofing, a fabric is exposed to $(\text{CH}_3)_2\text{SiCl}_2$ vapour. The vapour reacts with hydroxyl groups on the surface of the fabric or with traces of water to form the waterproofing film $[(\text{CH}_3)_2\text{SiO}]_n$, by the reaction

$$n[(\text{CH}_3)_2\text{SiCl}_2 + 2\text{OH}^- \rightarrow 2\text{Cl}^- + \text{nH}_2\text{O} + [(\text{CH}_3)_2\text{SiO}]_n$$
Where n stands for a large integer. The waterproofing film is deposited on the fabric layer upon layer. Each layer is 6.0 Å thick [the thickness of the $(\text{CH}_3)_2\text{SiO}$ group]. How much (in g) $(\text{CH}_3)_2\text{SiCl}_2$ is needed to waterproof one side of a piece of fabric, 1.00 m by 3.70 m, with a film 300 layers thick ? The density of the film is 1.0 g/cm³. (Si = 28)

Molar mass of $(\text{CH}_3)_2\text{SiO}$ = 74 g/mol

Molar mass of $(\text{CH}_3)_2\text{SiCl}_2$ = 129 g/mol

(Prayas JEE Chemistry M-1)

INTEGER TYPE QUESTIONS

- Cis-platin $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$, a compound used in cancer treatment is prepared by the reaction of ammonia with potassium tetrachloroplatinate.

$$\text{K}_2\text{PtCl}_4 + 2\text{NH}_3 \rightarrow 2\text{KCl} + \text{Pt}(\text{NH}_3)_2\text{Cl}_2$$
 - How many grams of cis-platin are formed from 41.5 gm K_2PtCl_4 and 34 gm NH_3 , if the reaction takes place in 90% yield? (Ans. x)
 - What is the maximum mass (in gm) of KCl which can be produced if initially total 9 moles of reactant are taken. Assuming 100% reaction ? (Ans. y)
What is the value of (x + y)? [Pt = 195, K = 39, Cl = 35.5]

(Prayas JEE Chemistry M-1)

- The number of Alkoxy groups in an organic compound $\text{A}(\text{OR})_x$ may be determined by the sequential reaction.

$$\text{A}(\text{OR})_x + x\text{HI} \rightarrow \text{A}(\text{OH})_x + x\text{RI}$$

$$\text{RI} + \text{Ag}^+ + \text{H}_2\text{O} \rightarrow \text{ROH} + \text{AgI}(\text{s}) + \text{H}^+$$
When 4.8 gm of organic compound $\text{A}(\text{OR})_x$ (molar mass = 240 gm mol⁻¹) is treated as above, 9.4 gm AgI is precipitated. Then, calculate number of alkoxy groups in the compound $\text{A}(\text{OR})_x$? (I = 127, Ag = 108)

(Prayas JEE Chemistry M-1)


- A sample of ammonia contains only H^1 and H^2 isotopes of hydrogen in 4 : 1 ratio and N^{14} and N^{15} isotopes of nitrogen in 3 : 1 ratio. How many neutrons are present in 1.785 mg of ammonia? (Answer in the order of 10^{18}) (N_A = 6×10^{23})

(Prayas JEE Chemistry M-1)

- In order to determine the composition of a mixture of halides containing MBr_2 & NaI , 14 gm mixture was dissolved in water. To this solution, AgNO_3 solution of certain molarity was added gradually. The mass of precipitate produced (in gm) were measured and it was plotted against volume of AgNO_3 solution added (in ml). If it is known that AgI is precipitated first and precipitation of Br^- does not start until the already precipitating I^- precipitates completely. Find out the value of $\text{AB} \times \text{CD}$ where:

AB = Atomic weight of metal 'M'


CD = Mole percent of NaI in original mixture.



(Molar mass of NaI = 149.89g/mol, Br = 79.9g/mol, AgI = 234.77g/mol)

(Prayas JEE Chemistry M-1)

- A mixture of gases liberated upon decomposition of 33.1 gm $\text{Pb}(\text{NO}_3)_2$ is dissolved in 10ml of water. What is the mass (in g) of 0.1M KOH solution with density 1.05 g/ml required to neutralise this acid. The reactions involved are: (at. mass of Pb = 207)



(Prayas JEE Chemistry M-1)

- Water is the working fluid in a thermal plant for producing electricity. Coal is Combusted for Generating heat as per reaction, $\text{C} + \text{O}_2 \rightarrow \text{CO}_2$. 0.01% of the released CO_2 gas is absorbed in water and gets converted into weak acid, H_2CO_3 which dissociate to give H^+ as $\text{H}_2\text{CO}_3 \rightarrow 2\text{H}^+ + \text{CO}_3^{2-}$. The % dissociation of acid is 5%. Assume no ionisation of H_2O . If in a certain application $[\text{H}^+]$ concentration can maximum be 10^{-5} M, then
 - Calculate maximum no. of moles of $\text{H}^+(x)$ and $\text{CO}_3^{2-}(y)$ in water if 10^9 litres of H_2O is used.
 - Calculate maximum no. of mole of carbon (z) which can be burnt so that water remains fit to be used.

What is the value of xy^2/z ?

(Prayas JEE Chemistry M-1)

9. A mother cell disintegrates into 60 identical cells and each daughter cell of mother disintegrates into 24 smaller cells, The smallest cell is uniform cylindrical in shape with diameter $120\text{\AA}$ and each cell is $6000\text{\AA}$ long. Determine molar mass of the mother cell, if density of the smallest cell is 1.12 gm/cm^3 . Using scientific notation if your answer is $x \times 10^y$, then write the value of $[x] + y$, where $[]$ is an integer function. Take avogadro number as 6×10^{23} .

(Prayas JEE Chemistry M-1)

10. Once Tom & Jerry entered a chemistry lab where a chemist was preparing 3L H_2SO_4 solution. He labelled the solution as 10 m ($d_{\text{solution}} = 3.3\text{ gm/ml}$). As the chemist left the lab, a mischief came in Tom's mind. He tried to throw the solution on Jerry but failed. In doing so some of the H_2SO_4 solution fell on the floor, so he added water to make it again to 3L. The chemist returned back & got astonished when he saw the result of analysis, that were $d = 1.5\text{ gm/ml}$ and $\% \text{ w/w} = 98$. Find out the number of moles of H_2SO_4 which fell down on the floor.

(Prayas JEE Chemistry M-1)

11. A polymeric substance, tetrafluoroethylene, can be represented by the formula $(\text{C}_2\text{F}_4)_x$, where x is a large number. The material was prepared by polymerizing C_2F_4 in the presence of a sulphur bearing catalyst that serves as a nucleus upon which the polymer grew. The final product was found to contain 0.012% S. What is the value of x , if each polymeric molecule contains one sulphur atom? Assume that the catalyst contributes a negligible amount to the total mass of the polymer. ($F = 19$, $S = 32$)


(Prayas JEE Chemistry M-1)

SINGLE CHOICE TYPE QUESTIONS

12. Density of water varies with temperature as shown.
 $d_{t^\circ\text{C}} = d_{0^\circ\text{C}} - (0.002 \times t^\circ\text{C})$; where
 $d_{t^\circ\text{C}}$ = density at $t^\circ\text{C}$ and $d_{0^\circ\text{C}}$ = density at $0^\circ\text{C} = 1\text{ gm/ml}$
Calculate % change in molarity and molality respectively of pure water due to change in temperature from 0°C to 100°C .
(a) 10%, no change (b) no change, 20%
(c) 20%, no change (d) 20%, 20%

(Prayas JEE Chemistry M-1)

ATDB.uno



ANSWER KEY

CONCEPT APPLICATION

- | | | | | | | | | | |
|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 1. (a) | 2. (a) | 3. (d) | 4. (a) | 5. (a) | 6. (a) | 7. (a) | 8. (b) | 9. (b) | 10. (b) |
| 11. (a) | 12. (b) | 13. (c) | 14. (b) | 15. (a) | 16. (b) | 17. (b) | 18. (a) | 19. (c) | 20. (d) |
| 21. (d) | 22. (b) | 23. (a) | 24. (c) | 25. (b) | 26. (c) | 27. (c) | 28. (b) | 29. (c) | 30. (d) |
| 31. (a) | 32. (b) | | | | | | | | |

PRARAMBHAH (TOPICWISE)

- | | | | | | | | | | |
|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 1. (b) | 2. (b) | 3. (c) | 4. (c) | 5. (c) | 6. (b) | 7. (a) | 8. (d) | 9. (a) | 10. (a) |
| 11. (b) | 12. (b) | 13. (a) | 14. (a) | 15. (c) | 16. (c) | 17. (d) | 18. (b) | 19. (a) | 20. (d) |
| 21. (c) | 22. (b) | 23. (b) | 24. (c) | 25. (c) | 26. (c) | 27. (c) | 28. (a) | 29. (a) | 30. (a) |
| 31. (d) | 32. (a) | 33. (c) | 34. (d) | 35. (b) | 36. (c) | 37. (a) | 38. (d) | 39. (d) | 40. (a) |

PRABAL (JEE MAIN LEVEL)

- | | | | | | | | | | |
|-----------|---------|-----------|-----------|---------|---------|---------|----------------|---------|---------|
| 1. (c) | 2. (a) | 3. (c) | 4. (a) | 5. (c) | 6. (b) | 7. (a) | 8. (a) | 9. (a) | 10. (d) |
| 11. (a) | 12. (a) | 13. (b) | 14. (a) | 15. (b) | 16. (b) | 17. (a) | 18. (a) | 19. (a) | 20. (a) |
| 21. (a) | 22. (d) | 23. (c) | 24. (b) | 25. (b) | 26. (c) | 27. (b) | 28. (b) | 29. (c) | 30. (b) |
| 31. (c) | 32. (a) | 33. (c) | 34. (c) | 35. (c) | 36. (a) | 37. (b) | 38. (c) | 39. (a) | 40. (d) |
| 41. (c) | 42. (a) | 43. [300] | 44. [1.6] | 45. [8] | 46. [4] | 47. [8] | 48. [30M, 75m] | 49. [2] | |
| 50. [180] | | | | | | | | | |

PARIKSHIT (JEE ADVANCED LEVEL)

- | | | | | | | | | | |
|-----------|-------------|-------------|-------------|-------------|------------|--------------|-----------|-------------|-------------|
| 1. (c) | 2. (d) | 3. (c) | 4. (c) | 5. (a) | 6. (b) | 7. (a) | 8. (a) | 9. (c) | 10. (b) |
| 11. (a) | 12. (c) | 13. (a,b,c) | 14. (b,c,d) | 15. (a,c) | 16. (b,d) | 17. (b,d) | 18. (a,b) | 19. (a,b,d) | 20. (a,b,c) |
| 21. (a,c) | 22. (a,c,d) | 23. (a,b,d) | 24. (a,b) | 25. (b,c,d) | 26. (a) | 27. (b) | 28. (b) | 29. (b) | 30. (c) |
| 31. (b) | 32. (c) | 33. (b) | 34. (b) | 35. (b) | 36. (d) | 37. [1811.5] | 38. [0.3] | 39. [73.6] | |
| 40. [3] | 41. [28] | 42. [3] | 43. [75] | 44. [50] | 45. [2400] | 46. [32] | | | |

PYQ's (PAST YEAR QUESTIONS)

- | | | | | | | | | | |
|-----------|-----------|---------|------------|-----------|------------|---------|----------|----------|---------|
| 1. (b) | 2. (c) | 3. (b) | 4. [8] | 5. (b, c) | 6. (a) | 7. (d) | 8. (b) | 9. [11] | 10. (d) |
| 11. [200] | 12. (c) | 13. [1] | 14. [2130] | 15. (c) | 16. [6.15] | 17. (a) | 18. [14] | 19. (d) | 20. (a) |
| 21. (a) | 22. [56] | 23. (b) | 24. [13] | 25. [3] | 26. (d) | 27. [8] | 28. [50] | 29. [24] | 30. (b) |
| 31. (b) | 32. [100] | 33. (c) | 34. [63] | 35. [80] | 36. (c) | | | | |

PW CHALLENGERS

- | | | | | | | | | | |
|------------|-----------|----------|--------|----------|-----------|-----------|----------|---------|---------|
| 1. [1.82] | 2. [1.16] | 3. [474] | 4. [2] | 5. [471] | 6. [4994] | 7. [2100] | 8. [250] | 9. [16] | 10. [5] |
| 11. [2667] | 12. (c) | | | | | | | | |

Some Basic Concepts of Chemistry