

PRAKAS

JEE 2026


ATDB.uno

PHYSICAL CHEMISTRY

SOLUTIONS

Lecture - 10

FAISAL RAZAQ





Topics to be covered

ATDB.uno

A

Van't Hoff factor





Osmotic Pressure $\rightarrow \pi = CRT$
 $\rightarrow \pi = h\rho g$

$$1 \text{ L} = 10^{-3} \text{ m}^3 //$$

$$\text{gm/cc or gm/ml} = \frac{10^{-3} \text{ kg}}{10^{-6} \text{ m}^3} = 10^3 \text{ kg/m}^3 //$$

$$R = 8.314 \text{ J/K-mol} //$$

if ($\pi = \text{atm}$)

$$C = \text{mol/L}$$

$$R = 0.0821 \text{ atm-L/K-mol}$$

$$T = \text{kelvin}$$

if ($\pi = \text{N/m}^2$ or pascals)

$$C = 10^3 \text{ mol/m}^3$$


$$R = 8.314 \text{ J/K-mol}$$

$$T = \text{kelvin}$$

$\pi = \text{osmotic pressure}$

$$\pi' > \pi$$

$$(\pi' - \pi) = h \rho g$$



To carry out reverse osmosis we need to increase pressure on higher

ATDB.uno solution side continuously.

Hypertonic, Hypotonic and isotonic solutions



Solutions having more osmotic pressure are hypertonic solutions. What the solutions with low osmotic pressure.

Solⁿ A is hypertonic

Solⁿ B is hypotonic

$$C_1 R T_1 > C_2 R T_2$$

ATDB.uno



$$C_1 T_1 > C_2 T_2$$

$$C_1 > C_2 \rightarrow \pi_1 = \pi_2 \text{ if } \pi_1 > \pi_2$$

$$\pi_1 = \pi_2 \text{ if } \pi_1 > \pi_2$$

Solutions having same osmotic pressure are isotonic.



$$\pi_1 = \pi_2$$

$$C_1 R T_1 = C_2 R T_2$$

$$C_1 T_1 = C_2 T_2$$


$$C_1 = C_2 \rightarrow T_1 = T_2$$

ATDB.uno

Osmotic Pressure of solution having two or more solutes



Osmotic pressure of a solⁿ having two or more solutes is equal to the sum of partial osmotic pressures.



$$\pi_{\text{net}} = \pi_1 + \pi_2$$

$$\pi_{\text{net}} = \sum \pi_i$$

ATDB.uno

$$\pi_1 = C_1 R T = \frac{n_A}{V} R T$$


$$\pi_2 = C_2 R T = \frac{n_B}{V} R T$$

*

$$\pi_{\text{net}} = \frac{(n_A + n_B) R T}{V}$$

Osmotic pressure of a resultant solution when two solutions of same substance of different osmotic pressure are mixed

$$\text{Net conc} = \frac{\text{net mole}}{\text{net volume}} = \left(\frac{\frac{\pi_1 V_1}{RT} + \frac{\pi_2 V_2}{RT}}{V_1 + V_2} \right)$$



$$\pi_{\text{net}} RT = \frac{\pi_1 V_1 + \pi_2 V_2}{V_1 + V_2}$$

$$\pi_1 = C_1 RT = \frac{n_1}{V_1} RT$$

$$\pi_2 = C_2 RT = \frac{n_2}{V_2} RT$$



Question

Two solutions of glucose have osmotic pressures 1.5 atm and 2.5 atm. 1 L of first solution is mixed with 3 L of second solution. Calculate the Osmotic pressure of resultant solution.

$$\pi_{net} = \frac{\sum(\pi V)}{\sum V}$$

$$\pi_{net} = \frac{\sum n RT}{V}$$

A solⁿ is having more than one solute

A solution is formed by adding two or more solutions.

$$= \frac{1.5 \times 1 + 2.5 \times 3}{4} = 2.25 \text{ atm}$$


Question

$$\bar{\kappa}_{net} = C_{net} R T$$



$$M_1, V_1$$

+



$$M_2, V_2$$



$$\bar{\kappa}_{net} = ?$$

ATDB.uno

$$\bar{\kappa}_{net} = \left(\frac{M_1 V_1 + M_2 V_2}{V_1 + V_2} \right) R T$$

Concept of ice separation



$$\Delta T_F = K_F m$$

$$= K_F \cdot \left(\frac{\text{mole of solute}}{\text{Weight of H}_2\text{O}} \right)$$

ATDB.uno

Question

A solution contains 62 gm ethylene glycol ($\text{CH}_2\text{OH}-\text{CH}_2\text{OH}$)
 in 250 gm H_2O is cooled upto -10°C . If $K_f = 1.86 \text{ K Kg mol}^{-1}$,
 then amount of water (in gm) separated as ice is -

A) 32

B) 48

✓ C) 64

D) 16

$$\Delta T_f = 10 = K_f \cdot m \quad \left[\text{JEE-Mains 2019} \right]$$

$$10 = 1.86 \times \left(\frac{62}{W} \right)$$

$$W = 0.186 \text{ Kg} = 186 \text{ g}$$

$$\text{weight of ice separated} = 250 - 186 = 64 \text{ gm.}$$



Van't Hoff factor (i)

1 mole salt (NaCl) is dissolved
in 1 Kg water. Find out ΔT_b .

$$\Delta T_b = K_b \cdot m = K_b \times \frac{1}{1} \text{ ATDB.uno}$$

$$\Delta T_b = K_b$$

$$\text{Van't Hoff factor} = \frac{\text{Obs CP}}{\text{Th CP}}$$

$$\text{Obs CP} = i \cdot \text{Th CP}$$

$$RLVP = \frac{p^0 - p_s}{p^0} = \frac{i n_{\text{solute}}}{i n_{\text{solute}} + n_{\text{solvent}}} \approx \frac{i n_{\text{solute}}}{n_{\text{solvent}}}$$

$$\Delta T_b = i K_b \cdot m$$


$$\Delta T_f = i K_f \cdot m$$

$$\pi = i CRT$$

ATDB.uno



Van t Hoff factor during dissociation



ATDB.uno

$n =$ इतने logo mein solute toota.

$$(\Delta T_b)_{th} = K_b \left(\frac{C}{W} \right)$$

$$(\Delta T_b)_{obs} = K_b \frac{C[1+(n-1)\alpha]}{W}$$

if the dissociation is $\alpha =$ इतने मे से α toota

$$\text{moles of solute after diss} = C(1-\alpha) + nC\alpha = C[1+(n-1)\alpha]$$




$$i = \frac{(\Delta T_b)_{obs}}{(\Delta T_b)_{th}} = \frac{\cancel{k_b} \cdot \frac{c [1 + (n-1)\alpha]}{\cancel{w}}}{\cancel{k_b} \cdot \frac{\cancel{c}}{\cancel{w}}} = 1 + (n-1)\alpha$$

$$i = 1 + (n-1)\alpha$$

n = [itne logo mei toota
 α = degree of dissociation

Van't Hoff factor during association

$n=2$ dimer
 $n=3$ trimer



ATDB.uno

β = degree of association

$$\text{Total moles of solute} = c(1-\beta) + \frac{c\beta}{n} = c \left[1 - \left(1 - \frac{1}{n}\right)\beta \right]$$

$$\begin{aligned} (\Delta T_f)_{th} &= K_f \left(\frac{c}{w} \right) \\ (\Delta T_f)_{obs} &= K_f \frac{c \left[1 - \left(1 - \frac{1}{n}\right)\beta \right]}{w} \end{aligned}$$

$$i = 1 - \left(1 - \frac{1}{n}\right)\beta$$





Dissociation $\rightarrow [1 + (n-1)\alpha]$

$n = \text{Jitne logo mei toota}$
 $\alpha = \text{dod}$

Association $\rightarrow [1 - (1 - \frac{1}{n})\beta]$

$n = \text{Jitne log jude}$
 $\beta = \text{doa}$

ATDB.uno

Question

Molal depression constant for the solvent is 4 K kg mol^{-1} .



The depression in freezing point of the solvent for 0.03 mol/kg


solution of K_2SO_4 is (assuming complete dissociation).

A) 0.12 K

B) 0.36 K

C) 0.18 K

D) 0.24 K



$$\alpha = 1$$

$$\alpha = 1 + (3 - 1)$$

$$\checkmark \checkmark$$

$$i = 3$$

$$\Delta T_f = i K_f m = 3 \times 4 (0.03) = 0.36 \text{ K}$$



TELEGRAM GROUP BY FAISAL SIR



ATDB.uno

@FAISALSIRPW



THANK YOU

ATDB.uno