

श्री
Balaji

M.S. Chouhan


Advanced Problems in

ORGANIC CHEMISTRY

for

JEE

ATDB.uno



8th
edition

☐ Published by:
SHRI BALAJI PUBLICATIONS
(EDUCATIONAL PUBLISHERS & DISTRIBUTORS)

6, Gulshan Vihar,
Opp. Mahalaxmi Enclave,
Jansath Road, Muzaffarnagar (U.P.)
Phone : 0131-2660440 (O), 2600503 (R)
website : www.shribalajibooks.com
email : sbjpub@gmail.com

- ☐ First edition : 2008
- ☐ Second edition : April 2008
- ☐ Third edition : April 2009
- ☐ Seventh edition: April 2013
- ☐ Eighth edition : April 2014

☐ © All Right Reserved with Author

☐ Price : ₹ 480.00


☐ Typeset by :
Sun Creation
Muzaffarnagar

- ☐ All the right reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the authors and publisher. Any violation/breach shall be taken into legal action.

*Dedicated to the
Lotus feet of
Revered Guruji*

SHRI KESARAM JI MAHARAJ





Advanced Problems in
**ORGANIC
CHEMISTRY**

Preface

It is a matter of great pleasure for me to present the eighth edition of **"Advanced Problems in Organic Chemistry for JEE"** before JEE aspirants. During my teaching experience, I felt that the facts may be made more and more clear to the students through problematic approach. Although an ocean of material in Organic Chemistry is available with the students, yet the approach to design the problems has been changed in recent years and if one tries to swim in this ocean, it will be a very difficult task. To make the students more familiar with trends and tricks how to solve problems, the present problem book has been presented. In the current scenario of stiff competition especially for JEE, one must be clear that almost all the sincere applicants are well equipped with the facts of subject, yet the winner is one who knows how to use these equipments with accuracy and efficiency. As an experienced teacher, I would like to suggest students three golden rules to score high in Organic Chemistry:

1. Don't get behind
2. Work out a number of problems of different types
3. Revise through short notes / learning chart.

I hope that the present book will cater to the needs of JEE aspirants & as a matter of fact, they will enjoy the present venture and I would feel rewarded if this book is found helpful to the students and teachers in real terms. All attempts have been made to make the book error free however a few misprints may inadvertently creep.

I acknowledge the blessing and support of my mother Smt. Raj Kanwar, father Shri B.S. Chouhan, brother Dr. V.S. Chouhan, my wife and daughter. They inspired me all the time during the preparation of this book.

The support and valuable suggestions from my colleagues especially Mr. N. Avasthi, Mr. V.K. Jaiswal, Mr. Nitin Jain, Mr. N.K. Sethia, Mr. Vikash Gupta, Mr. Pankaj Joshi, Dr. S. Kothari, Mr. Vineet Khatri, Mr. Ashish Mishra, Mr. Manish Arora, Mr. Govind Khandelwal, Mr. Rahul Pareek, Mr. Rahul Malav, Mr. Akshay Choudhary, Mr. Hanuman Sahay, Mrs. Neha Joshi, Mrs. Neetu Jha, Mr. Kamlesh Gupta, Mr. Kumud Ranjan and Mr. K.D. Tiwari are highly acknowledged. I also pay my sincere thanks to all the esteemed members of **M/s Shri Balaji Publications** in bringing out this book in such a nice form.

In the last, constructive criticism and valuable suggestions from the readers are most welcome to make the book more useful.

M.S. CHOUHAN

email: mahen_chouhan@yahoo.com

Mob.(Only for SMS) : 09828025625

A few words to the JEE Aspirants

Dear JEE aspirants,

I hope that this collection of problems will surely help you during your preparation for JEE. In this book, each chapter consists of two levels :

Level 1 - includes the problems having only one option correct. These problems are based on different facts and their twists.

Level 2 - includes unique approach which may be used to solve the problems altogether different from the prevailing trend followed by JEE. These approaches will undoubtedly help you in the quick revision of the key facts and their applications.

I wish all of you a grand success in the ensuing Joint Entrance Examination. Your valuable suggestions and constructive criticism for the betterment of the book are welcome.

M.S. Chouhan


e-mail : mahen_chouhan@yahoo.com



Contents

1	General Organic Chemistry	1-70
2	Isomerism (Structural & Stereoisomerism)	71-144
3	Grignard Reagent	145-164
4A	Hydrocarbons (Alkanes)	165-180
4B	Hydrocarbons (Alkenes)	181-244
4C	Hydrocarbons (Alkynes)	245-250
5A	Alkyl Halides (Substitution Reaction)	251-284
5B	Alkyl Halides (Elimination Reaction)	285-314
5C	Alkyl Halides	315-327
6	Alcohol, Ether and Epoxide	328-372
7	Aldehydes and Ketones	373-423
8	Aldol and Cannizaro Reaction	424-439
9	Carboxylic Acid and their Derivatives	440-456
10	Amines	457-463
11	Carbene and Nitrene	464-482
12	Aromatic Compounds	483-552
13	Practical Organic Chemistry	553-557
14	Biomolecules	558-567
15	IUPAC Names	568-594
🎯	Learning Chart	

ATDB.uno




1


GENERAL ORGANIC CHEMISTRY

LEVEL - 1
ATDB.uno


1. How many 2° Hydrogen atoms are present in the given following compound ?







- (a) 2 (b) 5 (c) 7 (d) 8
2. Identify which functional group are **Not** present in the given following compound ?




- (a) Ketone (b) Ester (c) Amide (d) Ether
3. Correct energy profile for amine inversion and hybridization of nitrogen in transition state is:



2 ORGANIC Chemistry for IIT-JEE



4.



(i)



(ii)







(iii)



Compare the heats of combustion of above compounds:

- (a) (i) > (ii) > (iii) (b) (i) > (ii) > (iii) (c) (i) > (i) > (ii) (d) (ii) > (iii) > (i)

5. Which of the following is not a resonance structure of the others ?



6. Rank the hydrogen atoms (H_a , H_b , H_c) in the following molecule according to their acidic strength.

(a) $a > b > c$ (b) $b > a > c$ (c) $b > c > a$ (d) $c > b > a$ 

Compare the bond lengths a and b .


(a) $a = b$ (b) $b > a$ (c) $b < a$

(d) Impossible to predict

GENERAL ORGANIC CHEMISTRY


3

8. The number of $sp^2 - sp^2$ sigma bonds in the compound given below is :







- (a) 1 (b) 3 (c) 4 (d) 5

9. The total number of lone pair of electrons in the given molecule is :




- (a) 2 (b) 3 (c) 4 (d) 5

10. Which of the following rings is highly strained ?







11. The functional groups present in Cortisone are :






- (a) Ether, alkene, alcohol (b) Alcohol, ketone, alkene, ether
(c) Alcohol, ketone, amine (d) Ether, amine, ketone




12. Select the acid with the highest K_a (i. e., lowest pK_a).



13. Most stable carbocation among the given example is :



14. Set the following in increasing order of their pK_a values.



(a) $y < x < z$

(b) $x < y < z$


(c) $y < z < x$

(d) $x < z < y$


4

ORGANIC Chemistry for IIT-JEE

15. Which is the major product of the following reaction ?



16. In the given pair identify most acidic compound in (A) and (B). Most basic in (C) and (D).




(a) A - I, B - II, C - I, D - II

(c) A - II, B - II, C - II, D - II

(b) A - II, B - I, C - I, D - II
























(d) A - I, B - II, C - I, D - I

17. Several factors (steric, electronic, orbital interactions etc.) can affect the inversion barrier of an amine. In the given pair which data is correctly placed ?



GENERAL ORGANIC CHEMISTRY

5

25. Rank the following in decreasing order of the acidic strengths.
- (a)  (b)  (c)  (d) 
26. Which of the following carbonyl compound is most stable?
 $\Delta G^\ddagger = 7.9 \text{ kcal/mol}$ $\Delta G^\ddagger = 22.9 \text{ kcal/mol}$
- (a)  (b)  (c)  (d) 
18. Select the response that correctly identifies the number of carbon atoms of each type of hybridization in the compound given below.
- (a)  (b)  (c)  (d) 
19. Circle represents most basic atoms in these molecule. Which of the following is correct representation?
- (a)  (b)  (c)  (d) 
27. Selected bond angle in the following molecule are shown. Which of the following is correct?
- (a)  (b)  (c)  (d) 
28. Which of the following acids would have a STRONGER CONJUGATE BASE?
- (a)  (b)  (c)  (d) 
20. Circle represent most acidic hydrogens in these molecules. Which of the following is correct representation?
- (a)  (b)  (c)  (d) 
28. Which statement about the following equilibrium is correct?
- (a) The equilibrium favors the products.
 (b) t-Butyl is the dominant anion in the equilibrium.
 (c) The equilibrium is shifted by resonance.
 (d) All of these.
39. Consider the following reaction involving two acids shown below. Formic acid and HF are the products of the reaction.
- $\text{K}^+ \text{F}^- + \text{HCOOH} \rightleftharpoons \text{HCOO}^- \text{K}^+ + \text{HF}$

8


ORGANIC Chemistry for IIT-JEE

Which of the following statements about this reaction are true ?


- (A) Formic acid is the strongest Bronsted acid in the reaction
 (B) HF is the strongest Bronsted acid in the reaction
 (C) KF is the strongest Bronsted base in the reaction
 (D) KO_2CH is the strongest Bronsted base in the reaction
 (E) The equilibrium favours the reactants
 (F) The equilibrium favours the products
 (G) Formic acid has a weaker conjugate base
 (H) HF has a weaker conjugate base

(a) A, D and F (b) B, D, and H (c) A, C, and H (d) B, D, E and H


30. Which one of the following compounds has a dipole moment significantly different from zero?



31. Which one of the following has the smallest heat of combustion ?




32. Rank the following substances in order of decreasing heat of combustion (maximum $\rightarrow$ minimum).



- (a) $1 > 2 > 4 > 3$ (b) $3 > 4 > 2 > 1$
 (c) $2 > 4 > 1 > 3$ (d) $1 > 3 > 2 > 4$

33. Which of the following has lowest pK_a value ?




GENERAL ORGANIC CHEMISTRY

9


34. Arrange the following (w, x, y, z) in decreasing order of their boiling points:




(w)



(x)



(y)



(z)


(a) $w > x > z > y$

(b) $w > x > y > z$


(c) $w > z > y > x$

(d) $w > z > x > y$


35. Arrange the following in increasing order of their acidic strength.



(a) $\text{III} < \text{I} < \text{IV} < \text{II}$




(c) $\text{I} < \text{III} < \text{IV} < \text{II}$




(b) $\text{II} < \text{I} < \text{IV} < \text{III}$

(d) $\text{II} < \text{III} < \text{I} < \text{IV}$



36. How many degrees of unsaturation are there the following compound ?




(a) 6

(b) 7

(c) 10

(d) 11

37. The heat of hydrogenation for 3-methylbutene and 2-pentene are -30 kcal/mol and -28 kcal/mol respectively. The heats of combustion of 2-methylbutane and pentane are -784 kcal/mol and -782 kcal/mol respectively. All the values are given under standard conditions. Taking into account that combustion of both alkanes give the same products, what is ΔH (in kcal/mol) for the following reaction under same conditions ?




(a) 0

(b) -4

(c) -2

(d) 2

38. Which of the following σ -bonds participate in hyperconjugation ?



(a) I and II

(b) I and V


(c) II and V

(d) III and IV

10

ORGANIC Chemistry for IIT-JEE




39.



Decreasing order of acidic strength of different ($-\text{OH}$) groups is :





- (a) $w > x > y > z$ (b) $w > z > x > y$ (c) $z > w > x > y$ (d) $z > x > w > y$

40. Arrange the following alcohols in decreasing order of the ease of ionization under acidic conditions.






- (a) $\text{I} > \text{III} > \text{II}$ (b) $\text{I} > \text{II} > \text{III}$ (c) $\text{II} > \text{III} > \text{I}$ (d) $\text{II} > \text{I} > \text{III}$

41. Among the isomeric amines select the one with the lowest boiling point.







42. Which one of the compounds shown below, is not an isomer of the others ?



43. Arrange the anions (p) CH_3^- , (q) NH_2^- , (r) OH^- , (s) F^- , in decreasing order of their basic strength.

- (a) $p > q > r > s$ (b) $q > p > r > s$ (c) $r > q > p > s$ (d) $r > p > q > s$

44. One among the following compounds will not give effervescence with sodium carbonate:







45. The carboxylic acid which has maximum solubility in water is:

- (a) phthalic acid (b) succinic acid
(c) malonic acid (d) salicylic acid


GENERAL ORGANIC CHEMISTRY

11

46. Among the following compounds, the most basic compound is :



47.







Arrange the (C - H) bonds x, y and z in decreasing order of their bond dissociation energies in homolysis.

- (a) $y > x > z$ (b) $z > x > y$ (c) $z > y > x$ (d) $y > z > x$



48. 23 g of sodium will react with methyl alcohol to give :

- (a) one mole of oxygen (b) 22.4 dm^3 of hydrogen gas at NTP
(c) 1 mole of H_2 (d) 11.2 L of hydrogen gas at NTP

49. Which of the following is most polar?







50.







The correct order of decreasing basic strengths of x, y and z is :

- (a) $x > y > z$ (b) $x > z > y$ (c) $y > x > z$ (d) $y > z > x$

51. Which of the following is the strongest Bronsted acid ?



52. Which of the following is the strongest Bronsted base ?



53. Which of the following is polar aprotic solvent ?

- (a) DMSO (b) Crown ether (c) DMG (d) All of these

12


ORGANIC Chemistry for IIT-JEE




54. Some pairs of acids are given below. Select the pair in which second acid is stronger than first
- $\text{CH}_3\text{CO}_2\text{H}$ and $\text{CH}_2\text{FCO}_2\text{H}$
 - $\text{CH}_2\text{FCO}_2\text{H}$ and $\text{CH}_2\text{ClCO}_2\text{H}$
 - $\text{CH}_2\text{ClCO}_2\text{H}$ and $\text{CH}_2\text{BrCO}_2\text{H}$
 - $\text{CH}_3\text{CH}_2\text{CHFCO}_2\text{H}$ and $\text{CH}_3\text{CHFCH}_2\text{CO}_2\text{H}$

55. $\text{H}-\text{C} \equiv \text{C} \overset{a}{-} \text{C} \equiv \text{C} \overset{b}{-} \text{CH}_3$;

Compare the bond lengths a and b :




- $a = b$
 - $a > b$
 - $b > a$
 - $a \gg b$
56. Which (isomeric) amine has lowest boiling point ?
- 1° amine
 - 2° amine
 - 3° amine
 - cannot predict

57.  $\xrightarrow{2\text{SbCl}_5}$ P ; P will be :




(d) mixture of (a) and (b)

58. Which of the following substances is not an isomer of 3-ethyl 2-methyl pentane ?




(d) All are isomers

59. Which of the following is an isomer of compound 1 ?



- 2
- 4
- 2 and 3
- all are isomers

60.  $\xrightarrow{\text{AgNO}_3}$ (A) ;

Which statement is incorrect in respect of the above reaction ?





- Product is aromatic
- Product has high dipole moment
- Product has less resonance energy
- Product is soluble in polar solvent

GENERAL ORGANIC CHEMISTRY





Multiple Choice Questions

13

61. Some pairs of ions are given below. In which pair, first ion is more stable than second?



62. Among the given pairs in which pair, first compound has higher boiling point than second?



63. Which of the following alcohol is the least soluble in water?

(a) Ethanol

(b) 1-Propanol

(c) 1-Butanol

(d) 1-Pentanol

64. Which of the following alcohols is expected to have a lowest pK_a value?





(a) Ethanol

(b) 1-propanol


(c) 2, 2, 2-trifluoroethanol

(d) 2-chloroethanol

65. Which of the following alkenes is the most stable?



66. Bond X is made by the overlap of which type of hybridized orbitals?







(a) sp and sp^3

(b) sp and sp^2

(c) sp^2 and sp^3

(d) none of these

67. Increasing order of acidic strength of given compounds is :



14

ORGANIC Chemistry for IIT-JEE

(a) III < I < IV < II

(b) II < I < IV < III

(c) I < III < IV < II

(d) I < III < II < IV



68. -COOH + NaHCO₃ $\xrightarrow{*}$ CO₂ + -COONa, $\bar{C}^*$ is with the product :

(a) CO₂(b) -COONa

(c) both

(d) none of these

69. Rank in the order of increasing acidity.



(a) III < I < II

(b) I < III < II


(c) III < II < I

(d) II < I < III

70. Which compound has the highest value of pK_a?

(a) Cl-CH₂-CH₂-COOH(b) CH₃-CH₂-COOH(c) CH₃-CH(Cl)-COOH(d) CH₃-C(Cl)(CH₃)-COOH

71. Consider the hydrogen atoms attached to three different carbon atoms (labeled 1, 2 & 3). Rank the attached hydrogen atoms in order from most acidic to least acidic.






(a) 2 > 1 > 3

(b) 1 > 2 > 3

(c) 2 > 3 > 1

(d) 3 > 2 > 1

72. Decreasing order of acidic strengths of following compounds is :



(a) x > y > z



(b) y > x > z



(c) z > y > x

(d) z > x > y

73. Among the given pairs, in which pair second compound is more acidic than first ?

(a) BrCH₂NO₂ and CH₃CH₃(b) CH₃-C(=O)CH₂CN and CH₃-C(=O)CH₃

(c)  & 
CH₃-C(=O)

(d)  & 

GENERAL ORGANIC CHEMISTRY

15


74. Which of the underlined atoms in the molecules shown below have sp -hybridization ?

- (u) CH_2CHCH_3 (v) CH_2CCHCl (w) CH_3CH_2^+ (x) $\text{H}-\text{C}\equiv\text{C}-\text{H}$
 (y) CH_3CN (z) $(\text{CH}_3)_2\text{C}=\text{N}-\text{NH}_2$
 (a) x and z (b) x, y, and z (c) u, w and x (d) v, x and y

75. Which of the following, is the product of the reaction between AlCl_3 and CH_3OCH_3 ?


- (a) $\text{Cl}-\text{Al}(\text{Cl})_2-\text{O}^+(\text{CH}_3)_2$
 (b) $\text{Cl}-\text{Al}^+(\text{Cl})_2-\text{O}^-(\text{CH}_3)_2$
 (c) $\text{Cl}-\text{Al}^+(\text{Cl})_2-\text{O}^-(\text{CH}_3)_2$
 (d) $\text{Cl}-\text{Al}(\text{Cl})_2-\text{CH}_3\text{OCH}_3$

76. Which of the following compounds contain at least one secondary alcohol ?




- (a) I, II, IV, VI (b) I, II (c) I, II, IV, V (d) I, III, V

77. Which of the following has the most negative heat of cycloaddition ?




78. Which of the following options is the correct order of relative stabilities of cations I, II and III as written below (most stable first) ?



- (a) I > II > III (b) II > III > I (c) III > I > II (d) I > III > II

79. What is the decreasing order of stability (most stable $\rightarrow$ least stable) of the following carbocations ?




- (a) 3 > 2 > 1 > 4 > 5 (b) 3 > 2 > 5 > 4 > 1
 (c) 1 $\approx$ 4 > 2 $\approx$ 5 > 3 (d) 3 > 1 $\approx$ 4 > 2 $\approx$ 5

16

ORGANIC Chemistry for IIT-JEE


80.



the hydrogen indicated by arrow will be easily removed as :


- (a) H^+ (b) $H^\ominus$ (c) $H^\bullet$ (d) H^{-2}

81. Rank the bond dissociation energies of the bonds indicated with the arrows. (from smallest to largest).




- (a) $1 < 2 < 3$ (b) $3 < 2 < 1$ (c) $2 < 3 < 1$ (d) $3 < 1 < 2$

82. Rank the following compounds in order of decreasing acid strength (most acidic $\rightarrow$ least acidic) .




- (a) $2 > 4 > 1 > 3$ (b) $1 > 2 > 3 > 4$ (c) $3 > 1 > 2 > 4$ (d) $3 > 1 > 4 > 2$

83. Rank the following compounds in order of increasing acidity (weakest acid first).



- (a) $2 < 3 < 1$ (b) $3 < 1 < 2$ (c) $1 < 2 < 3$ (d) $2 < 1 < 3$

84. Which of the following phenols has the largest pK_a value (i.e., is least acidic) ?



85. Among the given sets, which represents the resonating structures ?

- (a) $H - C \equiv \overset{+}{N} - \ddot{O}^-$ and $H - \ddot{O} - C \equiv N:$
 (b) $H - \overset{+}{O} = C = \ddot{N}^-$ and $H - \ddot{O} - C \equiv N:$
 (c) $H - C \equiv \overset{+}{N} - \ddot{O}^-$ and $H - C \equiv \ddot{O} - \ddot{N}:$
 (d) $H - \ddot{O} - C \equiv N^-$ and $H - \ddot{N} = C = \ddot{O}:$


GENERAL ORGANIC CHEMISTRY

17

86. Identify each species in the following equilibrium according to the code :

SA = stronger acid ; SB = stronger base ; WA = weaker acid ; WB = weaker base.

The pK_a of $(CH_3)_2NH$ is 36 ; the pK_a of CH_3OH is 15.2.



(a) WA WB

(b) WB WA

(c) SA SB

(d) SB SA

(e) WA WA

87. The hydrogen bonding is strongest in which one of the following set ?

(a) $F - H \cdots F$

(b) $O - H \cdots S$

(c) $S - H \cdots F$

(d) $F - H \cdots O$


88. Intermolecular hydrogen bonding is strongest in :

(a) methylamine

(b) phenol

(c) formaldehyde

(d) methanol



Identify most acidic hydrogen in given compound



(a) a

(b) b

(c) c

(d) d





90. Which of the following compounds would you expect to be strongest carbon acid ?



(c) $CH_2(CO_2Et)_2$

(d) $CH_3COCH_2COOC_2H_5$

91. 5-Bromo-2-cyclopropyl cyclohex-2-enol have correct structure is:





18

ORGANIC Chemistry for IIT-JEE


92. Rearrange the following in the increasing order of acidic strength:
- (i) benzoic acid (ii) *p*-methoxybenzoic acid (iii) *o*-methoxybenzoic acid
- (a) i < ii < iii (b) iii < i < ii (c) ii < i < iii (d) iii < ii < i

93. In the following acid-base reaction, in which can backward reaction if favoured?



94. Which compound posses highest dipole moment ?


- (a) naphthalene (b) phenanthrene
(c) anthracene (d) azulene



(E = activation energy)


Relation between activation energies of above reactions is :

- (a) $E_2 > E_1 > E_3$ (b) $E_3 > E_1 > E_2$ (c) $E_3 > E_2 > E_1$ (d) $E_1 > E_2 > E_3$
96. Rank the following solvents in decreasing order of ability to dissolve given compound.



- (a) $\text{Et}_2\text{O} > \text{H}_2\text{O} > \text{EtOH}$ (b) $\text{H}_2\text{O} > \text{EtOH} > \text{Et}_2\text{O}$
(c) $\text{H}_2\text{O} > \text{Et}_2\text{O} > \text{EtOH}$ (d) $\text{Et}_2\text{O} > \text{EtOH} > \text{H}_2\text{O}$

97.



Its basic strength is 10^{10} more than 1-dimethyl amino naphthalene.





Reason for high basic strength is :

- (a) resonance (b) steric inhibition of resonance
(c) ortho effect (d) hyperconjugation



GENERAL ORGANIC CHEMISTRY

19

98. In the given pair of compounds, in which pair second compound has higher boiling point than first compound ?



99.






; Dipole moments of given compound will be :

- (a) (A) = 6.87D, (B) = 4.11D (b) (A) = 4.11 D, (B) = 6.87 D
(c) (A) = 4.11 D, (B) = 4.11 D (d) (A) = 6.87 D, (B) = 6.87 D

100. Order of decreasing basic strengths of halides is :

- (a) $\text{F}^- > \text{Cl}^- > \text{I}^- > \text{Br}^-$ (b) $\text{F}^- > \text{Cl}^- > \text{Br}^- > \text{I}^-$
(c) $\text{I}^- > \text{Br}^- > \text{Cl}^- > \text{F}^-$ (d) $\text{I}^- > \text{Cl}^- > \text{Br}^- > \text{F}^-$

101. Among the xylenes, which is thermodynamically most stable ?



(d) All are equally stable

102. Heat of combustion of two isomer x and y are 17 kJ/mol and 12 kJ/mol respectively. From this information it may be concluded that :


- (a) isomer x is 5 kJ/mol more stable
(b) isomer y is 5 kJ/mol less stable
(c) isomer y has 5 kJ/mol more potential energy
(d) isomer x is 5 kJ/mol less stable

20 ORGANIC Chemistry for IIT-JEE


103. Rank the following substances in decreasing order of heat of combustion (most exothermic $\rightarrow$ least exothermic)



(A)




(B)




(C)

(a) $B > A > C$ (b) $A > B > C$ (c) $C > A > B$ (d) $C > B > A$


104.




(1)



(2)



(3)



(4)

Choose the statement that best describes given compounds.





(a) 1, 3, 4 represent same compound

(b) 1 and 3 are isomer of 2 and 4



(c) 1, 4 are isomer of 2 and 3

(d) All the structure represent the same compound


105. Decreasing order of acid strengths is :

(a) $B > A > C > D$ (b) $C > A > B > D$ (c) $C > A > D > B$ (d) $C > B > A > D$


106. Rank the following in decreasing order of basic strength is :

(a) $B > A > D > C$ (b) $D > A > B > C$ (c) $A > D > B > C$ (d) $A > D > C > B$


107. Among the given compound choose the two that yield same carbocation on ionization.




(A)



(B)



(C)



(D)

(a) A, C

(b) B, D

(c) A, B

(d) B, C

108. Oxalic acid

 pK_1

Malonic acid

 pK_2

Heptanedioic acid

 pK_3





where pK_1, pK_2, pK_3 are first ionization constants. Incorrect order is :

(a) $pK_1 > pK_2 > pK_3$ (b) $pK_1 < pK_2 < pK_3$ (c) $pK_3 > pK_2 > pK_1$ (d) $pK_3 > pK_1 > pK_2$





GENERAL ORGANIC CHEMISTRY

21





109. In sets a – d, only one of the set is incorrect regarding basic strength. Select it :




110. Dipole moment of which ketone is maximum ?



111. Correct order of basic strengths of given amines is :



112. Order of basic strength $\text{Ph} - \text{NH}_2$, $\text{Ph} - \text{NH} - \text{Me}$, $\text{Ph} - \text{N}(\text{Me})_2$, 

(A)

(B)

(C)

(D)




22

ORGANIC Chemistry for IIT-JEE

113. Carbon-carbon double bond length will be maximum in which of the following compounds ?

- (a) $\text{CH}_3 - \text{CH} = \text{CH}_2$ (b) $\text{CH}_3 - \text{CH} = \text{CH} - \text{CH}_3$
 (c) $\text{CH}_3 - \underset{\text{CH}_3}{\text{C}} = \underset{\text{CH}_3}{\text{C}} - \text{CH}_3$ (d) $\text{CH}_2 = \text{CH}_2$


114. Which has maximum dipole moment ?




Compare the basic strengths of compounds given:

- (a) (i) > (ii) > (iii) (b) (ii) > (i) > (iii)
 (c) (ii) > (iii) > (i) (d) (iii) > (ii) > (i)

116. For the following two reactions, which statement is true ?




- (a) Reaction A is faster and less exergonic than B
 (b) Reaction B is faster and more exergonic than A
 (c) Reaction A is faster and less endergonic than B
 (d) Reaction B is faster and more endergonic than A



Identify the site, where attack of H^+ is most favourable.

- (a) a (b) b (c) c (d) d

118. Rank the following alkenes on order of increasing λ_{max} .







- (a) $1 < 2 < 3$ (b) $1 < 3 < 2$ (c) $2 < 1 < 3$ (d) $2 < 3 < 1$


GENERAL ORGANIC CHEMISTRY

23


119. Which of the following cyclic amine has lowest $\Delta G^\ddagger$ for inversion ?




120. Rank in the order of increasing acidic strength:



(A)



(B)



(C)





(a) A < B < C

(b) A < C < B




(c) B < A < C

(d) B < C < A

121. Which one of the following dienes would you expect to be the most stable ?



122. Which metal catalyzed reaction would release the maximum amount of heat per CH₂ unit ?

(a) cyclopropane + H₂ → propane(b) cyclobutane + H₂ → butane(c) cyclopentane + H₂ → pentane(d) cyclohexane + H₂ → hexane

Compare basic strengths of the above compounds:






(a) A > B > C

(b) B > A > C

(c) C > A > B

(d) C > B > A


124. On reaction with acid, 4-pyrone gives a very stable cationic product. Which of the following structures shows the protonation site in that product ?



24

ORGANIC Chemistry for IIT-JEE

125. Which of the following is the most stabilized carbocation ?




126. Which carbocation is the most stable ?




127. Consider a positively charged C_2H_3 species in which the positively charged carbon is sp -hybridized, the uncharged carbon is sp^2 -hybridized and an empty p -orbital is perpendicular to the π system. What is the best description of this cation ?

- (a) vinyl (b) allenyl (c) alkyl (d) allyl


128. Which of the following reactions is not exothermic ?



129. List the following carbocation in order of decreasing stabilization energies.




130. For the following two acid-base reactions, which statement is true ?



- (a) I is favoured to the right, II is favoured to the left
 (b) I is favoured to the left, II is favoured to the right
 (c) I is favoured to the right, II is favoured to the right
 (d) I is favoured to the left, II is favoured to the left

131. Rank the hydrogen atoms (H_a , H_b , H_c) in the following molecules according to their acidic strengths:







- (a) $a > b > c$ (b) $b > a > c$ (c) $b > c > a$ (d) $a > c > b$





GENERAL ORGANIC CHEMISTRY

25


132. In which of the following reactions, backward reaction is favoured ?



133. Which carbocation is the most stabilized ?



134. Taking into account of hybridization and resonance effects, rank the following bonds in order of decreasing bond length.







(a) $\text{I} > \text{II} = \text{III}$

(b) $\text{II} > \text{III} > \text{I}$




(c) $\text{I} > \text{III} > \text{II}$

(d) $\text{II} = \text{III} = \text{I}$

135. Which one among the following carbocations has the longest half-life ?



136. Rank the following alkenes in order of decreasing heats of hydrogenation (largest first)



(a) $2 > 3 > 4 > 1$

(c) $1 > 3 > 4 > 2$





(b) $2 > 4 > 3 > 1$

(d) $1 > 4 > 3 > 2$






26

ORGANIC Chemistry for IIT-JEE





137. Which of the following reactions is most exothermic ?







138. Which energy diagram best represents the given reaction ?






139. Which one of the following is most stable ?



140. Which of the following is most stable ?



141. Compare relative stability of the following resonating structure.






(a) (i) > (ii) > (iii)

(b) (ii) > (i) > (iii)

(c) (i) > (iii) > (ii)

(d) (ii) > (iii) > (i)

142. Compare relative stability of the following resonating structure.




GENERAL ORGANIC CHEMISTRY


27




- (a) $p > q > r$
(c) $q > r > p$
(b) $q > p > r$
(d) $p > r > q$
143. Which of the following isomeric hydrocarbons is most acidic?
144. Which of the following orders is correct for heat of hydrogenation of these compounds?
145. Use the following data to answer the question below.
146. Calculate the resonance energy of anthracene.
147. How many double bond equivalents does amoxicillin (shown below) possess?
148. What is the oxidation state of osmium in 7B and 7C, respectively?
149. The most stable canonical structure among the given structure is

30


160. 

149. For the given compounds the correct order of resonance energy is :
(a) $\text{III} > \text{I} > \text{II}$ (b) $\text{II} > \text{I} > \text{III}$
(c) $\text{I} > \text{II} > \text{III}$ (d) $\text{III} > \text{II} > \text{I}$


161. 

150. Which of the following CH_3 groups has acidic hydrogen?
(a) 
(b) 
(c) 
(d) 


The correct stability order of the given canonical structures is :
(a) $\text{I} > \text{II} > \text{III} > \text{IV}$
(b) $\text{III} > \text{I} > \text{II} > \text{IV}$
(c) $\text{I} > \text{III} > \text{II}$ (d) $\text{II} > \text{III} > \text{I}$


162. 

151. In the above compound, how many sites are available for the attack of CH_3O^- ?
(a) 1 (b) 2 (c) 3 (d) 4



163. 

Which of the following orders of rotation barrier about the $\text{C} \equiv \text{C}$ bond, as indicated, is correct?
(a) $\text{I} > \text{II} > \text{III}$
(b) $\text{III} > \text{II} > \text{I}$
(c) $\text{III} > \text{I} > \text{II}$ (d) $\text{II} > \text{I} > \text{III}$

152. 

164. 

153. Which of the following orders of rotation barrier about the $\text{C} = \text{C}$ bond, as indicated, is correct?
(a) $\text{I} > \text{II} > \text{III}$
(b) $\text{III} > \text{II} > \text{I}$
(c) $\text{III} > \text{I} > \text{II}$ (d) $\text{II} > \text{I} > \text{III}$

165. Which of the following compound is not resonance stabilized?
(a) 
(b) 
(c) 
(d) 





GENERAL ORGANIC CHEMISTRY

31

166. Homologous compound have same:

- (a) General formula (b) Empirical formula
(c) Structural formula (d) Molecular formula

167. Most acidic is:




168. Which of the following substituents will decrease the acidic strength of phenol?

- (a) $-\text{NO}_2$ (b) $-\text{CN}$
(c) $-\text{CH}_3$ (d) $-\text{CHO}$

169. Which of the following structures possesses a cross-conjugated system?

- (a) $\text{CH}_2 = \text{CH} - \text{CH} = \text{CH} - \text{CH}_2$
- (c) $\begin{array}{c} \text{CH}_2 = \text{CH} - \text{CH} - \text{CH} = \text{CH}_2 \\ | \\ \text{CH} = \text{CH}_2 \end{array}$
- (b) $\begin{array}{c} \text{CH}_2 = \text{CH} - \text{C} = \text{CH}_2 \\ | \\ \text{CH}_2\text{CH}_3 \end{array}$
- (d) $\begin{array}{c} \text{CH}_2 = \text{CH} - \text{C} = \text{CH}_2 \\ | \\ \text{CH} = \text{CH}_2 \end{array}$

170. Examine the following resonating structures of formic acid for their individual stability and then answer the question given below.




Which of the following arrangements gives the correct order of decreasing stability of the above-mentioned resonance contributors?

- (a) $\text{II} > \text{I} > \text{III} > \text{IV}$
(b) $\text{I} > \text{II} > \text{III} > \text{IV}$
(c) $\text{IV} > \text{III} > \text{I} > \text{II}$
(d) $\text{IV} > \text{III} > \text{I} > \text{II}$

171. Which of the following is not resonating structure of each other?

- (a) $\text{CH}_3 - \text{N} = \text{C} = \text{S}$ and $\text{CH}_3 - \text{S} - \text{C} \equiv \text{N}$
 (b) $\text{CH}_3 - \overset{+}{\text{C}} = \text{O}$ and $\text{CH}_3 - \text{C} \equiv \overset{+}{\text{O}}$



- (d) $\text{CH}_2 = \text{CH} - \text{C} \equiv \text{N}$ and $\overset{+}{\text{CH}}_2 - \text{CH} = \text{C} = \text{N}^-$


172. In the molecule $\text{CH}_3\text{C} \equiv \text{CCH} = \text{CH}_2$, the maximum number of carbon atoms arranged linearly is:

- [illegible]

32

ORGANIC Chemistry for IIT-JEE

173. The stability order of the following carbocations is:




(a) $\text{II} > \text{IV} > \text{III} > \text{I}$

(b) $\text{IV} > \text{II} > \text{III} > \text{I}$

(c) $\text{II} > \text{III} > \text{I} > \text{IV}$

(d) $\text{I} > \text{III} > \text{II} > \text{IV}$

174. Total number of α -hydrogen in given compound is:




(a) 4

(b) 5


(c) 6

(d) 7


175. In which pair second ion is more stable than first?




(i)



(ii)



(iii)



(iv)

(a) (i) and (ii)

(b) (ii) and (iii)

(c) (ii) and (iv)

(d) (iii) and (iv)

GENERAL ORGANIC CHEMISTRY



33

ANSWERS — LEVEL 1


1.	(c)	2.	(d)	3.	(d)	4.	(a)	5.	(d)	6.	(d)	7.	(b)	8.	(c)
9.	(b)	10.	(c)	11.	(b)	12.	(b)	13.	(a)	14.	(b)	15.	(b)	16.	(b)
17.	(d)	18.	(c)	19.	(d)	20.	(d)	21.	(c)	22.	(c)	23.	(c)	24.	(a)
25.	(d)	26.	(d)	27.	(d)	28.	(a)	29.	(d)	30.	(a)	31.	(c)	32.	(c)
33.	(d)	34.	(d)	35.	(d)	36.	(d)	37.	(b)	38.	(b)	39.	(a)	40.	(c)
41.	(c)	42.	(d)	43.	(a)	44.	(c)	45.	(c)	46.	(d)	47.	(b)	48.	(d)
49.	(b)	50.	(b)	51.	(d)	52.	(d)	53.	(d)	54.	(a)	55.	(c)	56.	(c)
57.	(b)	58.	(b)	59.	(d)	60.	(c)	61.	(b)	62.	(b)	63.	(d)	64.	(c)
65.	(d)	66.	(c)	67.	(a)	68.	(a)	69.	(d)	70.	(b)	71.	(a)	72.	(d)
73.	(d)	74.	(d)	75.	(a)	76.	(d)	77.	(a)	78.	(a)	79.	(b)	80.	(a)
81.	(d)	82.	(d)	83.	(d)	84.	(c)	85.	(b)	86.	(a)	87.	(a)	88.	(b)
89.	(a)	90.	(d)	91.	(b)	92.	(c)	93.	(d)	94.	(d)	95.	(d)	96.	(d)
97.	(b)	98.	(d)	99.	(a)	100.	(b)	101.	(b)	102.	(d)	103.	(a)	104.	(a)
105.	(c)	106.	(c)	107.	(c)	108.	(b)	109.	(c)	110.	(c)	111.	(d)	112.	(c)
113.	(c)	114.	(c)	115.	(c)	116.	(a)	117.	(a)	118.	(d)	119.	(c)	120.	(c)
121.	(c)	122.	(a)	123.	(c)	124.	(c)	125.	(d)	126.	(b)	127.	(a)	128.	(d)
129.	(b)	130.	(a)	131.	(c)	132.	(d)	133.	(c)	134.	(d)	135.	(a)	136.	(d)
137.	(c)	138.	(d)	139.	(c)	140.	(c)	141.	(a)	142.	(d)	143.	(b)	144.	(c)
145.	(a)	146.	(b)	147.	(d)	148.	(b)	149.	(a)	150.	(b)	151.	(d)	152.	(b)
153.	(a)	154.	(a)	155.	(c)	156.	(c)	157.	(c)	158.	(b)	159.	(c)	160.	(c)
161.	(b)	162.	(c)	163.	(a)	164.	(a)	165.	(c)	166.	(a)	167.	(d)	168.	(c)
169.	(d)	170.	(b)	171.	(a)	172.	(d)	173.	(c)	174.	(c)	175.	(b)		

34

ORGANIC Chemistry for IIT-JEE


LEVEL-2

1. Rank in order of radical stability (1 = most stable).



2. Predict the acidity order for the three phenols shown below :






Acidity order : 1 (most) to 3 (least)



Acidity order :

.....

.....

3. Comprehension

GENERAL ORGANIC CHEMISTRY

35

A. Which of the phenol derivatives above is the strongest acid ?

- ☐ Compound A ☐ Compound B ☐ Compound C
☐ Compound D ☐ Compound E ☐ Compound F

B. Which of the phenol derivatives above is the weakest acid ?

- ☐ Compound A ☐ Compound B ☐ Compound C
☐ Compound D ☐ Compound E ☐ Compound F









C. Which of the mono-nitrophenol derivatives above is the strongest acid ?

- ☐ Compound A ☐ Compound D

D. Which of the carbon-substituted phenol derivatives above is the strongest acid ?

- ☐ Compound C ☐ Compound F

4. The following questions refer to the twelve compounds given below. You may enter as many as six choices in each answer box.





(a)		(b)		(c)	$\text{H}_3\text{C}-\text{C}(=\text{O})-\text{O}-\text{C}_2\text{H}_5$
(d)	$\text{H}-\text{F}$	(e)		(f)	
(g)		(h)		(i)	
(j)		(k)		(l)	

A. Which compound may serve only as H-bond acceptors ?

B. Which may serve both as H-bond donors and acceptors?

C. Which compounds will not participate in H-bonding ?

5. Consider the following compounds and answer A and B.

(I) 	(II) 	(III) 	(IV) 

A. Which of the compounds is the strongest Bronsted acid ?

(a) I

(b) II

(c) III

(d) IV

38

ORGANIC Chemistry for IIT-JEE

10. In the two questions below, you are asked to rank the relative strengths of illustrated acids and bases. Use your knowledge of resonance and inductive to answer this.

A. For the series of bases shown below, rank the set from strongest to weakest.

(i)					Strongest
	(a)	(b)	(c)	(d)	Weakest
(ii)					Strongest
	(a)	(b)	(c)	(d)	Weakest

B. For the series of acids shown below, rank the set from strongest to weakest.





(i)					Strongest
	(a)	(b)	(c)	(d)	Weakest
(ii)					Strongest
	(a)	(b)	(c)	(d)	Weakest

11. In each of the following sections four compounds are listed. (Decreasing order of acidic strength, 1 is strongest & 4 is weakest).

(a)	$\text{CH}_2(\text{CO}_2\text{C}_2\text{H}_5)_2$	$\text{CH}_3\text{COCH}_2\text{CO}_2\text{C}_2\text{H}_5$	$(\text{CH}_3\text{CO})_2\text{CH}_2$	$\text{RC}\equiv\text{CH}$
(b)	RCH_2NO_2	RSO_2CH_3	$(\text{C}_6\text{H}_5)_3\text{CH}$	RCOCH_3
(c)	$\text{CH}_2(\text{C}\equiv\text{N})_2$	$\text{CH}_2(\text{NO}_2)_2$	$\text{HC}\equiv\text{N}$	$\text{RCH}_2\text{CO}_2\text{C}_2\text{H}_5$

GENERAL ORGANIC CHEMISTRY

39

(d)				

12. Rank in the order of increasing basic strength.

A.

(a)

NC1CCC(=O)N1

(b)

CC1CCC(=O)N1

(c)

CC1CCC(=O)N1

B.

(a)

c1ccncc1

(b)

c1ccc(cc1)N2CCCC2

(c)

C1CCNCC1

(d)

C1CCNCC1

C.

(a)

NC(=N)N

(b)

CNC(=N)N






(c)

CNC(=O)N

(d)

CNC(=O)C

13. Compare acidic strength of the following (Write your answer in box).


A.					
	(a)	(b)	(c)		
B.					
	(a)	(b)	(c)		

40

ORGANIC Chemistry for IIT-JEE

C.						
	(a)	(b)	(c)			
D.						
	(a)	(b)	(c)	(d)		
E.						
	(a)	(b)	(c)			
F.						
	(a)	(b)	(c)			



14. Arrange the hydrogens in increasing order of their acidic strengths.



GENERAL ORGANIC CHEMISTRY

41

15. The compounds whose structures are shown below, incorporate a variety of functional groups. The question on the right ask you to identify which compounds have a specific functional group. For each compound that has the designed group, enter the appropriate number. The aromatic rings should not be counted as double bonds.


















A.	Which have carbon-carbon double bonds ?	
B.	Which have a ketone carbonyl group ?	
C.	Which have an aldehyde carbonyl group ?	
D.	Which have aromatic rings ?	
E.	Which have a hydroxy group ?	
F.	Which have ether groups ?	
G.	Which have an ester group ?	
H.	Which have an amide group ?	
I.	Which have a carboxylic acid group ?	

42

ORGANIC Chemistry for IIT-JEE

16.

Problem	A	B	C	D
1				
2				
3				
4				

A. Which is the strongest acid in 1?

- (a) A (b) B (c) C (d) D

B. Which is weakest acid in 1?

- (a) A (b) B (c) C (d) D

C. Which is the strongest acid in 2?

- (a) A (b) B (c) C (d) D

D. Which is weakest acid in 2?

- (a) A (b) B (c) C (d) D

E. Which is the strongest acid in 3?

- (a) A (b) B (c) C (d) D

F. Which is weakest acid in 3?

- (a) A (b) B (c) C (d) D

G. Which is the strongest acid in 4?

- (a) A (b) B (c) C (d) D







H. Which is weakest acid in 4?

- (a) A (b) B (c) C (d) D

GENERAL ORGANIC CHEMISTRY

43

17. For each of the six structural formulae (A through F), shown below, five questions are posed. The answer to each is a number that should be entered in the appropriate answer box.



 A	 B	 C
 D	 E	 F

- A.** (i) Number of sp^3 carbons :
(ii) Number of sp^2 carbons :
(iii) Number of sp carbons :
(iv) Number of carbon-carbon σ -bonds :
(v) Number of π -bonds to carbon :
- B.** (i) Number of sp^3 carbons :
(ii) Number of sp^2 carbons :
(iii) Number of sp carbons :
(iv) Number of carbon-carbon σ -bonds :
(v) Number of π -bonds to carbon :
- C.** (i) Number of sp^3 carbons :
(ii) Number of sp^2 carbons :
(iii) Number of sp carbons :
(iv) Number of carbon-carbon σ -bonds :
(v) Number of π -bonds to carbon :
- D.** (i) Number of sp^3 carbons :
(ii) Number of sp^2 carbons :
(iii) Number of sp carbons :
(iv) Number of carbon-carbon σ -bonds :
(v) Number of π -bonds to carbon :
- E.** (i) Number of sp^3 carbons :
(ii) Number of sp^2 carbons :
(iii) Number of sp carbons :
(iv) Number of carbon-carbon σ -bonds :
(v) Number of π -bonds to carbon :
- F.** (i) Number of sp^3 carbons :
(ii) Number of sp^2 carbons :
(iii) Number of sp carbons :
(iv) Number of carbon-carbon σ -bonds :
(v) Number of π -bonds to carbon :





44

ORGANIC Chemistry for IIT-JEE

18. Match the column (I) and (II). (Matrix)

Column (I)		Column (II)	
	Molecule		Property
(a)		(p)	cis-compound
(b)		(q)	trans-compound
(c)		(r)	Highest heat of combustion
(d)		(s)	lowest heat of combustion


19. Match the column (I) and (II).

Column (I)		Column (II)	
	Molecule		pK_a of Conjugate acid
(a)		(p)	0.8
(b)		(q)	5.33
(c)		(r)	10.65
(d)		(s)	10.95

GENERAL ORGANIC CHEMISTRY

45

20. The junctures centred on atoms A, B and C on the given structure.



- A. Which juncture has the greatest deviation from planarity ?
 (a) A (b) B (c) C (d) Cannot be predicted
- B. Of the carbon-carbon bonds, (shown above) numbered from 1 to 4, which represent the most favourable site for H_2 addition ?
 (a) 1 (b) 2 (c) 3 (d) 4

21. Select the most stable structure in each of the following

Part (A)			
(a)	(b)	(c)	(d)
Part (B)			
(a)	(b)	(c)	(d)
Part (C)			
(a) $H_2C = CH - CH = CH - CH_3$	(b) $H_2C = C = CH - CH_2 - CH_3$		
(c) $H_3C - CH = C = CH - CH_3$	(d) $H_2C = CH - CH_2 - CH = CH_2$		

22. Match the column I and II. (Matrix)

Column (I)		Column (II)	
(a)	$-NO_2$	(p)	$-m$ effect
(b)	$-O^-$	(q)	$+m$ effect
(c)	$-O - CH_3$	(r)	$+I$ effect
(d)	$-C \equiv N$	(s)	$-I$ effect

23. Match the column I and II. (Matrix)


Column (I)		Column (II)	
(a)	$H_3C - CH = CH - CH_3$	(p)	Dipole (<i>cis</i> > <i>trans</i>)
(b)	$H_3C - CH = CH - CN$	(q)	Dipole (<i>trans</i> > <i>cis</i>)
(c)	$H_3C - CH = CH - Cl$	(r)	Melting point (<i>trans</i> > <i>cis</i>)
(d)	$Cl - CH = CH - Cl$	(s)	Boiling point (<i>cis</i> > <i>trans</i>)

46


ORGANIC Chemistry for IIT-JEE

24. Identify the most stable structure in each of the following :


Part (A)




Part (B)




Part (C)




Part (D)



Part (E)




Part (F)




GENERAL ORGANIC CHEMISTRY

47



25. Identify the most basic compound in the following.



48

ORGANIC Chemistry for IIT-JEE

26. Identify the most acidic hydrogen containing compound from the following.




Part (A)			
(a)	(b)	(c)	(d)
Part (B)			
(a)	(b)	(c)	(d)
Part (C)			
(a)	(b)	(c)	(d)
Part (D)			
(a)	(b)	(c)	(d)
Part (E)			
(a)	(b)	(c)	(d)

GENERAL ORGANIC CHEMISTRY





64 65 66 67 68 69

49





Part (F)







Part (G)




Part (H)



Part (I)



27.



Give the type of hybridization present at each atom.

(i) N —.....

(ii) C₁ —.....

(iii) C₂ —.....

(iv) O_1 —.....


(v) CH₂ —.....

(vi) C₃ —.....

50

ORGANIC Chemistry for IIT-JEE

28. Predict the direction of the following equilibrium. Write your answer in the box given below.



(A)





(B)

(C)

(D)



29. Match the column I and II. (Matrix)

Column (I)		Column (II)	
(a)	NaHCO ₃ will react with	(p)	 Squaric acid
(b)	Na will react with	(q)	
(c)	NaOH will react with	(r)	
(d)	NaNH ₂ will react with	(s)	





30. Match the column I and II.

Column (I)		Column (II)	
Acid		pK _a	
(a)	CH ₃ — CO ₂ H	(p)	5.69
(b)	(CH ₃) ₃ N ⁺ CH ₂ CO ₂ H	(q)	4.27
(c)	(CH ₃) ₃ N ⁺ (CH ₂) ₄ CO ₂ H	(r)	1.83
(d)	Ö ₂ C — CH ₂ — CO ₂ H	(s)	4.80


GENERAL ORGANIC CHEMISTRY

51

31. Match the column I and II.

Column (I)		Column (II)	
(a)	 $\text{C}^{14}\text{O}-\text{O}-\text{H} + \text{NaHCO}_3 \rightarrow$	(p)	NH_3
(b)	 $\text{C}^{14}\text{O}-\text{O}-\text{H} + \text{NaHCO}_3 \rightarrow$	(q)	$^{14}\text{CO}_2$
(c)	 $\text{C}-\text{O}-\text{H} + \text{Na} \rightarrow$	(r)	CO_2
(d)	 $\text{S}-\text{O}-\text{H} + \text{NaNH}_2 \rightarrow$	(s)	H_2

32.



Sum of molecular mass of gas (A + C) is :


(a) 88

(b) 90

(c) 92

(d) 40

33.







Sum of molecular mass of gas A + B + C + D is :

52

ORGANIC Chemistry for IIT-JEE


34. Match the column I and II.

Column (I)		Column (II)	
	Molecule		Rotational free energy barrier
(a)		(p)	180 kJ/mol
(b)		(q)	88.3 kJ/mol
(c)		(r)	21 kJ/mol
(d)		(s)	Negative barrier


GENERAL ORGANIC CHEMISTRY

53


35. Consider the following reaction of boron trifluoride (BF_3) and acetone :



- A. What is the critical HOMO (nucleophile) of this reaction ?
- (a) non-bonding orbital on boron (b) σ -orbital of acetone
(c) π -orbital of acetone (d) non-bonding electron pair orbital on oxygen
- B. What is the critical LUMO (electrophile) of the reaction ?
- (a) p -orbital of BF_3 (b) σ -orbital of BF_3
(c) π^* orbital of acetone (d) non-bonding electron pair orbital on oxygen
- C. Which of the following is the correct product of this reaction ?
(Lone electron pairs are not shown explicitly).



36. Rank the following carbocations according to stability (1 = most stable, 5 = least stable).



Put the answer in the boxes.

54

Organic Chemistry


ORGANIC Chemistry for IIT-JEE

37. Among the given molecules, identify aromatic, anti-aromatic and non-aromatic molecules.

(1)	(2)	(3)	(4)
(5)	(6)	(7)	(8)
(9)	(10)	(11)	(12)
(13)	(14)	(15)	(16)
(17)	(18)	(19)	(20)
(21)	(22)	(23)	(24)
(25)	(26)	(27)	(28)
(29)	(30)	(31)	(32)
(33)	(34)	(35)	(36)

GENERAL ORGANIC CHEMISTRY

55



56



ORGANIC Chemistry for IIT-JEE

38. Among the given pairs, which is more reactive towards AgNO_3 (or) toward hydrolysis.




	Compound (A)	Compound (B)	Put the Answer here
1.			
2.			
3.			
4.			
5.			
6.	$\text{CH}_3 - \text{O} - \text{CH}_2 - \text{Cl}$	$\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{Cl}$	
7.			
8.			
9.			

GENERAL ORGANIC CHEMISTRY


57

10.			

39. Put the answer in boxes given as directed.

S.No.	Property	Molecules	Correct Answer	Name of force responsible for the property
A.	highest boiling point	NCl_3 ClNH_2 NH_4Cl NH_3		
B.	highest boiling point	  		
C.	most soluble in water	  		
D.	highest solubility in benzene	 		

40. Circle any conjugated portions of these molecules.




58


ORGANIC Chemistry for IIT-JEE

41. Arrange in the order as directed -


A. The given alkenes in the order of their stability (1- most stable, 3-least stable).




B. Arrange the following in the order of their acidic strength (1-most acidic, 4-least acidic)



C. Arrange the following molecules in order of expected boiling point. (1=highest bpt ; 4=lowest bpt.)



D. Arrange the following alkenes in order of their stability. (1 = most stable ; 5 = least stable).



GENERAL ORGANIC CHEMISTRY**59****42. Match the column. (Matrix)**

Column (I)		Column (II)	
	Compounds		Number of Benzylic hydrogen
(a)		(p)	2
(b)		(q)	3
(c)		(r)	4
(d)		(s)	5


43. Identify (+M) mesomeric & (-M) group of following.

	+M	-M	-I	+I

62


ORGANIC Chemistry for IIT-JEE

50. Consider the H-atoms in the molecule given below and answer the following.




- (A) Identify the type (1° , 2° or 3° alkyl, vinyl, allyl etc.) of these H-atoms.
 (B) Arrange them in the decreasing order of their ease of abstraction (easiest first).

51. Consider the molecule shown below and answer with respect to $H_a \rightarrow H_e$.




- (A) Identify the type of H-atom (1° , 2° , 3° alkyl, vinyl or allyl).
 (B) Arrange them in decreasing order of their bond energy.





52. Rank the following carbocations in order of stability (1 = most stable).



53. Rank the following alkenes according to energy (1 = lowest energy).



GENERAL ORGANIC CHEMISTRY**63****54. Match the column:**


Column (I)		Column (II)	
(Compounds)		(Double bond equivalent value)	
(a)		(p)	11
(b)		(q)	12
(c)		(r)	13
(d)		(s)	14
		(t)	15

64


ORGANIC Chemistry for IIT-JEE

SUBJECTIVE PROBLEMS


1. How many 2° carbon in the following ?




2. Find out the double bond equivalent (DBE) value of the given following compound:



3. Total number of functional groups present in the given following compound :




4. Total number of α -hydrogen in the given following compound is:




GENERAL ORGANIC CHEMISTRY**65**

5. How many carbon atom present in the parent chain in the given following compound?




6. Total number of DBE value in :



7. How many isomers of $C_4H_{10}O$ reacts with Na metal to evolve H_2 gas ? (excluding stereoisomer)






8.



[x]-crown-[y]-ether.

value of $\frac{x+y}{3} = ?$







9. Which of the given following compound will react with $NaHCO_3$ or soluble in $NaHCO_3$?



66

ORGANIC Chemistry for IIT-JEE

10. How many compound are stable after deprotonation ?



ATDB.uno

GENERAL ORGANIC CHEMISTRY

67

ANSWERS — LEVEL 2

1. a - 4; b - 3; c - 2; d - 1
2. a - 3; b - 2; c - 1
3. A - b; B - e; C - a; D - b
4. A - a, c, f, g, k, l; B - b, d, h, j; C - e, i
5. A - d; B - a
6. $C < B < A$
7. a - 2, 1, 4, 3; b - 1, 2, 3, 4; c - 3, 4, 1, 2; d - 3, 2, 4, 1
8. a - iv > ii > i > iii; b - iii > iv > i > ii
9. a - 4, 2, 3, 1; b - 3, 1, 4, 2; c - 2, 1, 3, 4; d - 2, 3, 4, 1
10. A (i) - d > b > c > a; (ii) - c > a > b > d
B (i) - c > d > b > a; (ii) - b > d > c > a
11. (a) 3 2 1 4; (b) 2 1 4 3; (c) 3 1 2 4; (d) 3 4 1 2
12. A - c < a < b; B - a < c < b; C - c < a < b
13. A - c < a < b; B - a < b < c; C - c < a < b; D - d < a < b; E - a < b; F - c < a < b
14. A - $H_c < H_a < H_b$; B - $H_d < H_c < H_b < H_a$
15. A - 1, 3, 4, 6, 7, 8, 9; B - 2, 5; C - 6; D - 3, 4, 6, 7, 9; E - 6, 8, 9; F - 6; G - 7; H - 9; I - 4, 7
16. A - b; B - a; C - d; D - b; E - a; F - d; G - d; H - a
17.

	(i)	(ii)	(iii)	(iv)	(v)
A	6	4	0	10	2
B	0	7	2	9	6
C	5	1	0	5	1
D	1	3	1	3	4
E	2	3	0	2	2
F	10	0	0	10	0
18. a - q; b - p, r; c - p, s; d - q
19. a - p; b - r; c - s; d - q
20. A - c; B - d
21. A - b; B - b; C - a
22. a - p, s; b - q, r; c - q, s; d - p, s
23. a - p, r, s; b - q, r; c - q, r; d - p, r, s

68

ORGANIC Chemistry for IIT-JEE


24. A - b; B - c; C - a; D - c; E - a; F - b; G - d
 25. A - d; B - a; C - d; D - c
 26. A - c; B - b; C - b; D - b; E - b; F - a; G - b; H - c; I - b
 27. i. - sp ; ii. - sp ; iii. - sp^2 ; iv. - sp^2 ; v. - sp^3 ; vi. - sp^3
 28. a - forward b - backward
 29. a - p, q, s; b - p, q, r, s; c - p, q, r, s; d - p, q, r, s
 30. a - s; b - r; c - q; d - p
 31. a - r; b - q; c - s; d - p
 32. b
 33. 65
 34. a - q; b - r; c - s; d - p
 35. A - d; B - a; C - a
 36. A - 3; B - 1; C - 5; D - 2; E - 4
 37. Aromatic— 3, 4, 5, 9, 12, 13, 15, 16, 17, 19, 22, 24, 25, 26, 27, 28, 29, 31, 32, 33, 34, 35, 37, 38, 39, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 53, 54, 56, 57, 58, 61, 62, 63, 66


Non-aromatic— 1, 6, 7, 8, 18, 20, 23, 30, 40, 55, 64, 65

Anti-aromatic— 2, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60



38. 1 - B; 2 - A; 3 - B; 4 - A; 5 - A; 6 - A; 7 - B; 8 - B; 9 - A; 10 - A

39. A.
- NH_4Cl
- , cation-anion interaction

- B.
- 
-
- CH_3
- , van der Waals' forces

- C.
- 
- , H-bonding (Also dipole-dipole)

- D.
- 
- NH, Aromatic stacking

- 40.
- 
- $\text{CH}_2 = \text{C} = \text{CH} - \text{CH} = \text{CH}_2$
- 
- 

41. A. A - 3, B - 2, C - 1,

- B. A - 1, B - 3, C - 2, D - 4

- C. A - 3, B - 1, C - 4, D - 2

- D. A - 1, B - 2, C - 3, D - 4

42. a - s; b - r; c - q; d - p

GENERAL ORGANIC CHEMISTRY

69


43.

+M

-M

-I

+I




✓

x

✓

x




✓

x

✓

x




x

✓

✓

x




x

✓

✓

x




x

✓

✓

x




✓

x

✓

x




x

✓

✓

x



x


✓

✓

x

70

ORGANIC Chemistry for IIT-JEE



✓

×

✓

×

44. A - PA; B - PP; C - NPA; D - PA

45. (i) B, (ii) B, (iii) B, (iv) A

46. (a) HCl ; (b) $\text{CH}_3 - \text{CH}_2 - \text{OH}$; (c) $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{CH}_3$;(d) $\text{CH}_3 - \text{CH}_2 - \text{CH}_3$; (e) $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{CH}_3$ 47. (a) BF_3 ; (b) HO^- ; (c) HO^- ; (d) HI ; (e) HOCl

48. In A, highly electronegative F-atoms are present at the periphery. In liquid term these F-atoms will repel each other due to partial negative charge and thus A will have lower b.pt.

49. $H_d > H_a > H_b > H_c$ 50. A- $H_a = 1^\circ$ alkyl; $H_b = 2^\circ$ alkyl; $H_c = 2^\circ$ allyl; $H_d = \text{vinyl}$; $H_e = 3^\circ$ allylB- Easiest to abstract: $H_e > H_c > H_b > H_a > H_d$ Hardest to abstract51. A- $H_a = 2^\circ$ alkyl; $H_b = 2^\circ$ alkyl; $H_c = \text{vinyl}$; $H_d = 3^\circ$ allyl; $H_e = 1^\circ$ alkylB- $H_c > H_e > H_a > H_b > H_d$

52. A-2; B-1; C-3; D-4

53. A-1; B-4; C-3; D-2

54. a-r; b-t; c-t; d-s

Subjective Solutions:

1. 21

2. 11

3. 3

4. 6

5. 4


6. 13

7. 4

8. 7


9. 2 (c, d)

10. 3 (a, c, f)



2

ISOMERISM (Structural & Stereoisomerism)

replace H_a with D and H_b with H






(X)

replace H_b with D and H_a with H

(Y)

Relation between (X) and (Y) is :


- (a) enantiomers (b) diastereomers
(c) E and Z isomer (d) constitutional isomer
2. Which of the following cyclopentane derivative is optically **inactive** ?




72

ORGANIC Chemistry for IIT-JEE

3. Which is the most stable conformer along the 2, 3 C – C bond axis of the compound ?




4. Assign double bond configurations to the following :




- (a) E (b) Z (c) E, E (d) Z, Z

5. Allegra, a common prescription drug with the structure shown below, is given for the treatment of seasonal allergies. How many stereogenic carbon does Allegra possess ?




- (a) 1 (b) 2 (c) 3 (d) 4
6. How many meso isomers of $C_4H_8Cl_2$ will be ?
- (a) 0 (b) 1 (c) 2 (d) 3
7. The stable form of *trans*-1, 4-dimethylcyclohexane is represented as:



ISOMERISM

73

8. Which of the following compound is non-resolvable (meso) compounds ?




9. $\text{HO}-\text{CH}_2-\text{CH}_2-\text{F}$
(2) (3)

Which conformer of above compound is most stable across C_2-C_3 ?


- (a) staggered (b) eclipsed (partially)
(c) gauche (d) fully eclipsed

10. The following molecule is fluorometholone, a steroidal anti-inflammatory agent. How many stereogenic centers does it contain ?




- (a) 5 (b) 6 (c) 7 (d) 8

11. How many chiral centers are there in these pine ar ar ip not drug ?




- (a) 9 (b) 8 (c) 7 (d) 6

12. How many chiral centers are in the following compound ?










- (a) 4 (b) 5 (c) 6 (d) 7

13. Among the following, the optically inactive compound is:







74







ORGANIC Chemistry for IIT-JEE



15. Which of the following compounds might be useful to the chemist trying to increase the optical purity of the (d) sample ?



16. Which of the following molecules is (are) chiral ?



(a) I and II

(b) III and IV





(c) II, IV and VI

(d) I, II, III and VI



ISOMERISM

73





17. The structure of (S)-2-fluorobutane is best represented by :




18. How are the following compounds related ?



- (a) Diastereomers
(b) Enantiomers
(c) Meso compounds
(d) Identical
19. Which one of the following is chiral ?
(a) 1, 1-Dibromo-1-chloropropane
(b) 1, 3-Dibromo-1-chloropropane
(c) 1, 1-Dibromo-3-chloropropane
(d) 1, 3-Dibromo-2-chloropropane
20. Among the following, the Newmann projections of meso-2, 3-butanediol are :



- (a) P, Q
(b) P, R
(c) R, S
(d) Q, S
21. The binaphthol (**Bnp**) is:







- (a) an optically active compound having chiral centre
(b) an optically inactive compound
(c) a meso compound
(d) an optically active compound without having chiral centre


76

ORGANIC Chemistry for IIT-JEE

22. Which of the following pairs of compounds is a pair of enantiomers ?



23. The maximum number of stereoisomers that could exist for the compound below ?




(a) 6

(b) 8

(c) 10

(d) 16

24. The following pair of compounds is best described as :




(a) identical

(b) diastereomers



(c) enantiomers

(d) none of the above



25. Determine the absolute configurations of the labeled carbons (a and b):

(a) $a = R$; $b = R$ (b) $a = R$; $b = S$ (c) $a = S$; $b = R$ (d) $a = S$; $b = S$


26. Which of the structures (a - d) will be produced if a "ring flip" occurs in the following compound in chair form ?



27. Which of the following compounds is most stable ?




28. Which is the most stable chair form of this compound ?




ISOMERISM

77




29. Which pairs of the salts would have identical solubilities in methanol ?




- (a) I & IV (b) I & III (c) I & II (d) II & IV

30. The following compounds differ in respect of :



- (a) their chemical and physical properties
(b) nothing
(c) the direction in which they rotate plane of polarized light
(d) their interactions with molecules

31. Indicate whether each of the following pairs are identical, or ?



- I (a) enantiomers (b) identical (c) enantiomers (d) enantiomers
- II (a) diastereomers (b) enantiomers (c) diastereomers (d) identical
- III (a) enantiomers (b) enantiomers (c) identical (d) identical

32. Which of the following is achiral ?






78

ORGANIC Chemistry for IIT-JEE



(d) a molecule of 3-methylheptane

33. Which of the following compounds are meso forms ?



(a) 1 only

(b) 3 only

(c) 1 and 2

(d) 2 and 3

34. The separation of a racemic mixture into pure enantiomers is termed as :





(a) Racemization

(b) Isomerization

(c) Resolution

(d) Equilibration

35. Rank of the following groups in order of R, S precedence (IV is highest) :



I II

II I

I II

III IV

(a) 3

2

4

2

(b) 1

4

2

3

(c) 3

4

1

2






(d) 3

4





2

1

36. Which of the following is a meso compound ?



37. Among the following structures, select E isomers (arrows indicate the bonds to be considered) ?



(a) 1 and 2

(b) 1 and 3


(c) 1 and 4

(d) 2 and 3

ISOMERISM

79


38. Which of the following compounds has a zero dipole moment ?



39. On Pluto, where everything is frozen, astronauts discovered two forms of butane gauche and anti. Assuming that there are no rotations around single bonds, which statement about the two forms is correct ?

- (a) They are enantiomers
 (b) They are diastereoisomers
 (c) They are meso compounds
 (d) The gauche form has two stereogenic centers, and the anti has only one

40. Which of the following will show optical activity ?




- (a) A, D and E (b) A and E only (c) B, C and D (d) All except C

41. Among the structures shown below, which has lowest potential energy ?




42. Which of the following molecules is/are chiral ?




- (a) I (b) II (c) III (d) I, II

43. A compound was synthesized by a student, but its structure was not identified. However, his wonderfully helpful instructor told him that it was a meso compound with 5 carbons and 2 stereogenic centers. Which of the following structures should the student consider as possibilities for his compound ?




- (a) I, II, IV (b) II, IV (c) I, III, V (d) II, IV, V

44. How many isomers are possible for the following molecule ?




- (a) 1 (b) 2 (c) 3 (d) 4

45. Which of the following molecules are chiral ?




- (a) I, II, III and IV (b) II, III and IV (c) II and IV (d) I and II


46. Which equilibrium is not rapid at room temperature ?




47. Which is the lowest energy conformation of butane ?



48. Which of the structures given below are chiral ?







Fischer projections




- (a) I, II, III (b) II, III, V (c) II, III (d) I, II

ISOMERISM**81**

49. Which of the following carboxylic acids could be resolved by reaction with an enantiomerically pure chiral amine ?



50. What is the relationship between the molecules in the following pairs ?







- (a) enantiomers (b) diastereomers (c) identical (d) structural isomers

51. What are the correct designations for the structure below ?







- (a) E, E (b) Z, E
(c) E, Z (d) No geometrical isomers are possible

52. Which of the following molecules are chiral ?



- (a) I and III (b) I and V (c) II and III (d) II, III, IV


53. Which one of the following isomeric structures has the lowest energy ?



82





ORGANIC Chemistry for IIT-JEE

54. The following compounds are identical with respect to :








- (a) molecular composition (b) boiling point
(c) melting point (d) IUPAC name

55. Among the following, the most stable isomer is :



56. The most stable conformation of the following compound is :



57. Which of the following molecules have non-zero dipole moments ?


- (I) gauche conformation of 1, 2-dibromoethane
(II) anti conformation of 1, 2-dibromoethane
(III) *trans*-1, 4-dibromocyclohexane
(IV) *cis*-1, 4-dibromocyclohexane
(V) tetrabromomethane
(VI) 1, 1-dibromocyclohexane

- (a) I and II (b) I and IV
(c) II and V (d) I, IV and VI

ISOMERISM

83

58. What is the maximum number of stereoisomers possible for discodermolide ?



- (a) 2^{14} (b) 2^{15} (c) 2^{16} (d) 2^{17}

59. An aqueous solution containing compounds A and B shows optical activity. A and B are stereoisomers. Which of the following possibilities cannot be correct ?

- (a) A has two chiral centers, but B does not have any because it has a symmetry plane
 (b) A and B are enantiomers
 (c) A and B are diastereomers
 (d) A and B are not present in equal amounts


60. Which of the following structures represents the lowest-energy form of (1S, 2S, 4R)-trimethyl-cyclohexane ?



61. Which one of the following is a diastereomer of (R)-4-bromo-cis-2-hexene ?

- (a) (S)-4-bromo-cis-2-hexene
 (b) (S)-5-bromo-trans-2-hexene
 (c) (R)-4-bromo-trans-2-hexene
 (d) (R)-5-bromo-trans-2-hexene

62. The structural formula of cocaine is shown below. How many stereogenic carbon atoms are there in this molecule ?




- (a) 1 (b) 2 (c) 3 (d) 4




84

ORGANIC Chemistry for IIT-JEE

63. Which of the following statements best describes the stereochemical relationships of compound I, II and III shown below ?




- (a) All compounds are chiral
 (b) None of the compounds is chiral
 (c) I and II are meso compounds
 (d) I and II are diastereomers, and III is a meso compound
 (e) I and II are chiral
64. What is the absolute configuration of the following molecules ? (NS = the molecule has no center) Note : For the purpose of this question only, the order of stereocenters is not specified ; i.e., R, S = S, R.



	I	II	III	IV
(a)	R	R, S	R	NS
(c)	R	R, S	NS	NS

	I	II	III	IV
(b)	R	R, R	S	R, R
(d)	R	R, S	R	R, S

65. The number of all the possible stereoisomers formed by the given compound is :




- (a) 2
 (c) 22


- (b) 3
 (d) 64

ISOMERISM**85**


66. The relationship among the following pairs of isomers is:




(I)



(II)



(III)



(IV)

I	A: Constitutional
II	B: Configurational
III	C: Conformational
IV	D: Optical


(a) I – A, II – B, III – C, IV – D

(b) I – A, II – A, III – C, IV – D

(c) I – B, II – A, III – B, IV – I

(d) I – C, II – B, III – C, IV – B

67. The structural formula of sativene is shown below. How many stereogenic centers are there in this molecule ?



(a) 2

(b) 3





(c) 4

(d) 5





86

ORGANIC Chemistry for IIT-JEE




68. Which of the following is the least stable conformer of cyclohexane ?



69. The S- enantiomer of ibuprofen is responsible for its pain-relieving properties. Which one of the following structures shown below is (S)-ibuprofen ?



70. Which of the following depict the same ?




- (a) 1 and 2
(c) 2 and 3

- (b) 1 and 3
(d) 1, 2, and 3

ISOMERISM


87

71. A naturally occurring substance has the constitution shown below. How many may have this constitution ?



- (a) 2 (b) 8 (c) 16 (d) 64

72. The absolute configurations of the two centers in the following molecule are :




- (a) 2(R), 3(S) (b) 2(R), 3(R) (c) 2(S), 3(S) (d) 2(S), 3(R)

73. The total number of stereoisomer possible for 2, 3-dichloro butane :

- (a) 2 (b) 3
(c) 4 (d) 5

74. Which of the following structure is not meso-2,3-butanediol ?




75. A solution of optically active 1-phenylethanol racemizes in acidified aqueous medium. It is due to :

- (a) enolization (b) carbonium ion formation
(c) carbanion formation (d) reversible oxidation-reduction

76. The most stable conformation of ethylene glycol is :

- (a) Anti (b) Gauche (c) Partially eclipsed (d) Fully eclipsed



The molecules represented by the above two structures are :

- (a) identical (b) enantiomers
(c) diastereomers (d) epimers


78. The correct order of priority of groups -SCH₃ (I), -NO₂ (II), -C≡CH (III) and -CH₂C₆H₅ (IV), on the basis of CIP classification, is (increasing order) :

- (a) I, III, II, IV (b) IV, III, II, I
(c) II, IV, I, III (d) III, IV, II, I

88

ORGANIC Chemistry for IIT-JEE

79. The configuration at C-2 and C-3 of the compound given :







(a) 2R, 3S

(b) 2S, 3R




(c) 2S, 3S

(d) 2R, 3R

80. Amongst the following amino acids, the (R) - enantiomer is represented by :



81. Which of the following is a meso compound ?



(d) All of these

82. Predict stereochemistry of product when *d* and *l*-amine reacts with *l*-acid:


(a) Diastereomers

(b) Meso

(c) Racemic

(d) Pure Enantiomer

83. How many chiral center (excluding N centres) are there in morphine?



(a) 4

(b) 5

(c) 6

(d) More than 6


84. Which dimethylcyclobutane is optically active ?

(a) *trans*-1, 2(b) *cis*-1, 2(c) *trans*-1, 3(d) *cis*-1, 3


ISOMERISM

85

- 85.** Which of the following is the enantiomer of the compound shown below?



- 86.** How many different stereoisomers are possible for the following compound?




- (a) 1 (b) 2 (c) 3 (d) 4

- 87.** The following compounds are best described as :



- (a) enantiomers
(b) diastereomers
(c) not stereoisomers
(d) conformational isomers (differing by single bond rotation)

- 88.** Rank the following substituent groups in order of decreasing priority according to the Cahn-Ingold-Prelog system :



- (a) $2 > 3 > 1$ (b) $1^1 > 3 > 2^2$ (c) $3 > 1^3 > 2^2$ (d) $2 > 1 > 3$


- 89.** Compare the stabilities of the following two compounds :

A : *cis*-1-Ethyl-3-methylcyclohexane

B : *trans*-1-Ethyl-3-methylcyclohexane

- (a) A is more stable
(b) A and B are of equal stability
(c) B is more stable
(d) No comparison can be made

90. What, if anything, can be said about the magnitude of the equilibrium constant K for the following equilibrium?




- (a) $K = 1$
 (c) $K > 1$
- (b) $K < 1$
 (d) No estimate of K can be made

90

ORGANIC Chemistry for IIT-JEE

91. What is the relationship between the two structures shown ?



- (a) Constitutional isomers
 (b) Stereoisomers
 (c) Different drawing of the same conformation of the same compound
 (d) Different conformation of the same compound

92. Which of the following statements is true ?

- (a) van der Waals' strain in *cis*-1, 2-dimethylcyclopropane is the principal reason for its decreased stability relative to the *trans* isomer
 (b) Cyclohexane gives off more heat per CH_2 group on being burned in air than any other cycloalkane
 (c) The principal source of strain in the boat conformation of cyclohexane is angle strain
 (d) The principal source of strain in the *gauche* conformation of butane is torsional strain


93. $\text{Ph}-\text{CH}=\text{NO}_2\text{H} \xrightarrow[3 \text{ days}]{\text{isomerises}}$ (x), Isomer (x) is :

- (a) $\text{Ph}-\text{NO}-\text{CH}_2\text{OH}$ (b) $\text{Ph}-\text{CH}_2-\text{NO}_2$
 (c) $\text{Ph}-\text{NH}-\text{CO}_2\text{H}$ (d) None

94. Which of the following will not show geometrical isomerism ?


- (a) $\text{CH}_3-\text{C}(\text{CH}_3)=\text{CH}-\text{CH}_3$ (b) $\text{Cl}_3\text{C}-\text{CH}=\text{CH}-\text{CH}_2-\text{CH}_3$
 (c) $\text{CH}_3-\text{CH}=\text{CH}-\text{CH}_3$ (d) $\text{CH}_3-\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-\text{CH}_3$

95. The two compounds shown below are :



- (a) diastereomers (b) enantiomers (c) epimers (d) regiomers

96. The molecular formula of diphenylmethane,




, is $\text{C}_{13}\text{H}_{12}$;

How many structural isomers are possible when one of the hydrogen is replaced by a chlorine atom ?

- (a) 6 (b) 4 (c) 8 (d) 7

97. Correct configuration of the following molecule is :



- (a) 2S, 3S (b) 2S, 3R (c) 2R, 3S (d) 2R, 3R


ISOMERISM

91

98. Maximum enol content is in : (a) CH3CHO (b) CH3COCH3 (c) CH3CH=CH2 (d) CH3C(=O)CH2CH3
99. Which of the following will have one of the stereoisomer meso ? (a) 2-chlorobutane (b) 2,3-dichlorobutane (c) 2,3-dichloropentane (d) 2-hydroxypropanoic acid
100. The correct decreasing order in the enol content of following molecules is : (a) CH3CHO (b) CH3COCH3 (c) CH3CH=CH2 (d) CH3C(=O)CH2CH3
101. Total number of stereoisomers of the compound 1-bromo-3-chlorocyclobutane is : (a) 0 (b) 1 (c) 2 (d) 3
102. Total number of stereoisomers of the 1,3-dichlorocyclohexane is : (a) 0 (b) 1 (c) 3 (d) 4
103. Total number of stereoisomers of the compound 1,4-dichlorocyclohexane is : (a) 0 (b) 1 (c) 2 (d) 4
104. Total number of stereoisomers of the compound 2,4-dichlorohexane is : (a) 0 (b) 1 (c) 3 (d) 4
105. In which of the following keto form is more dominating than enol form : (a) CH3CHO (b) CH3COCH3 (c) CH3CH=CH2 (d) CH3C(=O)CH2CH3
106. Among the following compounds, which will give maximum enol content in solution : (a) CH3CHO (b) CH3COCH3 (c) CH3CH=CH2 (d) CH3C(=O)CH2CH3
107. Which of the following has unstable enol form ? (a) CH3CHO (b) CH3COCH3 (c) CH3CH=CH2 (d) CH3C(=O)CH2CH3
108. Calculate enantiomeric excess of mixture containing 6g of (+) 2-butanol and 4g of (-) 2-butanol.

94


ORGANIC Chemistry for IIT-JEE



How many representations of lactic acid are possible in Fischer projection (d & l) ?


- (a) 8 (b) 12
(c) 24 (d) 36

121. Total number of stereoisomer formed by the given compound is :






- (a) 2 (b) 3
(c) 4 (d) 8

122. The number of stereoisomers formed by the given compound is :




- (a) 2 (b) 3
(c) 4 (d) 5

123. Which of the following compound does not undergo base - catalyzed exchange in D_2O even though it has an α -hydrogen?






(d) both (b) & (c)



Identify the product formed in the above reaction:


ISOMERISM

95



(d) None of these




- 125.** In 3-methyl-2-cyclohexenone which hydrogen cannot undergo deuterium exchange when it reacts with $\text{CH}_3\text{O}^-/\text{CH}_3\text{OD}$?



- (a) H_1, H_4
(c) H_3, H_2

- (b) H_4
(d) H_5, H_3

126.



(I)


(II)

(III)

The tautomer of II is :

- (a) I
(c) both I and III
- (b) III
(d) none of these

127.






In the enolization of the given molecule, the H-atom involved is :

- (a) α -H (b) β -H
(c) γ -H (d) cannot be enolized

96

ORGANIC Chemistry for IIT-JEE

128.



Among the given structure which can exhibit tautomerism ?




(a) I only

(b) II only

(c) III only

(d) none of these

129.



Identify the which can exhibit tautomerism ?

(a) I only

(b) II only

(c) III only

(d) all of these



(I)

(II)

Between the two tautomers which is more stable ?




(a) I

(b) II

(c) I = II

(d) none of these

131.



Correct stability order of the given tautomers is :




(a) I > II > III

(b) III > II > I

(c) II > I > III

(d) II > III > I

132.



Correct stability order of the given tautomers is :




(a) I > II > III

(b) III > II > I

(c) II > I > III

(d) II > III > I

133.



The correct order of enol contents x, y, z is :

ISOMERISM


97

(a) $x > y > z$

(b) $z > y > x$

(c) $y > x > z$

(d) $x > z > y$


The correct order of x, y, z is :

(a) $x > y > z$

(b) $z > y > x$

(c) $y > x > z$

(d) $x > z > y$




Among the given ketones, the one which does not enolize is :

(a) I



(b) II

(c) III

(d) none of these



The product of this reaction should be :




Among the given compounds, the correct order of enol content is :

(a) $I > II > III$

(b) $III > II > I$

(c) $II > I > III$

(d) $II > III > I$



Among the given compounds, the correct order of enol content is :

(a) $I > II > III$

(b) $III > II > I$


(c) $II > I > III$

(d) $II > III > I$


98

ORGANIC Chemistry for IIT-JEE


139.



(I)



(II)




(III)

Among the given compounds, the correct order of enol content is :

- (a) I > II > III (b) III > II > I (c) III > I > II (d) II > I > III





140.




How many geometrical isomers are possible for the above compound ?

- (a) 3 (b) 4 (c) 6 (d) 8

141. Which of the following compound will not show geometrical isomerism across the π -bond ?




142.



Choose the correct relation between l_1 and l_2 ?

- (a) $l_1 = l_2$ (b) $l_1 > l_2$ (c) $l_1 < l_2$ (d) $l_2 = 2l_1$


143.



Choose the correct relation between l_1 and l_2 ?

- (a) $l_1 = l_2$ (b) $l_1 > l_2$ (c) $l_1 < l_2$ (d) $l_2 = 2l_1$

144.




How many geometrical isomers are possible for the above compound ?

- (a) 0 (b) 2 (c) 3 (d) 4

ISOMERISM

99


145.



How many geometrical isomers are possible for the above compound ?

- (a) 0 (b) 2 (c) 3 (d) 4


146.



How many geometrical isomers are possible for the above compound ?

- (a) 0 (b) 2 (c) 3 (d) 4


147.



How many geometrical isomers are possible for the above compound ?

- (a) 0 (b) 2 (c) 3 (d) 4


148.



How many geometrical isomers are possible for the above compound ?

- (a) 0 (b) 2 (c) 3 (d) 4


149.



How many geometrical isomers are possible for the above compound ?

- (a) 0 (b) 2 (c) 3 (d) 4


150.



I and II are geometrical isomers of each other because

- (a) $l_1 = l_2$ (b) $l_1 > l_2$
(c) $l_2 > l_1$ (d) l_1 and l_2 cannot be compared.

151.




How many geometrical isomers are possible for this compound ?

- (a) 2 (b) 3 (c) 4 (d) 8


100

ORGANIC Chemistry for IIT-JEE




How many geometrical isomers are possible for this compound?

- (a) 2 (b) 3
(c) 4 (d) 6




How many geometrical isomers of this compound are possible ?

- (a) 2 (b) 3
(c) 4 (d) 6




- (a) chiral (b) C_2 axis of symmetry
(c) Optically active (d) Achiral



Relationship between above pair (A) & (B) is :

- (a) Enantiomer (b) Diastereomers (c) Identical (d) Structural isomer







From the above compound (A), (B), (C) & (D) chiral compound is :

- (a) A (b) B (c) C (d) D



ISOMERISM

101

157. Which of the following compound is achiral ?






158.



R and S configuration of compound (A) & (B) will be :

- (a) R, R (b) R, S
(c) S, R (d) S, S

159. Which of following compound has center of symmetry?






(d) All of these

102


ORGANIC Chemistry for IIT-JEE

160. Which mixture of structure in each beaker would rotate plane polarized light ?






(d) All of these

161. Which of following compound will rotate the plane polarized light at room temperature?



162. Which of the following having plane of symmetry ?





(d) All of these

ISOMERISM





103

163. Which of following compound is achiral ?







(d) All of these

164. Which of the following compound has plane of symmetry ?






(d) None of these

165. Which of following is isomer







166. Among the given pairs, in which pair second compound has less enol content than first compound?







(d) none of these

167. Which of the following is incorrect relation between given pairs ?



168. $\text{Ph}-\underset{\text{OH}}{\underset{\text{(A)}}{\text{CH}}}-\overset{\text{O}}{\parallel}{\text{C}}-\text{H} \xrightarrow[\text{H}_2\text{O}]{\text{HO}^\ominus} (\text{B})$; (A) and (B) are isomer and isomerization effectively

carried out by trace of base (B). Identify (B).



169. $\text{CH}_3-\text{CH}=\text{CH}-\text{H}=\text{CH}-\text{C}\text{I}_3$ total number of geometrical isomer is :






(a) 2

(b) 3

(c) 4

(d) 6


170. Identify most stable enol form of teric acid:




ISOMERISM

105


171. Which structure is most stable ?




172. Identify conformer of 2-methyl pentane :



173. The lowest energy conformer of  is:



174.



How many atoms will be bisect during plane of symmetry ?

- (a) 2 (b) 4
(c) 6 (d) 8




106

ORGANIC Chemistry for IIT-JEE

175. The number of all types of isomers of chlorobutane is :

- (a) 2 (b) 4
(c) 6 (d) 5

176. Which of the following pairs of compounds are not positional isomers ?



(d) All of these


177. Which of the following pairs of compounds are functional isomers ?



178. The isomeric alcohol which has a chiral carbon atom is:

- (a) *n*-butyl alcohol (b) *iso*-butyl alcohol (c) *sec*-butyl alcohol (d) *tert*-butyl alcohol

179. The pair of enantiomers among the following compound is:




- (a) I and IV (b) II and IV (c) II and III (d) I and II

180. Which of the following is chiral?

- (a) Cell phone (b) Spiral staircase (c) Scissor (d) All of these





181. In which of the following compound, possess plane of symmetry as well as centre of symmetry?








ISOMERISM

107


182. Which of the following compound has one of the stereoisomers as a meso compound?



183. For the following Newman projection







184. Which of the following is correct for the given compound?



- (a) It possess centre of symmetry
- (b) It possess C_4 axis of symmetry
- (c) It possess plane of symmetry
- (d) Compound is chiral

185. Which of the following molecules has axis of symmetry and a coaxial plane of symmetry?



(e) All of these

108



ORGANIC Chemistry for IIT-JEE

ANSWERS - LEVEL 1


1.	(b)	2.	(c)	3.	(b)	4.	(c)	5.	(a)	6.	(b)	7.	(c)	8.	(d)
9.	(c)	10.	(d)	11.	(b)	12.	(c)	13.	(a)	14.	(b)	15.	(b)	16.	(d)
17.	(c)	18.	(a)	19.	(b)	20.	(b)	21.	(d)	22.	(b)	23.	(c)	24.	(d)
25.	(a)	26.	(b)	27.	(d)	28.	(b)	29.	(a)	30.	(c)	31.	(c)	32.	(a)
33.	(b)	34.	(c)	35.	(c)	36.	(d)	37.	(c)	38.	(d)	39.	(b)	40.	(a)
41.	(a)	42.	(d)	43.	(a)	44.	(d)	45.	(a)	46.	(b)	47.	(c)	48.	(b)
49.	(c)	50.	(c)	51.	(d)	52.	(d)	53.	(e)	54.	(a)	55.	(d)	56.	(c)
57.	(d)	58.	(b)	59.	(a)	60.	(a)	61.	(c)	62.	(d)	63.	(e)	64.	(d)
65.	(b)	66.	(b)	67.	(d)	68.	(d)	69.	(d)	70.	(d)	71.	(d)	72.	(a)
73.	(b)	74.	(a)	75.	(b)	76.	(b)	77.	(a)	78.	(b)	79.	(c)	80.	(b)
81.	(d)	82.	(a)	83.	(b)	84.	(a)	85.	(a)	86.	(d)	87.	(a)	88.	(d)
89.	(a)	90.	(b)	91.	(b)	92.	(a)	93.	(b)	94.	(a)	95.	(b)	96.	(b)
97.	(a)	98.	(d)	99.	(c)	100.	(c)	101.	(c)	102.	(c)	103.	(c)	104.	(d)
105.	(d)	106.	(a)	107.	(c)	108.	(b)	109.	(d)	110.	(a)	111.	(c)	112.	(c)
113.	(a)	114.	(d)	115.	(b)	116.	(b)	117.	(c)	118.	(d)	119.	(b)	120.	(c)
121.	(a)	122.	(b)	123.	(d)	124.	(b)	125.	(b)	126.	(c)	127.	(c)	128.	(b)
129.	(d)	130.	(a)	131.	(c)	132.	(d)	133.	(d)	134.	(d)	135.	(b)	136.	(b)
137.	(c)	138.	(a)	139.	(d)	140.	(b)	141.	(b)	142.	(a)	143.	(c)	144.	(b)
145.	(b)	146.	(b)	147.	(b)	148.	(a)	149.	(b)	150.	(c)	151.	(a)	152.	(c)
153.	(b)	154.	(d)	155.	(c)	156.	(a)	157.	(c)	158.	(d)	159.	(d)	160.	(d)
161.	(b)	162.	(d)	163.	(d)	164.	(d)	165.	(d)	166.	(c)	167.	(d)	168.	(c)
169.	(b)	170.	(c)	171.	(c)	172.	(d)	173.	(b)	174.	(c)	175.	(d)	176.	(c)
177.	(b)	178.	(c)	179.	(c)	180.	(d)	181.	(d)	182.	(b)	183.	(b)	184.	(c)
185.	(e)														

ISOMERISM


109



1. Match the Column (I) and (II).

Column (I)		Column (II)	
Reaction		Stereoisomers	
(a)	$\text{CH}_3 - \text{CH} = \text{CH} - \text{CH} = \text{N} - \text{OH}$	(p)	2
(b)		(q)	4
(c)	$\text{CH}_3 - \text{CH} = \text{CH} - \text{CH} = \text{CH} - \text{CH} = \text{CH} - \text{CH}_3$	(r)	6
(d)	$\text{CH}_3 - \text{CH} = \text{CH} - \text{CH} = \text{CH} - \text{CH} = \text{CH} - \text{Ph}$	(s)	8

2. Match the Column (I) and (II)



Column (I)		Column (II)	
Group		Equilibrium Constant	
(a)	$\text{R} = -\text{H}$	(p)	38
(b)	$\text{R} = -\text{CH}_3$	(q)	23
(c)	$\text{R} = -\text{Et}$	(r)	18
(d)	$\text{R} = -\text{CH}(\text{CH}_3)_2$	(s)	1

110

ORGANIC CHEMISTRY

ORGANIC Chemistry for IIT-JEE

3. Match the Column (I) and (II). (Matrix)

Column (I)		Column (II)	
Molecule		Nature	
(a)		(p)	Chiral
(b)		(q)	Achiral
(c)		(r)	Meso
(d)		(s)	Compound containing even number of chiral centers





ISOMERISM**111****4. Match the Column (I) and (II). (Matrix)**

Column (I)		Column (II)	
Compound		Isomerism	
(a)		(p)	Geometrical isomerism
(b)		(q)	Optical isomerism
(c)		(r)	Compound containing plane of symmetry
(d)		(s)	Compound containing center of symmetry





112

ORGANIC Chemistry for IIT-JEE

5. Match the Column (I) and (II).

Column (I)		Column (II)	
Molecules		Relationship	
(a)		(p)	Identical
(b)		(q)	Enantiomer
(c)		(r)	Diastereomer
(d)		(s)	Structural Isomerism

6. Match the Column (I) and (II). (1 mark)

Column (I)		Column (II)	
Compound		Nature	
(a)		(p)	cis-compound
(b)		(q)	trans-compound
(c)		(r)	Optically active
(d)		(s)	Optically inactive





ISOMERISM**113****7. Match the Column (I) and (II). (Matrix)**

Column (I)		Column (II)	
Molecule		Property	
(a)		(p)	Chiral centers containing compound
(b)		(q)	Presence of stereocenter
(c)		(r)	Optically active compound
(d)		(s)	Compound containing plane of symmetry





8. Match the Column (I) and (II). (Matrix)

Column (I)		Column (II)	
Molecule		Property	
(a)		(p)	Polar molecule
(b)		(q)	Optically active
(c)		(r)	Optically inactive
(d)		(s)	Plane of symmetry

9. Match the Column (I) and (II). (Matrix)

Column (I)		Column (II)	
Molecule		Property	
(a)		(p)	Meso compound
(b)		(q)	Achiral
(c)		(r)	Chiral compound
(d)		(s)	Compound will show geometrical isomerism

ISOMERISM**115****10. Match the Column (I) and (II).**

Column (I)		Column (II)	
Modified Newmann Projection		Conformers	
(a)		(p)	Fully eclipsed
(b)		(q)	Partially eclipsed
(c)		(r)	Gauche
(d)		(s)	Staggered




116

ORGANIC Chemistry for IIT-JEE

11. Match the Column (I) and (II).

Column (I)		Column (II)	
Newmann Projection		Name of the Compound	
(a)		(p)	3-methyl pentane
(b)		(q)	n-butane
(c)		(r)	Methyl-cyclopentane
(d)		(s)	1,2,4-trimethyl cyclohexane

ISOMERISM**117****12. Match the Column (I) and (II). (Matrix)**

Column (I)		Column (II)	
Molecule		Property	
(a)		(p)	Rotates plane polarized light
(b)		(q)	Cannot rotate plane polarized light
(c)		(r)	Plane of symmetry
(d)		(s)	Centre of symmetry

13. Match the Column (I) and (II).

Column (I)		Column (II)	
Molecule		Stereocenters	
(a)	$\text{CH}_3 - \text{CH} = \text{CH} - \underset{\text{Br}}{\text{CH}} - \text{CH}_3$	(p)	1
(b)	$\text{H} - \text{C} \equiv \text{C} - \text{CH} = \text{CH} - \underset{\text{Br}}{\text{CH}} - \underset{\text{Br}}{\text{CH}} - \text{CH}_3$	(q)	2
(c)	$\text{Ph} - \overset{\text{O}}{\parallel}{\text{S}} - \text{CH} = \text{CH} - \underset{\text{CH}_3}{\text{CH}} - \text{CH}_3$	(r)	3
(d)	$\text{Ph} - \underset{\text{Cl}}{\text{CH}} - \text{Et}$	(s)	4





118

ORGANIC Chemistry for IIT-JEE

14. Match the Column (I) and (II).

Column (I)		Column (II)	
Molecule		Stereoisomers	
(a)		(p)	2
(b)		(q)	0
(c)		(r)	4
(d)		(s)	8

ISOMERISM**119****15. Match the Column (I) and (II). (Matrix)**

Column (I)		Column (II)	
Molecule		Property	
(a)		(p)	Meso Compound
(b)		(q)	Compound having even no. of chiral centres
(c)		(r)	Optically active compound
(d)		(s)	Compound having odd no. of chiral centres.

120

ORGANIC Chemistry for IIT-JEE

16. Match the Column (I), (II) and (III). (Matrix)

Column (I)		Column (II)		Column (III)	
Property		Molecule		No. of Chiral Center	
(a)		(p)	Optically active	(w)	0
(b)		(q)	Optically inactive	(x)	1
(c)		(r)	Plane of symmetry	(y)	2
(d)		(s)	Centre of symmetry	(z)	3

17.

(a)		(b)		(c)	
(d)		(e)		(f)	
(g)		(h)			


From the above compounds select :

(A)	two of which are chiral and contain chiral centre :	
(B)	two of which are achiral and contains chiral centre :	
(C)	two of which are chiral and does not contain chiral centre :	
(D)	two of which are achiral and does not contain chiral centre :	

ISOMERISM

121

18. Comprehension



Consider the given structures and answer A, B & C.


A. Which of the compound is optically active ?

- (a) P (b) Q (c) S (d) T

B. Which of the isomer is most stable ?

- (a) R (b) S (c) T (d) U

19. Identify relationship between following pairs :



















If they are enantiomer answer will be 1, if they are diastereomers answer will be 2, if they are constitutional isomers answer will be 3 and if they are identical present 4 as the answer. Sum of answer of each part a + b + c + d is :

122

ORGANIC Chemistry for IIT-JEE

20. In each of the following three questions a hydrocarbon is named. For each select from among the sixteen conformational structures (a through p) all structures that represent possible conformers of that compound. Write letters (a through p), corresponding to your selections, in each answer box.

A.	2-methylbutane	
B.	2,3-dimethylpentane	
C.	1-ethyl-1, 3-dimethyl cyclohexane	

(a)		(b)		(c)	
(d)		(e)		(f)	
(g)		(h)		(i)	
(j)		(k)		(l)	
(m)		(n)		(o)	
(p)					

ISOMERISM

123

21. Examine structures a through j, shown below, with respect to their symmetry or lack of it. Assume that the five-membered rings and the ring in compound g are planar. The wedge-hatched bonds in b, c, d & e designate specific configurations. Also, for the acyclic compounds assume stable anti conformations. Answer each of the following questions by writing letters (a through j), corresponding to your selections, in each answer box. If there is no structure that fits the description enter an x in the answer box.

A.	Which structures are chiral ?	
B.	Which structures have a plane of symmetry ?	
C.	Which structures have a center of symmetry ?	

(a)		(b)		(c)	
(d)		(e)		(f)	
(g)		(h)	$C_2H_5CHCl_2$	(i)	$C_2H_5CHClC_2H_5$
(j)	$C_2H_5CHClCH_3$				

22. (i) 1,2-dichlorocyclopropane = w
(ii) 1,3-dimethyl-cyclobutane = x
(iii) 2-bromo-3-chlorobutane = y
(iv) 1,3-dimethyl cyclohexane = z









Calculate total number of stereoisomer of the above compounds.

Sum of $w + x + y + z = \dots\dots$

126

ORGANIC Chemistry for IIT-JEE

27. The configuration of eight compounds, a through h are shown below, using various kind of stereo representations. To answer the question given below, write (a through h) indicating your choice.

(a)		(b)		(c)	
(d)		(e)		(f)	
(g)		(h)			

A.	Which of these configuration are achiral?	
B.	Which configuration has no stereogenic center?	
C.	Which configuration has more than one stereogenic center?	
D.	Which of these configuration are meso compound?	

28. The structural formula of ten compounds, (I) through (X) are drawn below, you may select any one of these structure.


Answer the following question about that compound.

ISOMERISM

127

I		II		III	
IV		V		VI	
VII		VIII		IX	
X					

- A. How many chiral centre are present in this compound ?
 (a) 0 (b) 1 (c) 2 (d) 3
 (e) 4 (f) 5
- B. Is this compound chiral or achiral ?
 (a) Chiral (b) Achiral
- C. What symmetry element are present in this compound ?
 (a) None (b) Plane of symmetry (c) Center of symmetry
29. The structure of one of the enantiomers of the amino acid cysteine is shown below.



Classify this structure as :

(a) R or S


(b) D or L

128


ORGANIC Chemistry for IIT-JEE

30. Identify the following double bonds either E, Z or None (N) in the compounds given below either.

A.



- B. (a) Bongkreic acid is a natural compound produced by *Pseudomonas cocovenenans*, and isolated from a mold that grows on banana, a ferment used in a coconut dish. (a) Label each double bond as E, Z or neither (N).




- (b) How many total stereoisomers (including all types) are possible for bongkreic acid ?

.....


- (c) How many sites of unsaturation are present in bongkreic acid ?

ISOMERISM**129**

31. Designate the following double bonds as E, Z or none (N) configuration in the boxes provided below.



32. The following compounds may exist as two or more stereoisomers. These may be classified as enantiomer pairs or meso compounds.




Answer the following question about the above structure.

- (A) Total number of stereoisomers :
 (B) Number of enantiomeric pairs :
 (C) Number of meso compounds :

130




ORGANIC Chemistry for IIT-JEE

33. Identify axis of symmetry in the given compound.









ISOMERISM

131



34. For each of the following pharmaceutical compounds, identify all stereogenic (i.e., all asymmetric carbon atoms) and label the configuration of each as being either (R) or (S).



132

ORGANIC Chemistry for IIT-JEE

35. Find relationship between given pair :

		Identical	Enantiomer	Diastereomer	Constitutional Isomer
1.					
2.					
3.					
4.					
5.					
6.					
7.					

ISOMERISM					
8.					
9.					
10.					

36. Comprehension

Structural formula of compound (A) is following:





- A. The correct statement(s) about the compound (A) is/are:
- The total number of stereoisomers possible for (A) is 3
 - The total number of mesoisomer possible for (A) is 1
 - The total number of pair of enantiomer possible for (A) is 1
 - All of these
- B. Number of plane of symmetry in cis-form of compound (A) is:
- 0
 - 1
 - 2
 - 3
37. Match the column. (Matrix)

Column (I)		Column (II)	
No. of Carbon		No. of structural isomer	
(a)	C_4H_{10}	(p)	2
(b)	C_5H_{12}	(q)	3
(c)	C_6H_{14}	(r)	5
(d)	C_7H_{16}	(s)	9


134

ORGANIC Chemistry for IIT-JEE








38. Match the column. (Matrix)





Column (I)		Column (II)	
Compound		% of enol content	
(a)		(p)	100 %
(b)		(q)	76 %
(c)	$\text{CH}_3 - \overset{\text{O}}{\parallel}{\text{C}} - \text{CH}_2 - \overset{\text{O}}{\parallel}{\text{C}} - \text{CH}_3$	(r)	8%
(d)	$\text{CH}_3 - \overset{\text{O}}{\parallel}{\text{C}} - \text{CH}_2 - \overset{\text{O}}{\parallel}{\text{C}} - \text{O} - \text{Et}$	(s)	Keto-Enol is not possible

39. Draw a most stable conformation (N - C) bond in the following compound.




40. Find total number of stereoisomers for each compound given below :



ISOMERISM**135**

41. Find the total number of stereoisomer for each compound :





42. Match the column :

Column (I)		Column (II)	
Pair		Isomeric Relationship	
(a)		(p)	Chain
(b)	$\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \overset{\text{O}}{\parallel} \text{C} - \text{OH}, \text{CH}_3 - \underset{\text{CO}_2\text{H}}{\text{CH}} - \text{CH}_3$	(q)	Positional


136


ORGANIC Chemistry for IIT-JEE


(c)		(r)	Functional
(d)		(s)	Metamers


43. Find sum of stereoisomer of following compound.

Number of stereoisomers

(a)  $\longrightarrow$ (a)

(b)  $\longrightarrow$ (b)


(c) $\text{HOCH}_2\text{---CH---CH---CH---CH}_2\text{---CH---CH---CH---CH}_2\text{---OH}$ $\longrightarrow$ (c)


(d)  $\longrightarrow$ (d)

ISOMERISM

137


44.



Prostaglandin E₁ 1 is a compound produced by the body to regulate a variety of processes including blood clotting, fever, pain and inflammation.

- A. Which of the following functional groups is not contained in 1 ?
 (a) A ketone (b) An alcohol (c) A carboxylic acid (d) An alkene
 (e) A nitrile
- B. How many asymmetric (stereogenic) centres are present in compound 1 ?
 (a) 3 (b) 4 (c) 5 (d) 6
- C. How many sp^2 hybridised carbon atoms are present in compound 1 ?
 (a) 1 (b) 2 (c) 3 (d) 4
- D. What is the geometric configuration about the double bond in compound 1 ?
 (a) E (b) Z

45.



The synthetic steroid ethynylestradiol (1) is a compound used in the birth control pill.


- A. How many sp^3 hybridised carbon atoms are present in compound (1) ?
 (a) 8 (b) 9 (c) 10 (d) 11 (e) 12
- B. How many sp^2 hybridised carbon atoms are present in compound (1) ?
 (a) 4 (b) 5 (c) 6 (d) 7 (e) 8
- C. How many sp hybridised carbon atoms are present in compound (1) ?
 (a) 2 (b) 4 (c) 6 (d) 8 (e) 10
- D. Which of the following functional group is contained in compound (1) ?
 (a) A ketone (b) An alcohol (c) A carboxylic acid (d) An ester
- E. How many asymmetric (stereogenic) centres are present in compound (1) ?
 (a) 2 (b) 3 (c) 4 (d) 5

138


ORGANIC Chemistry for IIT-JEE

SUBJECTIVE PROBLEMS

1. Number of chiral isomers are:





2.



Number of stereoisomer are

3.




(C)

Sum of number of stereoisomer (C) Degree of unsaturations in (D).


4. How many 5 membered parent chain alkane are possible for C_7H_{16} ?

5. Theoretical possible geometrical isomer of




ISOMERISM**139**

6. Total number of possible structural isomers of $C_5H_{11}Br$.
 7. Total number of plane of symmetry present in given compound is




8. Total number of isomers for $C_4H_6Br_2$ containing cyclobutane ring are (including stereoisomer) ?
 9. Total number of structural isomers of C_9H_{18} containing cyclohexane ring.
 10. How many structural isomer are possible for $C_4H_{10}O$ (only alcohol).
 11. Number of structural isomer of C_6H_{14} is .

12.



→ (x) (Number of plane of symmetry)



→ (y) Number of isomer of 1,3-dichlorocyclopentane)

Sum of (x+y=?)

13. Find out the total number of stereocentre in the given compound.
 $CH_3-CH=CH-\underset{\text{Br}}{\underset{|}{CH}}-\underset{\text{Cl}}{\underset{|}{CH}}-CH_3$
14. Find out the total number of stereoisomers of the given following compound.
 $Cl-CH=CH-\underset{\text{H}}{\underset{|}{\overset{\text{Cl}}{\text{C}}}}-CH=CH-Cl$
15. Find the total number of isomers of C_7H_{14} (only 5-membered ring).

140

ORGANIC Chemistry for IIT-JEE

ANSWERS — LEVEL 2

1. a - q; b - p; c - r; d - s
2. a - s; b - r; c - q; d - p
3. a - p, s; b - q, r, s; c - q, r, s; d - p, s
4. a - p, q; b - p, q; c - p, q; d - p, r
5. a - r; b - r; c - p; d - s
6. a - p, r; b - q, s; c - q, r; d - p, s
7. a - q, r; b - q, s; c - p, q, r; d - q, s
8. a - r, s; b - p, q; c - r, s; d - p, r, s
9. a - q; b - q, s; c - p, q, s; d - q, s
10. a - p; b - q; c - s; d - r
11. a - p; b - q; c - r; d - s
12. a - q, r; b - p; c - p; d - q, r
13. a - r; b - s; c - r; d - p
14. a - s; b - r; c - q; d - p
15. a - q, r; b - r, s; c - q, r; d - p, q
16. (a - p - x); (b - q, r - y); (c - p - x); (d - q, r - w)
17. A - b, h; B - a, g; C - c, e; D - d, f
18. A - d; B - a
19. $a + b + c + d = 13$
20. A - b, d, e; B - a, c, f, h; C - i, k, p
21. A - e, f, j; B - a, c, d, g, h, i, b; C - None
22. $w + x + y + z = 12$
23. A - (c & e), (b & e); B - (a & b) or (c & d); C - (a & c) or (a & e), (b & c), (b & e), (c & d) and (d & e)
24. A - (a & c) (b & f); B - (a & d) or (c & d), (a & e) (c & d); C - (d & e)
25. a - True; b - True; c - True; d - False
26. b, e, f, g, h, i
27. A - d, h; B - d; C - f, h; D - h
28.

Compound	A	B	C
I	c	a	a
II	c	b	b
III	c	a	a
IV	c	b	b
V	c	a	a
VI	c	a	a
VII	c	a	a
VIII	e	b	b
IX	e	a	a
X	e	b	b
29. (a) (R) (b) (L)

ISOMERISM**141****30A.** 1 - N; 2 - Z; 3 - E; 4 - Z; 5 - Z; 6 - E; 7 - N**B.** (a) 1 - Z; 2 - E; 3 - E; 4 - Z; 5 - Z; 6 - E (b) 2^9 (c) 10**31.** 1 - Z; 2 - N; 3 - E**32.** (a) A - 10, B - 4, C - 2

(c) A - 4, B - 2, C - 0

(e) A - 5, B - 2, C - 1


(g) A - 4, B - 2, C - 0

(b) A - 5, B - 0, C - 5


(d) A - 2, B - 1, C - 0

(f) A - 4, B - 1, C - 2


(h) A - 4, B - 1, C - 2

33. (1)


(2)




(3)




(4)




(5)

(7) C_3 axis, C_2 axis


(6)

(8) C_3 -axis(9) C_3 axis

(11)

(12) C_2 -axis


(10)

(13) C_2 -axis


142

ORGANIC Chemistry for IIT-JEE


34. (1)




(2)




(3)




(4)



(5)





(6)



35.

Identical Enantiomer Diastereomer Constitutional Isomer



ISOMERISM**143**


144

ORGANIC Chemistry for IIT-JEE

36. (A) - (d) (B) - (b)

37. a - p; b - q; c - r; d - s

38. a - s; b - p; c - q; d - r



40. (1) 16 (2) 4 (3) 16 (4) 4 (5) 4
 (6) 4 (7) 4 (8) 3 (9) 4 (10) 4
 (11) 2

41. a - 2, b - 4, c - 3, d - 4, e - 8, f - 2

42. a - s; b - p; c - q; d - r


43. a - 2^5 , b - 2^5 , c - $2^7 + 2^3$, d - 2^9

44. A - e; B - b; C - d; D - a

45. A - e; B - c; C - a; D - b; E - d

Subjective Problems:

1. 3 (c, f, h) 2. 6 3. 5 4. 5 5. 2 6. 8 7. 3
 8. 6 9. 12 10. 4 11. 5 12. 4 13. 4 14. 4 15. 8









3

GRIGNARD REAGENT







LEVEL - 1

1. What is the major product of the following reaction?




146


ORGANIC Chemistry for IIT-JEE



3. Reaction- 1 ;






Reaction- 2 ;



What is the ratio of (x/y) in above problem ?

- (a) 1.5 (b) 2 (c) 2.5 (d) 3

4. In which of the following reaction 2° alcohol is obtained as a product ?



(d) Both (a) and (b)




5. What product would you expect to obtain from Grignard reaction when an excess of phenylmagnesium bromide reacts with dimethyl carbonate $\text{CH}_3\text{OCOOCH}_3$?

- (a) (b) (c) (d)

Grignard Reagent






147

6. In which of the following reactions product formed is same ?







- (a) (i) and (ii) (b) (ii) and (iii) (c) (i) and (iii) (d) (i), (ii) and (iii)

7. Which of the following reaction sequences would be the best for synthesis of 2-pentanone ?







Number of moles (x) of Grignard reagent consumed in the above reaction is :

- (a) 2 (b) 3 (c) 4 (d) 5





Which of the following combinations can not be used to prepare alcohol given above ?





150

ORGANIC Chemistry for IIT-JEE


17. In which of the following reaction an acid-base reaction takes place ?



18.  Product (B) in this reaction is :




19. All of the following compounds react with ethylmagnesium bromide. Alcohols are formed from three of the compounds. Which one does not give an alcohol ?




20. A student was carrying out a Grignard reaction between PhMgBr and ethyl benzoate. She ran out of anhydrous ether just after the Grignard reagent was made. Which of the following solvents can still be used to dissolve the ethyl benzoate for its reaction with already formed PhMgBr ?

- (a) acetone (b) ethyl acetate (c) absolute alcohol (d) benzene




Number of equivalents of Grignard reagent (x) used in reaction (1) is :






- (a) 3 equivalent (b) 4 equivalent (c) 5 equivalent (d) 6 equivalent

Grignard Reagent**151**

The given product can not be obtained in the above reaction. Identify the correct product obtained.




23. Which of the following gives two isomers of 3° alcohol, when treated with phenyl magnesium bromide ?



26. When carboxylic acid reacts with organolithium reagents to give ketones, side reaction sometimes occur. For example,

152



ORGANIC Chemistry for IIT-JEE



Value of (x) in above reaction is :

- (a) 2 (b) 3 (c) 4 (d) 5

27. Which of the following alcohol can not be prepared by the reaction of acid chloride with excess of Grignard reagent followed by acidification ?





Product (B) of the above reaction is :




29. The reaction of elemental sulphur with Grignard reagent followed by acidification leads to the formation of

- (a) mercaptan (b) sulphoxide (c) thioether (d) sulphonic acid



Grignard Reagent


153




Number of moles of CH_3MgBr consumed in above reaction is :


- (a) 2 (b) 4 (c) 6 (d) 8

32. End product of the given reaction is :







33. Which of the following compound is not a suitable solvent for Grignard reaction ?



34.  Predict major product of the reaction :








35. Which of the following reaction sequences would be the best for synthesis of t-butyl alcohol ?



154

ORGANIC Chemistry for IIT-JEE

36. What is the major product of the following reaction ?



37. $\text{H} - \text{C}(\text{CH}_3) = \text{O} \xrightarrow[(2) \text{H}^+]{(1) \text{PhMgBr}}$ Products; Product obtained in this reaction are :





(a) diastereomers

(b) racemic





(c) pure enantiomer

(d) meso

38. $\text{CH}_3\text{CO}_2\text{Et} + (\text{CH}_2)_5(\text{MgBr})_2 \xrightarrow[(2) \text{H}^+]{(1) \text{PhMgBr}}$ $\text{C}_7\text{H}_{14}\text{O}$; compound (A) will be :












39. $\text{Cyclohexene} - \text{C}(=\text{O}) - \text{Ph} \xrightarrow[(2) \text{H}^+]{(1) \text{PhMgBr/CuCl}}$ (A) ; A will be : $\text{C}_{19}\text{H}_{20}\text{O}$












40. $\xrightarrow[(n\text{-Bu} = n\text{-butyl group})]{n\text{-Bu}_2\text{CuLi}}$ Product of the reaction will be :

Grignard Reagent

155



42. Ethyl acetoacetate when reacts with one mole methyl magnesium iodide then product of reaction will be :




44. For the sequence of reactions, $\text{A} \xrightarrow[\text{ether}]{\text{C}_2\text{H}_5\text{MgI}} \text{B} \xrightarrow{\text{H}_2\text{O}/\text{H}^+} \text{tert-Pentyl alcohol}$. The compound A in the sequence is :

(a) 2-Butanone




(b) Acetaldehyde

(c) Acetone

(d) Propanal



Same product (A) will form in both reactions. A is :




156

ORGANIC Chemistry for IIT-JEE

46. Which of the following Grignard reagent can be prepared ?

- (a) $\text{Br} - \text{Mg} - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{O} - \text{H}$ (b) $\text{Br} - \text{Mg} - \text{CH}_2 - \text{CH}_2 - \text{SH}$
 (c) $\text{BrMg} - \text{CH}_2 - \text{CH}_2 - \text{NH}_2$ (d) $\text{BrMg} - \text{CH}_2 - \text{CH}_2 - \text{N}(\text{CH}_3)_2$

47. In the reaction sequence :



- (a) (b) (c) (d)

48. $(\text{C}_2\text{H}_5\text{O})_2\text{CO} \xrightarrow[\text{H}_3\text{O}^+]{\text{CH}_3\text{MgBr}(\text{excess})} \text{A}$. A (alcohol) can also be obtained by :

- (a) $\text{CH}_3\text{CH}_2\text{CHO} \xrightarrow[\text{H}_3\text{O}^+]{\text{CH}_3\text{MgBr}(2\text{mol})}$ (b) $\text{CH}_3\text{C}(=\text{O})\text{OC}_2\text{H}_5 \xrightarrow[\text{H}_3\text{O}^+]{\text{CH}_3\text{MgBr}(2\text{mol})}$
 (c) $\text{CH}_3\text{C}(=\text{O})\text{CH}_3 \xrightarrow[\text{H}_3\text{C}]{\text{CH}_3\text{MgBr } 1\text{mol}} \rightarrow$ (d) as in (b) and (c)


49. The principal product of the reaction between methyl butanoate and 2 moles of CH_3MgBr after hydrolysis is :

- (a) $\text{C}_3\text{H}_7\text{COCH}_3$ (b) $\text{C}_3\text{H}_7\text{C}(\text{OH})(\text{CH}_3)_2$
 (c) $\text{C}_3\text{H}_7\text{CHOHCH}_3$ (d) $\text{C}_3\text{H}_7\text{COCH}(\text{CH}_3)_2$

50. Which of the following compounds will form hydrocarbon on reaction with Grignard reagent ?

- (a) $\text{CH}_3\text{CH}_2\text{OH}$ (b) CH_3CHO (c) CH_3COCH_3 (d) $\text{CH}_3\text{CO}_2\text{CH}_3$





51. What is the product (B) of the following reaction sequence ?




- (a) (b) (c) (d)

Grignard Reagent**157**


52. Which, if any, of the following pairs of reagents could be used to prepare 2-phenyl-2-butanol?



53. What is the product of the following reaction?




54.




- (a) $\text{Cyclopropylidenetriphenylphosphorane}$
- (b) $\text{Ph}-\text{C}\equiv\text{C}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{OH}$
- (c) $\text{Ph}-\text{C}\equiv\text{C}-\text{CH}_2-\text{CH}_2-\text{OH}$
- (d) $\text{Ph}-\text{CH}_2-\text{C}\equiv\text{C}-\text{CH}_2-\text{CH}_2-\text{OH}$

55. What sequence of steps represents the best synthesis of 4-heptanol ($\text{CH}_3\text{CH}_2\text{CH}_2$)₂CHOH?

- (a) $\text{CH}_3\text{CH}_2\text{CH}_2\text{MgBr}$ (2 moles) + formaldehyde ($\text{H}_2\text{C}=\text{O}$) in diethyl ether followed by H_3O^+
- (b) $\text{CH}_3\text{CH}_2\text{CH}_2\text{MgBr}$ + butanal ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}=\text{O}$) in diethyl ether followed by H_3O^+
- (c) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{MgBr}$ + acetone [$(\text{CH}_3)_2\text{C}=\text{O}$] in diethyl ether followed by H_3O^+
- (d) $(\text{CH}_3\text{CH}_2\text{CH}_2)_2\text{CHMgBr}$ + formaldehyde ($\text{H}_2\text{C}=\text{O}$) in diethyl ether followed by H_3O^+



- (a) diastereomers
- (b) racemic
- (c) single stereoisomer
- (d) meso




What is the value of x in the above reaction ?

- (a) 1 (b) 2
(c) 3 (d) 4


- 58.** 0.40 g of an organic compound (A), (M.F.- C_5H_8O) reacts with x mole of CH_3MgBr to liberate 224 mL of a gas at STP. With excess of H_2 , (A) gives pentan-1-ol. The correct structure of (A) is :

- (a) $\text{CH}_3 - \text{C} \equiv \text{C} - \text{CH}_2 - \text{CH}_2 - \text{OH}$
 (b) $\text{CH}_3 - \text{CH}_2 - \text{C} \equiv \text{C} - \text{CH}_2 - \text{OH}$
 (c) $\text{H} - \text{C} \equiv \text{C} - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{OH}$
 (d) $\text{H} - \text{C} \equiv \text{C} - \text{CH}_2 - \text{CH} - \text{CH}_3$



End product (X) of the above reaction is

- (a) $\text{CH}_2 = \text{CH} - \text{CH}_2 - \overset{\text{CH}}{\underset{\parallel}{\text{C}}} - \text{CH}_3$
- (b) $\text{H}_2\text{C} = \text{CH} - \text{CH} = \underset{\text{CH}_3}{\underset{|}{\text{C}}} - \text{CH}_3$
- (c) $\text{H}_2\text{C} = \text{CH} - \text{CH}_2 - \overset{\text{OH}}{\underset{\text{CH}_3}{\underset{|}{\text{C}}}} - \text{CH}_3$
- (d) $\text{H}_2\text{C} = \text{CH} - \text{CH}_2 - \text{CH} - \underset{\text{CH}_3}{\underset{|}{\text{CH}_2}} - \text{OH}$




- (a) $\text{CH}_2 - \underset{\text{O}}{\text{CH}} - \text{CH}_2 - \text{CH}_3$ (b) $\text{CH}_3 - \text{O} - \text{CH}_2 - \text{CH}_2 - \text{CH}_3$
- (c) $\text{H}_2\text{C} = \text{CH} - \text{CH}_2 - \text{O} - \text{CH}_3$ (d) $\text{H}_2\text{C} - \underset{\text{O}}{\text{CH}} - \text{CH}_3$





Grignard Reagent

159





61. Compound A was treated with a large excess of CH_3MgBr . The resulting product was exposed to $\text{POCl}_3/\text{pyridine}$ to give compound B, as one of many products :



Which of the following compounds can be A ?



62. Identify product Z in the following reaction sequence :



160**ORGANIC Chemistry for IIT-JEE****ANSWERS — LEVEL 1**

1.	(b)	2.	(d)	3.	(b)	4.	(d)	5.	(c)	6.	(d)	7.	(c)	8.	(c)
9.	(d)	10.	(c)	11.	(b)	12.	(b)	13.	(b)	14.	(c)	15.	(b)	16.	(b)
17.	(d)	18.	(d)	19.	(d)	20.	(d)	21.	(c)	22.	(b)	23.	(b)	24.	(c)
25.	(c)	26.	(b)	27.	(d)	28.	(d)	29.	(a)	30.	(b)	31.	(b)	32.	(b)
33.	(d)	34.	(b)	35.	(c)	36.	(c)	37.	(b)	38.	(b)	39.	(b)	40.	(a)
41.	(b)	42.	(c)	43.	(b)	44.	(c)	45.	(c)	46.	(d)	47.	(b)	48.	(d)
49.	(b)	50.	(a)	51.	(a)	52.	(a)	53.	(c)	54.	(b)	55.	(b)	56.	(a)
57.	(c)	58.	(c)	59.	(b)	60.	(c)	61.	(d)	62.	(a)				

ATDB.uno


Grignard Reagent

161




1. Comprehension

Grignard reagent is usually prepared by





Grignard reagent




Grignard reagent

Grignard reagent acts as a strong base. Grignard reagent carry out nucleophilic attack in absence of acidic hydrogen. Grignard reagent form complex with its ether solvent. Complex formation with molecule of ether is an important factor in the formation and stability of Grignard reagent.













A. What is the correct order of reactivity of halides with magnesium ?



B. Which of the following will undergo acid-base reaction with Grignard reagent ?



C. Which of the following reactants give primary alcohol as a major product when reacts with $RMgX$ followed by acidification ?



162

ORGANIC Chemistry for IIT-JEE

F. Which of the following Grignard reagents is not possible ?

- (a) $\text{HS}-\text{CH}_2-\text{CH}_2-\text{CH}_2\text{MgBr}$ (b) $\text{HO}-\text{CH}_2-\text{CH}_2-\text{CH}_2\text{MgBr}$
 (c) $\text{NH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2\text{MgBr}$ (d) All of these

G. How many different Grignard reagents when react with EtOH, give *n*-butane as product (excluding stereoisomerism).

- (a) 1 (b) 2 (c) 3 (d) 4

2. Match the column I and II. (Matrix)

Column (I)		Column (II)	
Reactant		Product	
(a)	$\text{PhMgBr} + \text{Cl}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{Et} \xrightarrow{\text{H}^+}$ (excess)	(p)	$\text{Ph}-\text{CH}_2-\text{OH}$
(b)	$\text{PhMgBr} + \text{H}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{Et} \xrightarrow{\text{H}^+}$ (excess)	(q)	$\begin{array}{c} \text{Ph}-\text{CH}-\text{Ph} \\ \\ \text{OH} \end{array}$
(c)	$\text{PhMgBr} + \text{H}-\overset{\text{O}}{\parallel}{\text{C}}-\text{H} \xrightarrow{\text{H}^+}$ (excess)	(r)	$\begin{array}{c} \text{OH} \\ \\ \text{Ph}-\text{C}-\text{Ph} \\ \\ \text{Ph} \end{array}$
(d)	$\text{PhMgBr} + \text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{Et} \xrightarrow{\text{H}^+}$ (excess)	(s)	$\begin{array}{c} \text{OH} \\ \\ \text{Ph}-\text{C}-\text{Ph} \\ \\ \text{CH}_3 \end{array}$


3. Match the column I and II. (Matrix)

Column (I)		Column (II)	
Reaction		Reactant	
(a)	$\text{PhMgBr} + (\text{A}) \xrightarrow{\text{H}^+} 1^\circ\text{alcohol}$	(p)	$\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_3$
(b)	$\text{PhMgBr} + (\text{B}) \xrightarrow{\text{H}^+} 2^\circ\text{alcohol}$	(q)	$\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_3$
(c)	$\text{PhMgBr} + (\text{C}) \xrightarrow{\text{H}^+} 3^\circ\text{alcohol}$	(r)	$\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{H}$
(d)	$\text{PhMgBr} + (\text{D}) \xrightarrow{\text{H}^+} \text{C}_6\text{H}_6$	(s)	$\text{H}-\overset{\text{O}}{\parallel}{\text{C}}-\text{H}$




Match the missing reactant A, B, C, D

Grignard Reagent**163**

4. Match the column I and II. (Matrix)

Column (I)		Column (II)	
Reaction		Moles of PhMgBr used	
(a)	$\text{PhMgBr} + \text{Et}-\text{O}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{Et} \xrightarrow{\text{H}^+} 3^\circ\text{alcohol}$	(p)	1
(b)	$\text{PhMgBr} + \text{HO}-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_3 \xrightarrow{\text{H}^+} 3^\circ\text{alcohol}$	(q)	2
(c)	$\text{PhMgBr} + \text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_3 \xrightarrow{\text{H}^+} 3^\circ\text{alcohol}$	(r)	3
(d)	$\text{PhMgBr} + $  $\xrightarrow{\text{H}^+} 3^\circ\text{alcohol}$	(s)	4

5. When 20 g of a compound (M.W. = 44, O_4) reacts with excess of CH_3MgBr , 14.6 L of CH_4 is obtained at STP. What is the structural formula of (A)?




(d) Both (a) & (b)

164

ORGANIC Chemistry for IIT-JEE

SUBJECTIVE PROBLEMS

1.




How many geometrical isomer of (X) is possible ?

2. How many isomer of C_4H_8O when reacts with CH_3MgBr followed by acidification to give 2° alcohol (only consider carbonyl isomers)?
(including stereoisomer)

3.

Total number of $RMgX$ are consumed in the following reaction




4. How many isomers of $C_4H_{10}O$ reacts with CH_3MgBr to evolve CH_4 gas ? (Excluding stereoisomer)

ANSWERS — LEVEL 2

1. A – c; B – d; C – a; D – b; E – b; F – d; G – b
2. a – r; b – q; c – p; d – s
3. a – s; b – r; c – q; d – p
4. a – r; b – q; c – p; d – s
5. (d)

Subjective Problems

1. 4 2. 2 3. 7 4. 4

**4A**

HYDROCARBONS (ALKANES)




- On halogenation, an alkane gives only one monohalogenated product. The alkane may be :
 - 2-methyl butane
 - 2, 2-dimethyl propane
 - cyclopentane
 - both (b) and (c)
- Which of the following compounds can be best prepared by Wurtz-reaction ?
 - Iso-butane
 - n*-butane
 - n*-pentane
 - Iso-pentane
- A hydrocarbon A (V.D. = 36) forms only one monochloro substitution product. A will be :
 - iso-pentane
 - neo-pentane
 - cyclohexane
 - methyl-cyclohexane
- Ethyl iodide and *n*-propyl iodide are allowed to undergo Wurtz reaction. The alkane which will not be obtained in this reaction is :
 - butane
 - propane
 - pentane
 - hexane
- $$\text{CH}_3 - \underset{\text{CH}_3}{\text{CH}} - \text{CH}_2 - \text{CH}_3 \xrightarrow[\text{h}\nu]{\text{Cl}_2}$$

Number of chiral centers generated during monochlorination in the above reaction :

- 1
- 2
- 3
- 4


166

ORGANIC Chemistry for IIT-JEE




Above conversion can be achieved by :

- (a) Zn / H^+ (b) LiAlH_4
 (c) $\text{Mg} / (\text{ether})$ then H_2O (d) all of these




Give the total number of monochloro products(including stereoisomers), which are possible in the above reaction.

- (a) 2 (b) 3 (c) 4 (d) 5

To obtain high yields of CH_3Cl , the ratio of CH_4 to Cl_2 must be :

- (a) high (b) low
 (c) equal (d) can't be predicted


9. Double bond equivalent of cubane is :




Cubane

- (a) 4 (b) 5 (c) 6 (d) 7

10. How many bond cleavages are required to convert cubane to a non-cyclic skeleton ?






Cubane



Non-cyclic skeleton

- (a) 2 (b) 3 (c) 4 (d) 5

11. Draw an energy profile diagram for a three step reaction in which first step is slowest and last step is fastest. (Assume that reaction is exothermic)




(d) None of these



HYDROCARBONS (ALKANES)**167**

12. $\text{CH}_3-\underset{\text{CH}_3}{\text{CH}}-\text{CH}_2-\text{CH}_3 \xrightarrow[h\nu]{\text{Cl}_2} (x) = \text{Number of monochloro product including stereoisomers.}$

(a) 4 (b) 5 (c) 6 (d) 7




Product (P) is :




Double bond equivalent (degree of Unsaturation) of (A) is :

(a) 1 (b) 2 (c) 3 (d) 4

15. Arrange the following alkanes in decreasing order of their heats of combustion.




(a) (i) > (ii) > (iii) (b) (iii) > (i) > (ii)
(c) (iii) > (ii) > (i) (d) (i) > (iii) > (ii)



Product of the above reaction will be :



(a) Racemic mixture (b) Diastereomers
(c) Meso (d) Constitutional isomers





Which of the following compound will not be obtained as a product in the above reaction ?

168


ORGANIC Chemistry for IIT-JEE




18. Following are the structures of four isomer of hexane. Among the names given below, which correctly identifies the fifth isomer ?



- (a) 2-Methyl pentane
(b) 2-Ethyl butane
(c) 2,3-Dimethyl butane
(d) 3-Methyl pentane
19. Which of the following describes the relationship between the methyl groups in the chair conformation of the substance shown below ?




- (a) Trans
(b) Anti
(c) Gauche
(d) Eclipsed
20. Compare the stabilities of the following two compounds (A) and (B):
A : *cis*-1-ethyl-3-methyl cyclohexane
B : *trans*-1-ethyl-3-methyl cyclohexane
(a) A is more stable
(b) B is more stable
(c) A and B are of equal stability
(d) No comparison can be made
21. Which conformation of ethane has the lowest potential energy ?
(a) Eclipsed
(b) Skew
(c) Staggered
(d) All will have equal potential energy
22. Ethane is subjected to combustion process. During the combustion the hybrid state of carbon changes from :
(a) sp^2 to sp^3
(b) sp^3 to sp
(c) sp to sp^3
(d) sp^2 to sp^2



Above reaction is an example of :


HYDROCARBONS (ALKANES)**169**

- (a) isomerization (b) polymerization
(c) cracking (d) de-hydrogenation
24. Which of the following has highest chlorine content ?
(a) Pyrene (b) DDT (c) Chloral (d) Gammaxene
25. Pure methane can be prepared by :
(a) Wurtz reaction (b) Kolbe electrolysis method
(c) soda-lime de-carboxylation (d) reduction with H_2
26. Calcium carbide + heavy water $\longrightarrow$?
The product of the above reaction is :
(a) C_2H_2 (b) CaD_2 (c) $Ca(OD)_2$ (d) CD_4




Arrange the compounds I, II and III in decreasing order of their heats of combustion:

- (a) $II > I > III$ (b) $I > II > III$
(c) $III > II > I$ (d) $III > I > II$
28. An alkane (mol. wt. = 86) on bromination gives only two monobromo derivatives (excluding stereoisomers). The alkane is :



29. Order of the bond strength of C — H bonds involving sp , sp^2 and sp^3 hybridized carbon atoms is :
(a) $sp > sp^2 > sp^3$ (b) $sp^3 > sp^2 > sp$
(c) $sp^2 > sp^3 > sp$ (d) $sp^2 > sp > sp^3$



Among the structures given, select the enantiomers :

- (a) I and II (b) I and III
(c) II and III (d) I, II and III

170

ORGANIC Chemistry for IIT-JEE

31.



(I)



(II)




(III)





The correct order of reactivity of I, II & III towards addition reactions is :

- (a) $I > III > II$ (b) $I > II > III$ (c) $III > II > I$ (d) $III > I > II$

32.



Product (A) of above reaction is :




33. Which of the following reactants is suitable for preparation of methane and ethane by using one step only ?

- (a) $H_2C = CH_2$ (b) CH_3OH
(c) $CH_3 - Br$ (d) $CH_3 - CH_2 - OH$

34. How many carbon atoms does an alkane (not a cycloalkane) need before it is capable to exist in enantiomeric form ?

- (a) 4 (b) 5 (c) 6 (d) 7

35. Among the following free radical bromination reactions, select those in which 2° halide is the major product —



- (a) P, Q, R, S (b) P, R, U (c) P, R, S, T (d) P, Q, R, S, T

36. (A) + $Cl_2 \xrightarrow{h\nu}$ monochloro product

To maximise the yield of monochloro product in the above reaction ?

- (a) Cl_2 must be added in excess
(b) Reactant (A) must be added in excess
(c) Reaction must be carried out in dark
(d) Reaction must be carried out with equimolar mixture of Cl_2 and A

37. $CH_3 - CH_2 - CH_2 - CH_3 \xrightarrow{Br_2/h\nu}$






Major product in the above reaction is :

- (a) Racemic mixture (b) Meso
(c) Diastereomers (d) Constitutional isomers

HYDROCARBONS (ALKANES)

171

38. Select the chain propagation steps in the free-radical chlorination of methane.




(a) 2, 3, 5

(b) 1, 3, 6

(c) 3, 5

(d) 2, 3, 4




The number of possible monobromo products is (excluding stereoisomers):

(a) 4

(b) 5

(c) 8

(d) 10



$\text{Br}^\bullet$ will abstract which of the following most readily?


(a) a

(b) b


(c) c

(d) d

41. Arrange the following compounds in decreasing order of their heats of combustion :



(i)



(ii)




(iii)

(a) (iii) > (ii) > (i)

(b) (ii) > (i) > (iii)

(c) (iii) > (i) > (ii)

(d) (i) > (ii) > (iii)



Arrange the hydrogens a, b, c, d, in decreasing order of their reactivities towards chlorination:

(a) $a > b > c > d$ (b) $b > c > d > a$ (c) $b > c > a > d$ (d) $c > b > a > d$

43. On catalytic reduction (H_2/Pt) how many alkenes will give n-butane ?

(a) 1

(b) 2

(c) 3

(d) 4

44. On catalytic reduction (H_2/Pt) how many alkenes will give 2-methylbutane ?

(a) 1


(b) 2

(c) 3

(d) 4


172

ORGANIC Chemistry for IIT-JEE




How many dichloro products are formed in the above reaction (including stereoisomers)?

- (a) 5 (b) 6
(c) 7 (d) 9




Product of the above reaction will be :

- (a) Racemic mixture (b) Diastereomers
(c) Meso (d) Constitutional isomers






Product of the above reaction will be :

- (a) Diastereomers (b) Racemic mixture
(c) Meso (d) Constitutional isomers






Products obtained in above Wurtz reaction is :



- (d) Both (a) and (b)


49. Rank the transition states that occur during the following reaction steps in order of increasing stability (least $\rightarrow$ most stable) :



- (a) $1 < 2 < 3$ (b) $2 < 3 < 1$
(c) $1 < 3 < 2$ (d) $2 < 1 < 3$

HYDROCARBONS (ALKANES)**173**

50. Which of the following does not represent major product of that reaction ?



ATDB.uno

ANSWERS — LEVEL 1

1.	(d)	2.	(b)	3.	(b)	4.	(b)	5.	(b)	6.	(d)	7.	(b)	8.	(a)
9.	(b)	10.	(d)	11.	(c)	12.	(c)	13.	(b)	14.	(c)	15.	(c)	16.	(b)
17.	(d)	18.	(d)	19.	(c)	20.	(a)	21.	(c)	22.	(b)	23.	(a)	24.	(a)
25.	(c)	26.	(c)	27.	(c)	28.	(c)	29.	(a)	30.	(c)	31.	(b)	32.	(b)
33.	(c)	34.	(d)	35.	(b)	36.	(b)	37.	(a)	38.	(c)	39.	(b)	40.	(a)
41.	(d)	42.	(c)	43.	(c)	44.	(c)	45.	(c)	46.	(a)	47.	(a)	48.	(d)
49.	(c)	50.	(d)												


174

ORGANIC Chemistry for IIT-JEE



1. Comprehension



For the given question (1, 2, 3), consider the following reaction.




- A. Light is involved in which step of the reaction :
 - (a) Initiation only
 - (b) Termination only
 - (c) Propagation only
 - (d) Propagation and Termination
- B. Which halogen will give the best yield of a single monohalogenation product ?
 - (a) F_2
 - (b) Cl_2
 - (c) Br_2
 - (d) I_2
- C. How many monohalo derivatives are possible (excluding stereoisomers) ?
 - (a) 3
 - (b) 4
 - (c) 5
 - (d) 6

2. Comprehension

Halogenation is a substitution reaction where halogen replaces one or more hydrogens of hydrocarbon.



Chlorination is exothermic and transition state resembles with products



Bromination is endothermic and transition state resembles with products

Chlorine free radical make 1° , 2° , 3° radicals with almost equal ease, whereas bromine free radicals have a clear preference for the formation of tertiary free radicals. So, bromine is less reactive, and more selective whereas chlorine is less selective and more reactive.

The relative rate of abstraction of hydrogen by $\text{Br}^\cdot$ is

$$3^\circ > 2^\circ > 1^\circ$$

$$(1600) \quad (82) \quad (1)$$


The relative rate of abstraction of hydrogen by $\text{Cl}^\cdot$ is :

$$3^\circ > 2^\circ > 1^\circ$$

$$(5) \quad (3.8) \quad (1)$$


HYDROCARBONS (ALKANES)**175****Consider the above argument and answer A to G :**

- A.** 1-halo-2,3-dimethyl butane will be obtained in better yields, if halogen is :
 (a) Br₂ (b) Cl₂ (c) I₂ (d) Can't be predicted







Above product will obtained in better yield if X is

- (a) Cl₂ (b) I₂ (c) Br₂ (d) Can't be predicted




Major product in the above reaction is :

- (a)  (b) 
 (c)  (d) 

- D.** Which of the following will give five monochloro products, when allowed to react with Cl₂ in presence of sun light (excluding free radical) :


- (a) n-pentane (b) Iso-pentane (c) 2-methyl-pentane (d) 3-methyl pentane



What is the value of x (% yield of product)?

- (a) 18 % (b) 82 % (c) 90 % (d) 60 %

- F.** What would be the product ratio x/y in the chlorination of propane if all the hydrogen were abstracted at equal rate ?









- (a) $\frac{1}{3}$ (b) $\frac{3}{1}$ (c) $\frac{9}{1}$ (d) $\frac{1}{9}$

- G.** How many dichloro products (including stereoisomers) will be formed when R-2-chloropentane reacts with Cl₂ in presence of UV radiation ?


- (a) 5 (b) 6 (c) 7 (d) 8

176

ORGANIC Chemistry for IIT-JEE



- (a) Meso compound (b) Racemic mixture (c) Diastereomers (d) Optically active



Product (A) is :

- (a) $\text{Ph}-\text{CO}_2\text{H}$ (b) $\text{Ph}-\text{CH}=\text{OP}$ (c) $\text{Ph}-\text{CH}=\text{CH}_2$ (d) 

6. Match the column I with column II and with column III.





Column (I)		Column (II)		Column (III)	
Compound		Mono-chloro products		Monochloro products	
		(excluding stereoisomerism)		(including stereoisomerism)	
(a)		(p)	1	(w)	1
(b)	$\text{CH}_3-\text{CH}(\text{CH}_3)-\text{CH}_2-\text{CH}_3$	(q)	2	(x)	3
(c)	$\begin{array}{c} \text{CH}_3 \text{ CH}_3 \\ \quad \\ \text{CH}_3-\text{C}-\text{C}-\text{CH}_3 \\ \quad \\ \text{CH}_3 \text{ CH}_3 \end{array}$	(r)	3	(y)	5
(d)	$\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{CH}_3$	(s)	4	(z)	6

HYDROCARBONS (ALKANES)**177**

7.

A.	$R\text{-}2\text{-chloropentane} \xrightarrow[h\nu]{\text{Cl}_2}$ Optically active di-chloro products (<i>P</i>)
B.	 $\xrightarrow[h\nu]{\text{Cl}_2}$ Optically active dichloro products (<i>Q</i>)
C.	$R\text{-}2\text{-chlorobutane} \xrightarrow[h\nu]{\text{Cl}_2}$ Optically active di-chloroproducts (<i>R</i>)









Sum *P* + *Q* + *R* is :**8. Match the column I and II.**

Column (I)		Column (II)	
Reaction		Type of Reaction	
(a)		(p)	Meso compound
(b)		(q)	Diastereomers
(c)		(r)	Racemic
(d)		(s)	Optically inactive due to absence of chiral center


178

ORGANIC Chemistry for IIT-JEE

9. Match the column :

Column (I)		Column (II)	
Reaction		Product	
(a)		(p)	
(b)		(q)	
(c)		(r)	
(d)		(s)	

10. How many distinct monochlorinated products, (including stereoisomers) may be obtained when the alkane shown below is heated in the presence of Cl_2 ?



- (a) 1 (b) 2 (c) 3 (d) 4 (e) 6

11. How many distinct monochlorinated products, (including stereoisomers) may be obtained when the alkane shown below is heated in the presence of Cl_2 ?



- (a) 2 (b) 4 (c) 5 (d) 6 (e) 8


HYDROCARBONS (ALKANES)

[illegible]

179

12. Match the column :

Column (I)		Column (II)	
	Wurtz reaction		Number of dimerization product
(a)	$\text{CH}_3 - \text{Cl} \xrightarrow[\text{dry ether}]{\text{Na}}$	(p)	5
(b)	$\text{CH}_3 - \text{Cl} + \text{CH}_3 - \text{CH}_2 - \text{Cl} \xrightarrow[\text{dry ether}]{\text{Na}}$	(q)	6
(c)	$\text{CH}_3 - \text{Cl} + \text{CH}_3 - \text{CH}_2 - \text{Cl} + \text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{Cl} \xrightarrow[\text{dry ether}]{\text{Na}}$	(r)	3
(d)	$\text{H}_2\text{C} = \text{CH} - \text{CH} = \text{CH} - \text{CH}_2 - \text{Cl} + \text{CH}_3 - \text{CH}_2 - \text{Cl} \xrightarrow[\text{dry ether}]{\text{Na}}$	(s)	1




180

ORGANIC Chemistry for IIT-JEE

ANSWERS — LEVEL 2

1. A – a; B – c; C – b
2. A – b; B – c; C – a; D – c; E – c; F – b; G – c
3. a, b, c
4. a
5. c
6. a – q – x; b – s – z; c – p – w; d – q – x
7. $P + Q + R = 10$
8. a – q; b – p; c – r; d – s
9. a – p; b – s; c – q; d – r
10. a
11. e
12. a – s; b – r; c – p; d – q
13. 9




ATDB.uno

**4B**

HYDROCARBONS (ALKENES)






1. (R)-3-bromocyclopentene (shown below) reacts with Br_2/CCl_4 to form two products, Y and Z, Y is not optically active (does not rotate plane-polarized light). What is the structure of Y?






182


ORGANIC Chemistry for IIT-JEE



Which of the following products cannot be obtained in ozonolysis of *o*-xylene?








Product (A) of the reaction is :







HYDROCARBONS (ALKENES)






183







9. The reaction of propene with H_3O^+ will proceed with which of the following intermediates ?




10. Which of the following bromides is the major product of the reaction shown below, assuming that there are no carbocation rearrangement ?



11. Which of the following reactions results in the formation of a pair of diastereomers ?




12. What is a likely product of the reaction shown ?




184


ORGANIC Chemistry for IIT-JEE



13. Which of the following, when undergoing addition of HBr, will form ONLY a pair of diastereomers ?




14. How many transition states and intermediates will be formed during the course of following reaction ?




- (a) 3 transition states and 3 intermediates (b) 4 transition states and 3 intermediates
(c) 3 transition states and 2 intermediates (d) 5 transition states and 4 intermediates


15. Product of which of the following reactions, is racemic mixture ?



16. The product(s) of the following reaction can best be described as :




- (a) a racemic mixture (b) a single enantiomer
(c) a pair of diastereomers (d) an achiral molecule
17. Taking into account the stability of various carbocations and, as well as the rules governing mechanisms of carbocation rearrangements, which reaction is most likely to occur during the given reaction ?




HYDROCARBONS (ALKENES)


185




18. Consider the following reaction in which the intermediate carbocation loses H^+ to give the final product ?




Which of the following energy profiles best represents the overall reaction ?



19. Methyl vinyl ether, $H_2C=CH-OCH_3$, reacts with Br_2/CH_3OH . If methanol is reacting as water would, and if this reaction follows a typical mechanism of electrophilic addition, what would be the expected product ?




20. 2, 4-hexadiyne (C_6H_6) is allowed to react with Li in $NH_3(lig)$. The product obtained is treated with 1 equivalent of Cl_2 in CCl_4 . Which of the following constitutional isomers are possible products ?



- (a) I and II
(b) II and III
(c) I and V
(d) I and III


21. Which of the following is the best stereochemical representation when reaction between 1-methylcyclohexene and NBS react in aqueous dimethyl sulfoxide ?




186

ORGANIC Chemistry for IIT-JEE

22. Which of the following is among the major products of the reaction of (E)-3-methyl-2-pentene with BH_3 in THF followed by the addition of $\text{H}_2\text{O}_2/\text{HO}^-$?




23. Compare rate of dehydration of (i), (ii) and (iii) by conc. H_2SO_4 .




- (a) (i) > (iii) > (ii) (b) (i) > (ii) > (iii)
(c) (ii) > (i) > (iii) (d) (ii) > (iii) > (i)

24. How many products will be formed in this reaction ?




- (a) 10 (b) 2
(c) 3 (d) 4

25. (A). Product (A) of the reaction is:




26. $\text{CH}_3-\text{CH}(\text{CH}_3)_2 + \text{H}_2\text{C}=\text{CH}_2 \xrightarrow[2-5^\circ\text{C}]{\text{HF}} (\text{A})$; (A) is:





HYDROCARBONS (ALKENES)

187


27. Predict the product (A) of the following reaction




28.




29. Di-imide (N_2H_4) is used to reduce double bond of:




30.




End product of the reaction is:




31.




Product (A) is:



32.




Compound (A) is:




190

ORGANIC Chemistry for IIT-JEE


40. The major product of the following reaction sequence is :



41. Which one of the following compounds gives acetone $(CH_3)_2C=O$ as one of the product of its ozonolysis ?




42. Addition of HCl to 3, 3-dimethyl-1-butene yields two products, one of which has a rearranged carbon skeleton. Among the following carbocations, select the possible intermediates in that reaction ?




43. Conversion of cyclohexene to cyclohexane can be conveniently achieved by :




44. *Trans*-cyclohexane-1,2-diol can be obtained by the reaction of cyclohexene with :



45. Bromination of (E)-2-butenedioic acid gives









46. The major product formed during the reaction of 1-methyl cyclopentene with CH_3CO_3H is







47. $CH_2=CH-CO_2H \xrightarrow[2. H_2O_2, HO^-]{1. NaOH} (A) \xrightarrow{\text{electrolysis}} (B)$; Product (B) of the reaction is :
- (a) CH_3-CH_3 (b) $H_2C=CH_2$
(c) $H-C \equiv C-H$ (d) $CH_2=CH-CH=CH_2$

HYDROCARBONS (ALKENES)










191



Cyclohexene








50. What is the major product expected from the following reaction ?






192

ORGANIC Chemistry for IIT-JEE



54. Which compound is a possible product from addition of Br_2 to 1-butene ?



55. Addition of Br_2 to *cis*-2-butene would give a product which is:

(a) achiral

(b) racemic

(c) meso

(d) optically active

56. Addition of Br_2 to *trans*-2-butene would give a product which is:

(a) achiral

(b) racemic

(c) meso

(d) optically active

57. Addition of OsO_4 to cyclopentene would give a product which is:

(a) achiral

(b) racemic

(c) meso

(d) optically active


58. Addition of BH_3 followed by H_2O_2 to *trans*-2-butene would give a product which is:

(a) achiral

(b) racemic

(c) meso

(d) optically active





(a) $\text{H}_2\text{O}/\text{H}^+$

(b) $\text{BH}_3 \cdot \text{THF}/\text{H}_2\text{O}_2 - \text{OH}^-$

(c) $\text{Hg}(\text{OCOCH}_3)_2 \cdot \text{THF}/\text{NaBH}_4 \cdot \text{NaOH}$





(d) All are possible

60. The major product of the following reaction is :








HYDROCARBONS (ALKENES)

193








62. Which of the following is a major product of the reaction shown below?




63. In methyl alcohol solution, bromine reacts with ethylene (ethene) to yield $\text{BrCH}_2\text{CH}_2\text{OCH}_3$ in addition to 1, 2-dibromoethane because
- the methyl alcohol solvates the bromine
 - the ion formed initially may react with Br^- or $\text{C}_2\text{H}_5\text{OH}$
 - this is a free radical reaction
 - the reaction follows Markovnikov's rule

64. Which of the following compound was the starting material for the oxidation shown below?



65. Which series of reactions will achieve the following transformation?



- 1
(a) Cl_2/CCl_4
(c) Cl_2/CCl_4

- 2
(b) Br_2
(d) $\text{NBS}/h\nu$


- 1
(b) HBr
(d) $\text{NBS}/h\nu$

- 2
(c) Cl_2/CCl_4
(d) Cl_2/CCl_4


194

ORGANIC Chemistry for IIT-JEE


66. Taking into account the stability of various cycloalkanes and carbocations, as well as the rules governing mechanisms of carbocation rearrangements, what is the most likely product of this reaction ?




67. A triene is treated with ozone followed by zinc in acetic acid to give the following three products. What is the structure of the triene ?



68. Which of the following compound would yield trialkylborane shown below when treated with BH_3/THF ?



- (a) 2-methylbut-1-ene (b) 2-methylbut-2-ene
(c) 3-methylbut-1-ene (d) 3-methylbut-1-yne
69. If the following compound is treated with Pd/C in excess of hydrogen gas, how many stereoisomers of the product will be obtained ?






HYDROCARBONS (ALKENES)

195


- (a) 1 (b) 2
(c) 3 (d) 4

70. Which is the most precise designation of stereochemistry for the products formed in the electrophilic addition of DBr to 1-methylcyclohexene ? (D = ^2H , an isotope of hydrogen)



(d) both (a) and (b)

71. Consider the addition of HBr to 3,3-Dimethyl-1-butene shown below. What is the best mechanistic explanation for the formation of the observed product ?



- (a) Protonation of the alkene followed by a hydride shift and addition of bromide to the carbocation
(b) Double bond shift in the alkene following by the protonation and addition of bromide to the carbocation
(c) Addition of bromide to the alkene followed by a double bond shift and protonation
(d) Protonation of the alkene followed by a methyl shift and addition of bromide to the carbocation


72. Propene $\text{CH}_3\text{CH} = \text{CH}_2$ can be converted into 1-propanol by oxidation. Indicate which sets of reagents amongst the following is ideal to effect the above conversion ?

- (a) KMnO_4 (alkaline) (b) Osmium tetroxide ($\text{OsO}_4/\text{CH}_2\text{Cl}_2$)
(c) B_2H_6 and alk. H_2O_2 (d) O_3/Zn

73. Which is the most suitable reagent among the following distinguish compound (3) from the others ?

- (1) $\text{CH}_3\text{C} \equiv \text{C} - \text{CH}_3$ (2) $\text{CH}_3\text{CH}_2 - \text{CH}_2 - \text{CH}_3$
(3) $\text{CH}_3\text{CH}_2\text{C} \equiv \text{CH}$ (4) $\text{CH}_3\text{CH} = \text{CH}_2$
(a) Bromine in carbon tetrachloride (b) Bromine in acetic acid solution
(c) Alk. KMnO_4 (d) Ammonical silver nitrate









74. The principal organic product formed in the reaction given below is :



- (a) $\text{CH}_3 - \text{CHBr}(\text{CH}_2)_8\text{COOH}$ (b) $\text{CH}_2 = \text{CH}(\text{CH}_2)_8\text{COBr}$
(c) $\text{CH}_2\text{BrCH}_2(\text{CH}_2)_8\text{COOH}$ (d) $\text{CH}_2 = \text{CH}(\text{CH}_2)_7\text{CHBrCOOH}$

196


ORGANIC Chemistry for IIT-JEE

75. When 2-butyne is treated with Pd – BaSO₄; the product formed will be :
 (a) *cis*-2-butene (b) *trans*-2-butene (c) 1-butene (d) 2-hydroxy butane
76. In the reaction, $\text{CH}_3\text{C} \equiv \text{C} - \text{CH}_3 \xrightarrow[\text{(ii) Zn/H}_2\text{O}]{\text{(i) X}} \text{CH}_3 - \underset{\text{O}}{\underset{\text{||}}{\text{C}}} - \underset{\text{O}}{\underset{\text{||}}{\text{C}}} - \text{CH}_3$, X is :
 (a) HNO₃ (b) O₂ (c) O₃ (d) KMnO₄
77. Which of the following alkene on catalytic hydrogenation given *cis* and *trans*-isomer ?
 (a)  (b) 
 (c)  (d) all of these
78. In the reaction of hydrogen bromide with an alkene (in the absence of peroxides), the first step of the reaction is the to the alkene.
 (a) fast addition of an electrophile (b) slow addition of an electrophile
 (c) fast addition of a nucleophile (d) slow addition of a nucleophile
79. Which of the following alcohols cannot be prepared from hydration of an alkene ?
 (a)  (b) 
 (c)  (d) 
80. Which of the species shown below is the most stable form of the intermediate in the electrophilic addition of Cl₂ in water to cyclohexene to form a halohydrin ?
 (a)  (b) 
 (c)  (d) 
81. The reaction, $(\text{CH}_3)_2\text{C} = \text{CH}_2 + \text{Br}^\bullet \longrightarrow (\text{CH}_3)_2\dot{\text{C}} - \text{CH}_2\text{Br}$ is an example of a/an step in a radical chain reaction.
 (a) initiation (b) termination
 (c) propagation (d) heterolytic cleavage


HYDROCARBONS (ALKENES)

197


82. Which of the following most accurately describes the first step in the reaction of hydrogen chloride with 1-butene ?





83. Which of the following best describes the flow of electrons in the acid-catalyzed dimerization of $(\text{CH}_3)_2\text{C}=\text{CH}_2$?



84. Hydroboration of 1-methylcyclopentene using B_2D_6 , followed by treatment with alkaline hydrogen peroxide, gives



85.








The correct statements with respect to the above pair of reactions are that

- (I) the reactions are stereospecific
 (II) (X) is erythro and (Y) is threo isomer
 (III) (X) is threo and (Y) is erythro isomer
 (IV) each of (P) and (Q) gives a mixture of (X) and (Y)
- (a) I and II (b) I and III (c) I and IV (d) II and IV

198

ORGANIC Chemistry for IIT-JEE


86. The products *P* and *Q* in the following sequence of reactions, are



87. 4-Pentenoic acid when treated with I_2 and $NaHCO_3$ gives :

- (a) 4, 5-diiodopentanoic acid (b) 5-iodomethyl-dihydrofuran-2-one
(c) 5-iodo-tetrahydropyran-2-one (d) 4-pentenoliodide


88. Product (*B*) of the reaction is:




89. Product (*C*) is :

- (a) $Ph - C \equiv CNa$ (b) $Ph - CH_2 - C \equiv CH$
(c) $Ph - C \equiv C - CH_3$ (d) $Ph - CH = C = CH_2$

90. Which of the following will give a mixture of *cis* and *trans*-1,4-dimethyl cyclohexane, when undergo catalytic hydrogenation ?




91. An optically active compound *A* with molecular formula C_8H_{14} undergoes catalytic hydrogenation to give meso compound, the structure of (*A*) is :




HYDROCARBONS (ALKENES)

199




How many products will be formed in above reaction ?


- (a) 2 (b) 4
(c) 3 (d) 6



- (a) Racemic (b) Diastereomers
(c) Meso (d) Pure enantiomers




- (a) Racemic (b) Diastereomer
(c) Meso (d) E and Z isomer




Rate of reaction towards reduction using (H_2/Pt) :

- (a) $a > b$ (b) $a = b$
(c) $b > a$ (d) Reduction of given molecule is not possible



Product A of the above reaction is :




























- (a) $\text{R} - \overset{\text{O}}{\parallel} \text{C} - \text{R}$ (b) $\text{R}' - \text{CHO}$
(c) $\text{R} - \text{CO}_2\text{H}$ (d) both (a) and (b)



MCPBA $\longrightarrow$ Metachloroperbenzoic acid

200






ORGANIC Chemistry for IIT-JEE



Product (C) in the above reactions is :

HYDROCARBONS (ALKENES)

201









Unknown (A) in the above reaction is :

(a) 2, 2, 3-trimethyl pentane





(b) 2, 2, 4-trimethyl pentane

(c) 2, 2-dimethyl hexane






(d) *n*-octane



Product (C) of the reaction is:




107. What is the major product expected from the following reaction ?



202

ORGANIC Chemistry for IIT-JEE

108. Choose the correct product of this reaction :



- (a)
- (b)
- (c)
- (d)


109. A; Product A is:

- (a)
- (b)
- (c)
- (d)

110. Product is :

- (a)
- (b)
- (c)
- (d)

111. Choose the correct product of the following reactions :



- (a)
- (b)
- (c)
- (d)

112. How many stereoisomeric tetrabromides will be formed in the following reaction ?




- (a) 2 (b) 3 (c) 4 (d) 6

HYDROCARBONS (ALKENES)

203

113. How many stereoisomeric pentabromides will be formed in the following reaction ?



- (a) 2 (b) 3
(c) 4 (d) None of these

114. (Z) (major)

Identify (Z) in the above sequence of reactions :

- (a) (b)
(c) (d)

115. $\text{CH}_3 - \text{CH} - \text{CO}_2\text{K}$
 $\text{CH}_3 - \text{CH} - \text{CO}_2\text{K}$ $\xrightarrow{\text{electrolysis}}$ (A) (Major)

Major product (A) of the above reaction :

- (a) (b)
(c) (d)

116. (A) + (B)

(only one enantiomer is taken)

Which of the following statement is correct about A and B ?






- (a) A and B are mixture of diastereomers
(b) A and B are mixture of enantiomers
(c) A and B are optically active
(d) B is racemic mixture

117. (C) (one of the product)





Identify the product (C):

204

ORGANIC Chemistry for IIT-JEE




Product (Y) of the above reaction is :



119. In the reaction $\text{M} \xrightarrow{\text{C} \equiv \text{C} - \text{Et}} \xrightarrow{\text{NH}_3} \text{J} \xrightarrow{\text{Br}_2/\text{CCl}_4} \text{Q}$; then Q is :



- (a) A pure compound which is optically inactive due to internal compensation
- (b) A binary mixture which is optically inactive due to external compensation
- (c) A binary mixture which is optically active
- (d) A pure compound which is optically inactive due to absence of chiral centre



Which (π -bond) will reduce first, when above compound undergoes catalytic hydrogenation ?


- (a) a
- (b) b
- (c) c
- (d) d

121. Compound A, which is a degradation product of the antibiotic vermiculine has following structure




HYDROCARBONS (ALKENES)

205




(d) None of these



Reagent (A) and (B) in above reaction are

- (a) $A = \text{RCO}_3\text{H}$, $B = \text{H}_2\text{O}_2$ (b) $A = \text{CCl}_4$, $B = \text{HIO}_4$
 (c) $A = \text{RCO}_3\text{H}$, $B = \text{O}_3$ (d) $A = \text{O}_3$, $B = \text{RCO}_3\text{H}$

123. Rank the following in the increasing order of rate of reaction with HBr.




- (a) $R > P > Q$ (b) $R > Q > P$
 (c) $P > R > S$ (d) $P > S > R$

124. Select the reaction(s) that would result in the formation of 2-bromopropane.





- (I) $\text{CH}_3\text{CH}=\text{CH}_2 + \text{HBr} \xrightarrow{\text{peroxide}}$ (II) $\text{CH}_3\text{CH}=\text{CH}_2 + \text{HBr} \xrightarrow{\text{CCl}_4}$
 (III) $\text{CH}_3\text{CH}_2\text{CH}_3 + \text{Br}_2 \xrightarrow{h\nu}$ (IV) $\text{CH}_3\text{CH}=\text{CH}_2 + \text{Br}_2 \xrightarrow{\text{CCl}_4}$
 (a) I and III (b) II and III
 (c) I, II, and III (d) I, II and III

125. Which of the following reactions generates the major product? Ignore stereoisomerism.







206





ORGANIC Chemistry for IIT-JEE



126. In the given selective hydrogenation which combination is incorrect ?




Compound (C) in above sequence of reaction is :







HYDROCARBONS (ALKENES)


207



Major product (A) is :



129. In the reaction given below, the product would be :




(a) a mixture of diastereomers

(b) optically active

(c) optically pure enantiomer

(d) a racemic mixture

130. Surprisingly, the reaction shown below goes through classical carbocation. What is the major product of this reaction?










(a) *trans*-1, 3-dibromocyclohexane

(b) *cis*-1, 3-dibromocyclohexane

(c) *trans*-1, 2-dibromocyclohexane

(d) *cis*-1, 2-dibromocyclohexane

131. The major product of the reaction given below is :



(a) (i) and (ii)

(b) (iii) and (iv)






(c) (v) and (vi)

(d) none of these

208

ORGANIC Chemistry for IIT-JEE


132. Which reaction will occur at the fastest rate ?



Above reaction is known as :


- (a) Wurtz reaction
(c) Fittig reaction

- (b) Wurtz fittig reaction
(d) Kolbe electrolysis




Product A is :


- (a) propane (b) propanol (c) propanoic acid (d) propene

135. Which of the following compound give diastereomers when treated with Br₂ in CCl₄?


Methylcyclopentane



1-Methylcyclopentene




3-Methylcyclopentene



4-Methylcyclopentene

136. A mixture of C₂H₆, C₂H₄ and C₂H₂ is bubbled through alkaline solution of copper (I) chloride, contained in Woulf's bottle. The gas coming out is :

- (a) original mixture (b) C₂H₆
(c) C₂H₆ and C₂H₄ mixture (d) C₂H₄ and C₂H₂



The number of possible products for x and y is :

- (a) 2, 4 (b) 3, 5
(c) 3, 6 (d) 3, 4





HYDROCARBONS (ALKENES)

209

138. Select the incorrect statement :

- (a) Bromine is more selective and less reactive
- (b) Chlorine is less selective and more reactive
- (c) Benzyl free radical is more stable than 2° free radical
- (d) Vinyl free radical more stable than allyl free radical


139. Which of the following compound does not evolve CO₂ gas, when undergo oxidative ozonolysis ?



140. *cis*-3-hexene $\xrightarrow{(a)}$ meso 3,4-hexanediol
trans-3-hexene $\xrightarrow{(b)}$ meso 3,4-hexanediol.

Choose pair of reagent (a, b) for above conversions.

- (a) Cold KMnO₄, OsO₄
- (b) Cold KMnO₄, RCO₃H/H₃O⁺
- (c) RCO₃H/H₃O⁺, cold KMnO₄
- (d) None of these




Product (C) of the above reaction is :

- (a) 1,3-hexadiene
- (b) 1,4-pentadiene
- (c) 1,3-butadiene
- (d) 1,3-heptadiene


142. How many carbon-hydrogen bond orbitals are available for overlap with the vacant *p*-orbital in ethyl carbocation ?

- (a) 0
- (b) 3
- (c) 5
- (d) 6




To achieve above conversion, the reagents used will be :

- (a) O₃/H₂O₂, HO⁻/Δ
- (b) HBr, alc. KOH, O₃, LiAlH₄, H⁺/Δ
- (c) HBr, *t*-BuOK, O₃, KMnO₄, Δ
- (d) HCl, KMnO₄ (cold), H⁺/Δ




210

ORGANIC Chemistry for IIT-JEE




145. Decreasing order of heat evolved upon catalytic hydrogenation of given reactants with a H_2 (Pd/C) is :



- (a) $b > c > a > d$ (b) $d > a > c > b$ (c) $d > c > a > b$ (d) $c > b > c > d$

146.



The correct order of heat of hydrogenation of given molecules is :


- (a) $d > c > a > b$

- (b) $d > a > b > c$


- (c) $b > a > c > d$

- (d) $d > a > b > c$


147.




Product (A) of the above reaction is :



148.



Product (A) is :




HYDROCARBONS (ALKENES)

211




- (a) Racemic mixture
(b) Diastereomers
(c) Meso
(d) Optically inactive due to absence of chiral center



Product (A) of the above reaction is :

- (a) $\text{CH}_3 - \overset{\text{CH}_3}{\underset{\text{HgOAc}}{\underset{|}{\text{C}}}} - \overset{\text{OEt}}{\underset{|}{\text{CH}}} - \text{CH}_3$ (b) $\text{CH}_3 - \overset{\text{CH}_3}{\underset{\text{OEt}}{\underset{|}{\text{C}}}} - \overset{\text{HgOAc}}{\underset{|}{\text{CH}}} - \text{CH}_3$
(c) $\text{CH}_3 - \overset{\text{CH}_3}{\underset{\text{OH}}{\underset{|}{\text{C}}}} - \overset{\text{HgOAc}}{\underset{|}{\text{CH}}} - \text{CH}_3$ (d) $\text{CH}_3 - \overset{\text{CH}_3}{\underset{\text{HgOAc}}{\underset{|}{\text{C}}}} - \overset{\text{OH}}{\underset{|}{\text{CH}}} - \text{CH}_3$




Product (B) of above reaction :

- (a) $\text{Me}_2\text{C}(\text{OH})\text{CH}_2\text{Me}$ (b) $\text{Me}_2\text{CH} - \underset{\text{OH}}{\underset{|}{\text{CH}}} - \text{Me}$
(c) $\text{Me} - \underset{\text{OH}}{\underset{|}{\text{CH}}} - \text{CMe}_3$ (d) $\text{HO} - \text{CH}_2 - (\text{CH}_2)\text{Me}$

152. In which of the following reaction, Markownikoff's rule is violated ?

- (a) $\text{CH}_3 - \text{O} - \text{CH} = \text{CH}_2 \xrightarrow[\text{CCl}_4]{\text{HBr}}$ (b) $\text{CH}_3 - \text{NH} - \text{CH} = \text{CH}_2 \xrightarrow[\text{CCl}_4]{\text{HBr}}$
(c) $\text{CH}_3 - \text{S} - \text{CH} = \text{CH}_2 \xrightarrow[\text{CCl}_4]{\text{HBr}}$ (d) $\text{O}_2\text{N} - \text{CH} = \text{CH}_2 \xrightarrow[\text{CCl}_4]{\text{HBr}}$

153. Decreasing order of rate of reaction of molecules towards electrophilic addition reaction is :



212


ORGANIC Chemistry for IIT-JEE

(a) $a > b > c > d$






(c) $d > b > c > a$

(b) $b > c > a > d$





(d) $b > d > c > a$







Product (A) is :



Product (B) of the above reaction is :



156. Which of the following compounds gives the same carbocation on ionization ?




(a) 1 and 3

(b) 2 and 4

(c) 1 and 2

(d) 1 and 4

157. For the following reactions the major products are shown :

These provide an example of 1 control at low temperature and 2 control at higher temperature.

- 1
(a) kinetic
(c) kinetic

- 2
thermodynamic
kinetic



- 1
(b) thermodynamic
(d) thermodynamic

- 2
kinetic
thermodynamic

HYDROCARBONS (ALKENES)


213

158. What is the product of 1, 4-addition in the reaction shown below ?




Dehydration of the above compound will give :

- (a) meso product (b) racemic mixture
(c) diastereomer (d) optically pure enantiomer



What is stereochemistry of product ?

- (a) Racemic mixture (b) Optically active
(c) Diastereomers (d) Meso product




End product formed in the above reaction is :


- (a) Optically active (b) Racemic (c) Meso (d) Diastereomer

162. How many moles of BH_3 are needed to react completely with 2 mole of 1-pentene in hydroboration-oxidation reaction ?

- (a) 2 mole (b) 3 mole
(c) 2/3 mole (d) 3/2 mole




Product (B) in the above reaction is :







214




ORGANIC Chemistry for IIT-JEE








Product of the above reaction is :







165. In which of the following reactions 1,3-butadiene will be obtained as a major product ?



(d) All of these




Product (A) is :



HYDROCARBONS (ALKENES)

215

168.




Bromination take place at :

- (a) a (b) b (c) c (d) d

169.





Which is incorrect statement about heats of combustion ?




- (c) Iso-butene > *trans*-2-butene > 1-butene (d) *n*-Hexane < *n*-Heptane < *n*-Octane

170.

Predict the major product of the reaction.





171.



- (a) Meso compound (b) Enantiomeric pair
(c) Diastereomers (d) Optically pure enantiomer

172.




216


ORGANIC Chemistry for IIT-JEE





Product (A) of above reaction is:


- (a) $\text{CH}_3\text{O}-\underset{\text{CO}_2\text{H}}{\text{CH}}-\underset{\text{CO}_2\text{H}}{\text{CH}_2}-\text{CH}_2-\text{CHO}$
- (b) $\text{CH}_3\text{O}-\text{CH}_2-\underset{\text{CO}_2\text{H}}{\text{CH}}-\text{CH}_2-\text{CO}_2\text{H}$
- (c) $\text{CH}_3\text{O}-\underset{\text{CO}_2\text{H}}{\text{CH}}-\text{CH}_2-\text{CH}_2-\text{CO}_2\text{H}$
- (d) $\text{CH}_3\text{O}-\underset{\text{CO}_2\text{H}}{\text{CH}}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CHO}$







- (a) Diastereomers (b) Racemic mixture (c) Meso (d) Optically pure enantiomer



- (a)  (b) 
- (c)  (d) 




In polyenes that contain differently substituted ($\text{C}=\text{C}$) double bonds, it is possible to hydrogenate chemoselectively one ($\text{C}=\text{C}$) double bond. Product is :

- (a)  (b) 
- (c)  (d) 

HYDROCARBONS (ALKENES)


217





(MCPBA $\rightarrow$ meta-chloro perbenzoic acid)

Stereochemistry of the product of above reaction is :



- (a) Meso (b) Racemic
(c) Diastereomers
(d) Optically inactive due to absence of chiral center.



Identify product (P).






A isomerise to B on addition of traces of acid H_2SO_4 . Compound (B) is :



Product (A) of the reaction is :







218

ORGANIC Chemistry for IIT-JEE










(d) None of these

180.



181.

Which of the following reactions do not represent the major product of given Birch reductions ?




(a) (i), (iii), (vi)

(b) (iv), (vi), (vii)

(c) (iv), (v), (vi)

(d) (i), (ii), (v), (vii)


182.




Product (A) is:

HYDROCARBONS (ALKENES)

219








Hint : Think carefully about the relative stabilization of developing positive charge, when the double bond reacts with an electrophile.




Correct statement about above reaction is:

(a) A = <i>cis</i> -2-chlorocyclohexanol,	B = cyclohexeneoxide
(b) A = <i>trans</i> -2-chloro cyclohexanol,	B = anti-diol
(c) A = <i>trans</i> -2-chlorocyclohexanol,	B = cyclohexeneoxide
(d) A = <i>cis</i> -2-chlorocyclohexanol,	B = anti-diol



220


ORGANIC Chemistry for IIT-JEE





187.  $\xrightarrow[\text{CCl}_4]{\text{Br}_2}$ stereochemistry of the product is:

- (a) Diastereomers (b) Racemic mixture
(c) Meso (d) Pure Enantiomers

188.  $\xrightarrow[\text{CCl}_4]{\text{Br}_2}$ Product/s obtained is/are :

- (a) Diastereomers (b) Racemic
(c) Meso (d) Optically pure enantiomers

189.  $\xrightarrow{\text{Ph}_3\text{P}=\text{CH}_2}$ $\xrightarrow{\text{Ph}_3\text{P}=\text{CH}_2}$ (x) ; Product (x) is :


- (a)  (b) 
(c)  (d) 

190. $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \underset{\text{CH}_3}{\overset{\text{CH}_3}{\text{C}}} - \text{CH}_2 - \text{OH} \xrightarrow[\Delta]{\text{H}^+} (\text{A})$; Product (A) is :
(major)

- (a) $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \underset{\text{CH}_3}{\text{C}} = \text{CH} - \text{CH}_3$
(b) $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{CH} = \text{C} \begin{matrix} \text{CH}_3 \\ \text{CH}_3 \end{matrix}$
(c) $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \underset{\text{CH}_3}{\text{C}} = \text{CH}_2$
(d) $\text{CH}_3 - \text{CH}_2 - \underset{\text{CH}_3}{\text{CH}} - \text{CH}_2 - \underset{\text{CH}_3}{\text{C}} = \text{CH}_2$




HYDROCARBONS (ALKENES)

221




Comment on optical activity of the products:

- (a) Racemic (b) Diastereomer
(c) Meso (d) Optically pure enantiomer



Alkene (A) will be :





- (a) *cis*-2-pentene (b) *cis*-3-hexene
(c) *cis*-4-octene (d) *trans*-2-hexene



Product (A) is

- (a) *trans*-2-butane (b) *cis*-2-butene (c) 1-butene (d) Iso-butene


195. In which of the following reactions, two products will be formed other than phosphonium ylide (POPh_3)



222


ORGANIC Chemistry for IIT-JEE





196. To carry out the given conversions, select the correct option:




- (a) $a = \text{Ag}_2\text{O}$, $b = \text{Zn}/\text{CH}_3\text{CO}_2\text{H}$, $c = \text{LiAlH}_4$
 (b) $a = \text{H}_2\text{O}_2$, $b = \text{CH}_3 - \text{S} - \text{CH}_3$, $c = \text{NaBH}_4$
 (c) Both (a) and (b)
 (d) None of these

197. The product (A) of given alkoxymercuration de-mercuration is :




- (a)  (b)  (c)  (d) 



End product of the reaction is :

- (a) $\text{H}_2\text{C} = \text{CH} - \underset{\text{CH}_3}{\text{C}} = \text{CH}_2$ (b) $\text{CH}_3 - \text{CH} = \text{CH} - \text{CH} = \text{CH}_2$
 (c) $\text{H}_2\text{C} = \text{CH} - \text{CH} = \text{CH}_2$ (d) $\text{H}_2\text{C} = \text{CH} - \text{CH}_2 - \text{CH} = \text{CH}_2$

199. Major product of the given reaction is :








- (a) $\text{CH}_3 - \underset{\text{I}}{\text{CH}} - \underset{\text{I}}{\text{CH}_2}$ (b) $\text{CH}_3 - \underset{\text{I}}{\text{CH}} - \text{CH}_3$
 (c) $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{I}$ (d) $\text{I} - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{I}$

HYDROCARBONS (ALKENES)**223**

200. The rate constant for a reaction can be increased by a the stability of the reactant or by b the stability of the transition state. Select the correct choice for *a* and *b*.





- (a) decreasing, decreasing (b) increasing, decreasing
(c) decreasing, increasing (d) increasing, increasing

201. Major product of the given reaction is :








202. + $\text{Ph}_3\text{P}=\text{CH}_2 \longrightarrow$ (A) Major

Major product (A) is :







203. In the given reaction, only one alkene undergo preferential oxidation by electrophilic ozone. Identify product (P) of the given reaction:






224

ORGANIC Chemistry for IIT-JEE




Product (B) is :





HYDROCARBONS (ALKENES)

225


208. An unknown alkene (A) reacts with 3 mole of H_2 gas in presence of platinum catalyst to form 1-isopropyl-4-methyl cyclohexane. When unknown alkene (A) is ozonized and reduced, following product are obtained




The alkene (A) is :



Product (C) is






210. The following reaction take place in high yields.




Use your knowledge of alkene chemistry to predict a product even though you have never seen this reaction before







What is the ratio of glyoxal to pyruvaldehyde obtained in the above reaction ?

- (a) 1 : 3
(c) 3 : 2







Which of the following product cannot be obtained in above reaction ?



(TMAO $\rightarrow$ trimethyl amine -N - oxide)


Product (A) is :






HYDROCARBONS (ALKENES)

227

214.




Product (A) of the reaction is :







(d) None of these




215.



Product (A) is :



216.







Arrange the above in the decreasing order of reactivity towards HBr :

(a) $a > b > c$ (b) $b > a > c$ (c) $b > c > a$ (d) $a > c > b$





228

ORGANIC Chemistry for IIT-JEE

217. Which reaction has the lowest $\Delta G^\ddagger$ or (Activation-Energy)?



218. Which of the following will rearrange ?







(a) 1

(b) 1 and 3


(c) All

(d) 1, 2, 4,

219. Which of the following is most likely to undergo a favorable hydride shift ?



220. Energy profile diagram for dehydration of 2-butanol using conc. H_2SO_4 is given below :



Product (b) of above reaction is :

(a) 1-butene

(b) *cis*-2-butene

(c) *trans*-2-butene




(d) *iso*-butene

HYDROCARBONS (ALKENES)**229**


221. How many alkene on catalytic hydrogenation given isopentane as a product ?

- (a) 2 (b) 3
(c) 4 (d) 5

222. Which of the following would not rearrange to a more stable form ?




223. Consider the following reaction.








In this reaction SbF_5 acts as:

- (a) an acid (b) a base
(c) a nucleophile (d) an electrophile

224.  $\xrightarrow{\text{Br}_2/h\nu}$ Major (X) $\xrightarrow[\Delta]{\text{conc. alic. KOH}}$ Major (Y) $\xrightarrow{\text{H-Br, peroxide}}$ Major (Z):

Product (Z) is:









Relation between (B) and (C) is:

- (a) Enantiomer (b) Diastereomer
(c) Geometrical isomer (d) Meso

230

ORGANIC Chemistry for IIT-JEE

226. The reaction of HBr with the following compound would produce :



- (a) Nucleophilic addition
(c) Electrophilic addition
(e) Free radical substitution

- (b) Nucleophilic substitution
(d) Electrophilic substitution

ATDB.uno

HYDROCARBONS (ALKENES)

64 50 58 5 62

231**ANSWERS — LEVEL 1**

1.	(c)	2.	(d)	3.	(c)	4.	(d)	5.	(b)	6.	(b)	7.	(c)	8.	(c)
9.	(c)	10.	(d)	11.	(b)	12.	(d)	13.	(c)	14.	(b)	15.	(b)	16.	(c)
17.	(d)	18.	(d)	19.	(b)	20.	(d)	21.	(b)	22.	(b)	23.	(b)	24.	(b)
25.	(b)	26.	(b)	27.	(d)	28.	(b)	29.	(d)	30.	(b)	31.	(c)	32.	(b)
33.	(a)	34.	(b)	35.	(b)	36.	(b)	37.	(b)	38.	(b)	39.	(b)	40.	(b)
41.	(d)	42.	(e)	43.	(c)	44.	(c)	45.	(a)	46.	(c)	47.	(c)	48.	(b)
49.	(b)	50.	(b)	51.	(b)	52.	(a)	53.	(b)	54.	(d)	55.	(b)	56.	(c)
57.	(c)	58.	(b)	59.	(c)	60.	(a)	61.	(b)	62.	(d)	63.	(a)	64.	(b)
65.	(d)	66.	(b)	67.	(d)	68.	(a)	69.	(c)	70.	(d)	71.	(d)	72.	(c)
73.	(d)	74.	(c)	75.	(a)	76.	(c)	77.	(d)	78.	(b)	79.	(d)	80.	(d)
81.	(c)	82.	(b)	83.	(a)	84.	(a)	85.	(a)	86.	(d)	87.	(b)	88.	(b)
89.	(c)	90.	(d)	91.	(b)	92.	(b)	93.	(a)	94.	(a)	95.	(a)	96.	(d)
97.	(b)	98.	(a)	99.	(c)	100.	(d)	101.	(b)	102.	(b)	103.	(d)	104.	(b)
105.	(b)	106.	(c)	107.	(d)	108.	(b)	109.	(d)	110.	(d)	111.	(c)	112.	(b)
113.	(a)	114.	(b)	115.	(c)	116.	(a)	117.	(d)	118.	(d)	119.	(b)	120.	(d)
121.	(b)	122.	(c)	123.	(a)	124.	(b)	125.	(d)	126.	(a)	127.	(b)	128.	(c)
129.	(d)	130.	(a)	131.	(c)	132.	(d)	133.	(c)	134.	(a)	135.	(d)	136.	(c)
137.	(b)	138.	(d)	139.	(d)	140.	(b)	141.	(b)	142.	(b)	143.	(b)	144.	(b)
145.	(b)	146.	(c)	147.	(b)	148.	(a)	149.	(d)	150.	(b)	151.	(a)	152.	(d)
153.	(c)	154.	(c)	155.	(b)	156.	(c)	157.	(a)	158.	(a)	159.	(b)	160.	(c)
161.	(b)	162.	(c)	163.	(b)	164.	(d)	165.	(d)	166.	(b)	167.	(b)	168.	(a)
169.	(c)	170.	(b)	171.	(b)	172.	(d)	173.	(b)	174.	(c)	175.	(b)	176.	(b)
177.	(b)	178.	(c)	179.	(b)	180.	(c)	181.	(b)	182.	(b)	183.	(c)	184.	(c)
185.	(c)	186.	(b)	187.	(a)	188.	(b)	189.	(b)	190.	(b)	191.	(a)	192.	(b)
193.	(c)	194.	(b)	195.	(c)	196.	(c)	197.	(b)	198.	(a)	199.	(b)	200.	(c)
201.	(c)	202.	(c)	203.	(b)	204.	(d)	205.	(b)	206.	(c)	207.	(b)	208.	(b)
209.	(b)	210.	(b)	211.	(c)	212.	(c)	213.	(b)	214.	(a)	215.	(b)	216.	(b)
217.	(d)	218.	(c)	219.	(a)	220.	(b)	221.	(b)	222.	(c)	223.	(d)	224.	(c)
225.	(b,c)	226.	(b)	227.	(c)										

232

ORGANIC Chemistry for IIT-JEE



1.

Reagents

A. HCl	B. Br ₂	C. Hg(OAc) ₂ in H ₂ O	D. B ₂ H ₆ (BH ₃) in ether
E. H ₂ O ₂	F. KMnO ₄ in H ₂ O	G. HOBr	H. NaBH ₄

In each reagent box write a letter designating the best reagent and condition selected from the above list of reagents.

Reactant	Reagent	Product
$(\text{CH}_3)_2\text{CHCH}=\text{CH}_2$ 3-methyl-1-butene	(i) ☐	$(\text{CH}_3)_2\text{CHCH}(\text{Cl})\text{CH}_3$ 2-Chloro-3-methyl butane
	(ii) ☐	$(\text{CH}_3)_2\text{CHCHBrCH}_2\text{Br}$ 1, 2-dibromo-3-methyl butane
	(iii) ☐	$(\text{CH}_3)_2\text{CHCHOHCH}_2\text{Br}$ bromo-3-methyl 2-butanol
	(iv) ☐	$(\text{CH}_3)_2\text{CHCH}(\text{OH})\text{CH}_3$ 3-methyl-2-butanol
	(v) ☐	$(\text{CH}_3)_2\text{CHCH}(\text{OH})\text{CH}_2\text{OH}$ 3-methyl-1, 2-butanediol

2. **Propene** ($\text{CH}_3 - \text{CH} = \text{CH}_2$) can be transformed to compounds (a to j) listed in the left-hand column.

Write letter designating the reagent, you believe will achieve desired transformation. In the case of a multi step sequence write the reagent in the order they are to be used.

	Desired Product	No. of Steps	Write options	Reagent List
a.	$\text{CH}_3\text{CHBrCH}_2\text{Br}$	one	A.	Hg(OAc) ₂ in H ₂ O
b.	$(\text{CH}_3)_2\text{CHOH}$	two	B.	B ₂ H ₆ in THF

HYDROCARBONS (ALKENES)**233**










c.	$\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$	two	C.	NaBH_4 in alcohol
d.	CH_3COCH_3	three	D.	Br_2 in CH_2Cl_2
e.	$\text{CH}_3\text{CH}_2\text{CHO}$	three	E.	H_2O_2 in aqueous base
f.	$\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{Br}$	one	F.	HOBr (NBS in aqueous acetone)
g.	$(\text{CH}_3)_2\text{CHBr}$	one	G.	HBr in CH_2Cl_2
h.	$\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{OH}$	two	H.	OsO_4 in ether
i.	$\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{Cl}$	three	I.	Thionyl chloride (SOCl_2)
j.	$\text{CH}_3 - \text{C} \equiv \text{CH}$	two	J.	NH_4Cl in aqueous acetone
			K.	NaOH in alcohol and reflux
			L.	NaNH_2 (strong base)

234

ORGANIC Chemistry for IIT-JEE

3. In each reaction box write a single letter designating the best reagent and condition selected from the list at bottom of the page.

(F.S., → first step, S.S. → second step, T.S. → third step)


Reaction	Reactant	Options	Product
1.		F.S. ☐ S.S. ☐ →	
2.		F.S. ☐ S.S. ☐ T.S. ☐ →	
3.		F.S. ☐ S.S. ☐ →	
4.		F.S. ☐ S.S. ☐ →	
5.		F.S. ☐ S.S. ☐ T.S. ☐ →	

A. $\text{NaBH}_4/\text{alcohol}$	B. $\text{Ph}-\text{CO}_3\text{H}/\text{CH}_2\text{Cl}_2$	C. PCC	D. $\text{CH}_3\text{ONa}/\text{CH}_3\text{OH}$
E. B_2H_6 in THF	F. $\text{H}_2\text{O}_2/\text{aq. NaOH}$	G. H_3PO_4 & heat	H. $\text{AlCl}_3/\text{C}_6\text{H}_6$
I. O_3 in CH_2Cl_2	J. Br_2 in CH_2Cl_2	K. 20% KOH & heat	L. $\text{Ph}-\text{Li}/\text{ether}$

HYDROCARBONS (ALKENES)

235

4. Match the reagents a-j with products A-J. There is one best product for each reaction.



The molecule (x) is the starting material for all reactions in problem. Do the ones you know first and then tackle the rest by deductive reasoning

Products		Reagents		Option	
A		B		(a) H_2O heat, pH 7	
C		D		(b)	
E		F		(c) tBuOK , polar aprotic solvent	
G		H		(d) (1) OsO_4 , CH_2Cl_2 (2) H_2O , NaOH , H_2O_2	
I		J		(e) Br_2 , CCl_4	
K		L		(f) NBS , $h\nu$, CCl_4	
M		N		(g) (1) H_3O^+ (2) NaOH , H_2O	
O		P		(h) (1) BH_3 , ether (2) H_2O_2	
Q		R		(i) (1) OsO_4 (2) NaOH , H_2O	
S		T		(j) $\text{H}_2/\text{Pd/C}(\text{EtOH})$	

236

ORGANIC Chemistry for IIT-JEE

5. Match the column:

Column (I)		Column (II)	
(a)	$\text{CH}_3 - \text{C} \equiv \text{C} - \text{CH}_3$	(p)	cis-product with $\text{H}_2 / \text{Pd} - \text{BaSO}_4$
(b)	$\text{CH}_3 - \text{CH}_2 - \text{C} \equiv \text{CH}$	(q)	Trans-product with $\text{Na} / \text{liq. NH}_3$
(c)	$\text{CH}_3 - \text{C} \equiv \text{CH}$	(r)	White with amm. AgNO_3
(d)	$\text{CH}_3 - \text{C} \equiv \text{C} - \text{Et}$	(s)	H_2 gas with Na

ATDB.uno

HYDROCARBONS (ALKENES)

237









6. Match the column I with column II and with column III (Matrix).





Column-I		Column- II		Column- III	
Reaction		Nature of product formed		Number of chiral center present in product. (Consider only one isomer in case of racemic mixture or Diastereomer)	
(a)	$\xrightarrow[\text{CCl}_4]{\text{Br}_2}$	(p)	Racemic mixture	(w)	0
(b)	$\xrightarrow[\text{CCl}_4]{\text{Br}_2}$	(q)	Meso	(x)	1
(c)	$\xrightarrow[\text{CCl}_4]{\text{Br}_2}$	(r)	Diastereomer	(y)	2
(d)	$\xrightarrow[\text{CCl}_4]{\text{Br}_2}$	(s)	Vicinal dihalide	(z)	3

238



ORGANIC Chemistry for IIT-JEE

7. Match the column I and II.

Column (I)		Column (II)	
	Reaction		Product
(a)		(p)	
(b)		(q)	
(c)		(r)	
(d)		(s)	



Sum of molecular mass of A, B, C, D (i.e. A + B + C + D) is equal to :

HYDROCARBONS (ALKENES)**239**

Total number of products A and B (i.e. A + B) is equal to :

10.


Reaction 1	Reaction 2
$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H} - \text{C} - \text{Br} \\ \\ \text{CH} \\ \\ \text{CH} \\ \\ \text{H} - \text{C} - \text{Br} \\ \\ \text{CH}_3 \end{array} \xrightarrow[\text{CCl}_4]{(\text{cis}) \text{Br}_2} (\text{P}) $	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H} - \text{C} - \text{Br} \\ \\ \text{CH} \\ \\ \text{CH} \\ \\ \text{H} - \text{C} - \text{Br} \\ \\ \text{CH}_3 \end{array} \xrightarrow[\text{CCl}_4]{(\text{trans}) \text{Br}_2} (\text{Q}) $
Reaction 3	Reaction 4
$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H} - \text{C} - \text{Br} \\ \\ \text{CH} \\ \\ \text{CH} \\ \\ \text{CH}_3 \end{array} \xrightarrow[\text{CCl}_4]{(\text{cis}) \text{Br}_2} (\text{R}) $	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H} - \text{C} - \text{Br} \\ \\ \text{CH} \\ \\ \text{CH} \\ \\ \text{CH}_3 \end{array} \xrightarrow[\text{CCl}_4]{(\text{trans}) \text{Br}_2} (\text{S}) $
Sum of products P, Q, R, S (i.e. P + Q + R + S) is equal to :	

11. Comprehension


Vladimir Markovnikov rule :

Alkenes undergo electrophilic addition reactions. It is triggered by the acid acting as a electrophile toward π -electrons of the double bond.





Markovnikov's rule states that when an unsymmetrically substituted alkene reacts with a hydrogen halide, the hydrogen atom adds to the carbon that has the greater number of hydrogen, e.g.,







Mechanism :






A. Which of the following is most reactive toward Markovnikov addition ?



B. What is the energy profile for the given reaction ?




C. In which of following reactions carbocation rearrangement is possible ?




(d) All of these

HYDROCARBONS (ALKENES)**241**


D. Identify the major products r_1 , r_2 and r_3 in the given reactions.



E. In which of the following reactions, product is racemic mixture ?




F. In which of the following reactions, diastereomers will be formed ?



242




ORGANIC Chemistry for IIT-JEE

12. Comprehension



Consider the above reaction and answer A to E.



A. What is electrophile in first step ?






B. What is nucleophile in first step ?






(b) 1-butene



C. What is electrophile in second step ?



D. What is nucleophile in second step ?







E. Which step is rate determining step ?

(a) attack of nucleophile CH_3OH (b) attack of electrophile H^+ (c) attack of nucleophile H_2O (d) attack of electrophile CH_3^+

HYDROCARBONS (ALKENES)

243

13. Match the column I and II :

Column (I)		Column (II)	
Conversion		Reagent	
(a)		(p)	$\text{SO}_2\text{Cl}_2 / h\nu$ (2 equivalent)
(b)		(q)	NBS (2 equivalent)
(c)		(r)	NBS then $\text{SO}_2\text{Cl}_2/h\nu$
(d)		(s)	$\text{SO}_2 \text{Cl}_2 / h\nu$ then NBS


244

ORGANIC Chemistry for IIT-JEE










ANSWERS — LEVEL 2

1. (i) – A; (ii) – B; (iii) – G; (iv) – C; (v) – F
2. a – D; b – A, C; c – B, E; d – A, C, F; e – B, E, F; f – F; g – G; h – I, K; i – B, E, I; j – D, L
3. Reaction 1 : B, D; Reaction 2 : E, F, C Reaction 3 : I, A
Reaction 4 : L, G Reaction 5 : B, L, C
4. a – C; b – D; c – A; d – F; e – I; f – J; g – E; h – H; i – B; j – G
5. a – p, q; b – r, s; c – r, s; d – p, q
6. a – r, s – z; b – p, s – y; c – p, s – y; d – q, s – y
7. a – r; b – p; c – q; d – s
8. $A + B + C + D = 312$
9. $A + B = 5$
10. $P + Q + R + S = 8$
11. A – b; B – c; C – d; D – b; E – d; F – d
12. A – b; B – b; C – c; D – b; E – b
13. a – q; b – p; c – s; d – r



ATDB.uno






4c HYDROCARBONS (ALKYNES)








Product (K) of the above reaction is :











Give the structural formula of compound (B) :



; This conversion can be achieved by :




6. Which alkyne will give 3-ethylhexane on catalytic hydrogenation ?






(d) All of these

7. Reactant P gives products Q or R.



The possible reagents are :



The correct statement with respect to the above conversion is/are :

(a) Q is obtained on treatment with reagent (I)












(b) R and Q are obtained on treatment with reagent (II)

(c) R is obtained on treatment with reagent (I)




HYDROCARBONS (ALKYNES)

247

(d) R is obtained on treatment with reagent (II)




11. Which combination is best for preparation of the compound (A) shown below ?



12. Which one of the following is the intermediate in the preparation of a ketone by hydration of an alkyne in the presence of sulfuric acid and mercury (II) sulphate ?


248

ORGANIC Chemistry for IIT-JEE








To carry out above conversion, (A) and (B) respectively, are :

- (a) $\text{NaNH}_2, \text{Cl}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{Br}$
 (b) $\text{NaNH}_2, \text{F}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{Br}$
 (c) $\text{NaNH}_2, \text{I}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{Br}$
 (d) $\text{NaNH}_2, \text{I}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{I}$




- (a) $\text{Ph}-\text{C}(\text{I})=\text{CH}-\text{I}$ (b) $\text{Ph}-\text{CH}(\text{I})-\text{CH}_2-\text{I}$
 (c) $\text{Ph}-\text{C}\equiv\text{C}-\text{I}$ (d) $\text{I}-\text{C}\equiv\text{C}-\text{H}$



- (a)  (b)  (c)  (d) 


16. Which of the following alkyne on treatment with H_2 (2 mole)/ Pt gives an optically inactive compound ?

- (a) 3-Methyl-1-pentyne (b) 4-Methyl-1-hexyne
 (c) 3-Methyl-1-heptyne (d) None of the above




- (a) Toluene (b) Ethyl-benzene (c) Benzene (d) Butyne

18. What is the final product, C, of the following reaction sequence ?







HYDROCARBONS (ALKYNES)

249



Compound (X) will be :



20. Choose the sequence of steps that describes the best synthesis of 1-butene from ethanol :


(a) (1) $\text{NaC} \equiv \text{CH}$; (2) H_2 , Lindlar Pd(b) (1) $\text{NaC} \equiv \text{CH}$; (2) Na , NH_3 (c) (1) HBr , heat ; (2) $\text{NaC} \equiv \text{CH}$; (3) H_2 , Lindlar Pd(d) (1) HBr , heat ; (2) $\text{KOC}(\text{CH}_3)_3$, DMSO ; (3) $\text{NaC} \equiv \text{CH}$; (4) H_2 , Lindlar catalyst21. Which alkyne yields butanoic acid ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$) as the only organic product on treatment with ozonolysis followed by hydrolysis?

(a) 1-Butyne

(b) 2-Pentyne

(c) 1-Pentyne

(d) 2-Hexyne



Carlina oxide


Unit of unsaturation in compound (A) ?

(a) 7

(b) 8

(c) 9

(d) 10



Product (C) of above reaction is:

(a) $\text{H}_2\text{C} = \text{CH}_2$ (b) $\text{CH}_3 - \text{C} \equiv \text{C} - \text{CH}_3$ (c) $\text{HC} \equiv \text{CH}$ (d) $\text{CH}_3 - \text{CH} = \text{CH} - \text{CH}_3$

24. To convert 1-butyne to 1-D-butanal, one would carry out the following steps :

(I) Sodium amide, then D_2O

(II) Disiamylborane, then hydrogen peroxide/sodium hydroxide

(III) The transformation can not be carried out with the indicated reagents.







(a) I, followed by II (b) II, followed by I (c) III

(d) II





250

ORGANIC Chemistry for IIT-JEE

25. An unknown compound (A) has a molecular formula C_4H_6 . When (A) is treated with excess of Br_2 a new substance (B) with formula $C_4H_6Br_4$ is formed. (A) forms a white ppt. with ammonical silver nitrate solution. (A) may be :
- (a) But-1-yne (b) But-2-yne
(c) But-1-ene (d) But-2-ene
26. One mole of 1,2-dibromopropane on treatment with X moles of $NaNH_2$ followed by treatment with ethyl bromide gave a pentyne. The value of X is :
- (a) One (b) Two (c) Three (d) Four




Product (A) is:



ANSWERS — LEVEL 1

1.	(b)	2.	(b)	3.	(b)	4.	(b)	5.	(b)	6.	(d)	7.	(c)	8.	(c)
9.	(b)	10.	(d)	11.	(b)	12.	(d)	13.	(c)	14.	(c)	15.	(c)	16.	(a)
17.	(c)	18.	(a)	19.	(a)	20.	(c)	21.	(b)	22.	(c)	23.	(c)	24.	(c)
25.	(a)	26.	(c)	27.	(c)	28.	(b)								









5A

ALKYL HALIDES

Substitution Reactions (S_{N1} , S_{N2} , S_{N1}')




1. Which of the following is not expected to be intermediate of the following reaction ?




252


ORGANIC Chemistry for IIT-JEE




3. Rate of S_N2 will be negligible in :



4. What is the major product obtained in the following reaction ?




5. $\text{Cl}-\text{CH}_2-\text{C}(\text{CH}_3)_2-\text{CH}_2-\text{CH}_2-\text{Cl} + \text{I} \xrightarrow{\text{DMF}}$ product; Major product of this reaction is:



6. Which of the following expressions is representative of the rate law for a S_N2 reaction ?


- (a) Rate = k [electrophile] (b) Rate = k [electrophile] [nucleophile]
 (c) Rate = k [nucleophile]² (d) Rate = k [electrophile]²

7. ; Major product of this reaction is :




ALKYL HALIDES (SUBSTITUTION)**253**


8. Which of the following alkyl halide undergo rearrangement in S_N1 reaction ?




9. Arrange the following three chlorides in decreasing order towards S_N1 reactivity.



10. Which compound undergoes nucleophilic substitution with NaCN at the fastest rate ?




11. Rank the following in order of decreasing rate of solvolysis with aqueous ethanol (fastest $\rightarrow$ slowest)




12. The reaction of 4-bromobenzyl chloride with sodium cyanide in ethanol leads to the formation of :



13. Which of the following reactant will not favour nucleophilic substitution reaction ?



14.



254









ORGANIC Chemistry for IIT-JEE

Conversion of I to II :

(a) takes place by S_N1 (b) takes place by S_N2 (c) takes place both by S_N1 and S_N2


(d) does not take place

15. Which is the correct reaction coordinate diagram for the following solvolysis reaction ?






(c) both (a) and (b)

(d) None of these




ALKYL HALIDES (SUBSTITUTION)


255






(d) None of these

18. Which of the following represents the correct graph for S_N2 reaction ?






19. Which of the following graph represents correct graph for S_N1 reaction :




ALKYL HALIDES (SUBSTITUTION)




257







20. Which of the following is most reactive toward S_N² reaction ?



21. Among the given pairs, in which pair first compound reacts faster than second compound in S_N¹ reaction ?







22. What is the major product of the following reaction ?




258

ORGANIC Chemistry for IIT-JEE



23. S_N1 and S_N2 products are same with (excluding stereoisomer) :





24. Consider the nucleophilic attacks given below. Select in each pair that shows the greater S_N2 reaction rate.





or




or



or





or



- A B C D
(a) (I) ; (IV) ; (VI) ; (VIII)
(c) (I) ; (III) ; (V) ; (VIII)

- A B C D
(b) (II) ; (III) ; (V) ; (VIII)
(d) (I) ; (III) ; (V) ; (VII)

25. Which of the two stereoisomers of 4*t*-butylcyclohexyl iodide ($^{127}\text{I}^-$) will undergo S_N2 substitution with $^{128}\text{I}^-$ faster, and why ?






- (a) A will react faster because it is the more stable of the two isomers
(b) A will react faster because it will yield a more stable product, and the transition state for both reactions is of the same energy
(c) A will react faster because the approach of $^{128}\text{I}^-$ can depart unhindered.
(d) B will react faster because it is less stable than A, and the transition state for both reactions is of the same energy

ALKYL HALIDES (SUBSTITUTION)

259

26. (Z)-2-Butene reacts with $\text{Br}_2/\text{H}_2\text{O}$. The resulting bromohydrin when treated with methoxide in methanol undergoes an intramolecular $\text{S}_{\text{N}}2$ reaction. Taking into consideration the stereochemical consequences of the reaction mechanism involved, choose the final product(s) of these transformations.






(a) (I) only

(b) (II) only







(c) (III) only

(d) Equal amounts of (I) and (II)

27. Rank the following species in order of decreasing nucleophilicity in a polar protic solvent (most $\rightarrow$ least nucleophilic):

(a) $3 > 1 > 2$ (b) $2 > 3 > 1$ (c) $1 > 3 > 2$ (d) $2 > 1 > 3$




28. Identify products of the given reactions:






262

ORGANIC Chemistry for IIT-JEE





















38. In the given pairs of alkyl-halide, in which pair the first compound is more reactive than second compound toward S_N2 reaction?



39. In the given pair of reaction in which pair the second reaction is more reactive than first toward S_N2 reaction?




40. Among the following pairs of reactions, which pair the second reaction is more reactive than first in S_N2 reaction?



ALKYL HALIDES (SUBSTITUTION)**263**

44. Increasing the concentration of an electrophile in a typical S_N2 reaction by a factor of 3 and the concentration of the nucleophile by a factor of 3 will change the reaction rate to :
- (a) increase by a factor of 6 (b) increase by a factor of 9
(c) decrease by a factor of 3 (d) remain about the same
45. Consider the following reaction and select the best choice that represents the reaction.




- (a)
- (b)
- (c)
- (d)

46. Product; Identify the product.





- (a)
- (b)
- (c)
- (d)

47. The reaction ,



proceeds by the..... mechanism.

- (a) S_Ni (b) S_N2 (c) S_E2 (d) S_{N1}
48. Consider the following anions.








When attached to sp^3 -hybridized carbon, their leaving group ability in nucleophilic substitution reaction decreases in the order :

- (a) $\text{I} > \text{II} > \text{III} > \text{IV}$ (b) $\text{I} > \text{II} > \text{IV} > \text{III}$ (c) $\text{IV} > \text{I} > \text{II} > \text{III}$ (d) $\text{IV} > \text{III} > \text{II} > \text{I}$

264

ORGANIC Chemistry for IIT-JEE





50. Reaction of *R*-2-butanol with *p*-toluenesulphonyl chloride in pyridine followed by reaction with LiBr gives:

- (a) *R*-2-butyl bromide
(c) *R*-2-butyl tosylate

- (b) *S*-2-butyl tosylate
(d) *S*-2-butyl bromide

51. The compound which undergoes S_{N}^1 reaction most easily is.



52. Addition of KI accelerates the hydrolysis of primary alkyl halides because :






- (a) KI is soluble in organic solvents
(b) the iodide ion is a weak base and a poor leaving group
(c) the iodide ion is a strong base
(d) the iodide ion is a powerful nucleophile as well as a good leaving group

53. Which of the following phrases are not correctly associated with S_{N}^1 reaction ?


- (1) Rearrangement is possible
(2) Rate is affected by polarity of solvent
(3) The strength of the nucleophile is important in determining rate
(4) The reactivity series is tertiary > secondary > primary
(5) Proceeds with complete inversion of configuration
(a) 3, 5
(b) 5 only
(c) 2, 3, 5
(d) 3 only

ALKYL HALIDES (SUBSTITUTION)

265



55. Energy profile diagram for an exothermic reaction, $A \xrightarrow{1} B \xrightarrow{2} C \xrightarrow{3} D$, is given below.









The rate determining step of the reaction is :

(a) $A \longrightarrow B$

(b) $B \longrightarrow C$

(c) $C \longrightarrow D$





(d) can not predict




The product X is :

266

ORGANIC Chemistry for IIT-JEE



58. Relative rate of reaction of the following amine with methyl iodide is:




(A)

(B)





(C)

(a) $A > B > C$ (b) $A > C > B$ (c) $B > C > A$ (d) $B > A > C$


59.







Major product which you expect in the above reaction is :



60.








Major product of the above reaction is :





ALKYL HALIDES (SUBSTITUTION)




267



- (a)  (b) 
 (c)  (d) 

62. The decreasing order of reactivity of the compounds given below towards solvolysis under identical conditions is:





- (a)  (b)  (c)  (d) None of these

64. (R)-2-octyl tosylate is solvolyzed in water under ideal S_N1 conditions. The product(s) will be:

- (a) R-2-octanol and S-2-octanol in a 1 : 1 ratio
 (b) R-2-octanol and S-2-octanol in a 1.5 : 1 ratio
 (c) R-2-octanol only
 (d) S-2-octanol only

270

ORGANIC Chemistry for IIT-JEE

75. Relative rate of reaction with H_2O .


(a) (i) > (ii) > (iii)

(b) (ii) > (i) > (iii)



(c) (iii) > (ii) > (i)

(d) (iii) > (i) > (ii)

76.




End product (P) of the above reaction is :

77. Which of the following statements is correct regarding the rate of hydrolysis of the compounds (A) and (B) by $\text{S}_{\text{N}}1$ reaction ?

ALKYL HALIDES (SUBSTITUTION)**271**


- (a) A reacts faster than B (b) B reacts faster than A
 (c) Both A and B reacts at the same rate (d) Neither A nor B reacts

78. What are reactant X and product Y in the following sequence of reactions ?



- Reactant X Product Y
- (a)
- (b)
- (c)
- (d)



79. Transition state of given S_N2 is :



- (a)
- (b)

272


ORGANIC Chemistry for IIT-JEE



80. $\text{C}_6\text{H}_{13}\text{Br} + \text{OH}^- \longrightarrow \text{C}_6\text{H}_{13}\text{OH} + \text{Br}^-$ is an example of:

- (a) Nucleophilic addition (b) Nucleophilic substitution
(c) Electrophilic addition (d) Electrophilic substitution
(e) Free radical substitution

81. Transition state 2 is structurally most likely as :



- (a) intermediate 1 (b) transition state 3
(c) intermediate 2 (d) product

ANSWERS — LEVEL 1

1.	(a)	2.	(b)	3.	(c)	4.	(a)	5.	(b)	6.	(b)	7.	(b)	8.	(d)
9.	(b)	10.	(a)	11.	(c)	12.	(a)	13.	(d)	14.	(d)	15.	(b)	16.	(c)
17.	(a)	18.	(a)	19.	(c)	20.	(d)	21.	(b)	22.	(a)	23.	(c)	24.	(c)
25.	(d)	26.	(d)	27.	(d)	28.	(a)	29.	(a)	30.	(d)	31.	(b)	32.	(d)
33.	(d)	34.	(c)	35.	A(a)	35.	B(b)	36.	(d)	37.	(b)	38.	(d)	39.	(b)
40.	(d)	41.	(d)	42.	(a)	43.	(c)	44.	(b)	45.	(c)	46.	(d)	47.	(a)
48.	(b)	49.	(b)	50.	(d)	51.	(b)	52.	(d)	53.	(a)	54.	(b)	55.	(a)
56.	(a)	57.	(b)	58.	(c)	59.	(c)	60.	(b)	61.	(c)	62.	(d)	63.	(a)
64.	(b)	65.	(c)	66.	(c)	67.	(d)	68.	(c)	69.	(d)	70.	(c)	71.	(c)
72.	(b)	73.	(d)	74.	(b)	75.	(c)	76.	(d)	77.	(b)	78.	(b)	79.	(d)
80.	(b)	81.	(c)												

ALKYL HALIDES (SUBSTITUTION)

273



1. **Statement-1** : Nucleophilicity order in polar-protic solvent is $I^- < Br^- < Cl^- < F^-$

Statement-2 : Due to bigger size of I^- it is less solvated in polar-protic solvent.



- (a) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.
 (b) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.
 (c) Statement-1 is true, statement-2 is false.
 (d) Statement-1 is false, statement-2 is true.

2. **Statement - 1** : $CH_3 - CH_2 - Cl + NaI \xrightarrow{\text{Acetone}} CH_3 - CH_2 - I + NaCl \downarrow$

Statement- 2 : Acetone is polar-protic solvent and solubility order of sodium halides decreases dramatically in order $NaI > NaBr > NaCl$. The last being virtually insoluble in this solvent and a 1° and 2° chloro alkane in acetone is completely driven to the side of Iodoalkane by the precipitation reaction.

- (a) Statement-1 is true, Statement-2 is true and Statement-2 is correct explanation for statement-1.
 (b) Statement-1 is true, Statement-2 is true and Statement-2 is Not the correct explanation for statement-1.
 (c) Statement-1 is true, Statement-2 is false.
 (d) Statement-1 is false, Statement-2 is true.
3. Encircle whichever of the following :
- (a) is the stronger nucleophile (aprotic solvent) : F^- or I^-
 (b) is the stronger nucleophile (protic solvent) : F^- or I^-
 (c) is the stronger base : F^- or I^-
 (d) is the stronger nucleophile (protic solvent) : NH_3 or NH_2NH_2
 (e) is the better leaving group : CH_3COO^- or $CH_3SO_3^-$
4. Encircle whichever of the following :


- (a) undergoes an S_N2 reaction more rapidly, $CH_3 - Br$ or $CH_3 - \overset{\overset{Br}{|}}{CH} - CH_3$
 (b) undergoes an S_N1 reaction more rapidly, $CH_3 - Br$ or $CH_3 - \overset{\overset{Br}{|}}{CH} - CH_3$
 (c) undergoes an E_2 reaction to give (Z)-1,2-diphenylpropene : $\begin{array}{c} \text{Ph} \\ \diagdown \\ \text{C} = \text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{CH}_3 \end{array}$




274

ORGANIC Chemistry for IIT-JEE






(d) reacts with NaI to give (Z)-1,2-diphenylpropene :



(e) undergoes an $\text{S}_{\text{N}}1$ reaction more rapidly,



5. Encircle whichever of the following :



ALKYL HALIDES (SUBSTITUTION)**275****6. Match the column :**

Alkyl halide			Relative rate (S_{N1})		Relative rate (S_{N2})
(a)	$\text{CH}_3 - \text{Br}$	(p)	1	(w)	1200
(b)	$\text{CH}_3 - \text{CH}_2 - \text{Br}$	(q)	1.05	(x)	40
(c)	$\begin{array}{c} \text{CH}_3 - \text{CH} - \text{Br} \\ \\ \text{CH}_3 \end{array}$	(r)	11	(y)	16
(d)	$\begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3 - \text{C} - \text{Br} \\ \\ \text{CH}_3 \end{array}$	(s)	1,200000	(z)	1





7. Matrix :

Column (I)		Column (II)	
Compound		Type of reaction	
(a)		(p)	S_{N1} reaction can take place
(b)		(q)	S_{N2} reaction can take place
(c)		(r)	S_{N1} is not possible
(d)		(s)	S_{N2} is not possible


276

ORGANIC Chemistry for IIT-JEE

8. Encircle whichever of the following :

9. Reactivity : Circle the reaction that reacts FASTER by S_N2 in each pair :

10. Consider the potential energy diagram given below



(X) Name the positions A-D


(Y) Answer the following questions .

- (i) Both reaction pathways are : EXOTHERMIC or ENDOTHERMIC
 (ii) Which step is the rate determining step (RDS) ? B or D
 (iii) Which product is most stable ? E or F
 (iv) In accordance with Hammonds postulate, exothermic reactions tend to have
 (a) early transition states that are reactant - like
 (b) late transition states that are reactant-like
 (c) early transition states that are product-like
 (d) late transition states that are product-like.


ALKYL HALIDES (SUBSTITUTION)

277


11. Select whether the following combinations of reactants will react by substitution (S_N1 or S_N2 mechanism), elimination (E_1 or E_2 mechanism)



- (a) S_N1 (b) S_N2 (c) E_1 (d) E_2




- (a) S_N1 (b) S_N2 (c) E_1 (d) E_2




- (a) S_N1 (b) S_N2 (c) E_1 (d) E_2




- (a) S_N1 (b) S_N2 (c) E_1 (d) E_2



- (a) S_N1 (b) S_N2 (c) E_1 (d) E_2



- (a) S_N1 (b) S_N2 (c) E_1 (d) E_2









- (a) S_N1 (b) S_N2 (c) E_1 (d) E_2

278

ORGANIC Chemistry for IIT-JEE

12. Examine the ten structural formulas shown in fig. & select that satisfy each of the following conditions. Write one or more (a through j) in each answer box.

(a)		(b)	$\begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\text{C}-\text{Cl} \\ \\ \text{CH}_3 \end{array}$	(c)	
(d)	CH_3-I	(e)		(f)	
(g)	$\begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\text{C}-\text{CH}_2-\text{Cl} \\ \\ \text{CH}_3 \end{array}$	(h)	$\begin{array}{c} \text{H}_2\text{C}=\text{C}-\text{CH}_2-\text{Cl} \\ \\ \text{CH}_3 \end{array}$	(i)	
(j)					

- A. Which compounds give an $\text{S}_{\text{N}}2$ substitution reaction on treatment with alcoholic NaSH ?
 B. Which compounds give an $\text{E}2$ elimination reaction on treatment with alcoholic KOH ?
 C. Which compounds do not react under either of the previous reaction conditions ?

ALKYL HALIDES



279

13. Select which reaction from the following reaction pairs will occur faster.






PART - 1	
Reaction A	<chem>CC1(I)CCCCC1.O>[DMSO]>CC1(O)CCCCC1</chem>
Reaction B	<chem>C1(I)CCCCC1.O>[DMSO]>C1(O)CCCCC1</chem>
PART - 2	
Reaction C	<chem>CC1(Cl)CCCCC1.[Na]I>[DMSO]>CC1(I)CCCCC1</chem>
Reaction D	<chem>C1CCCCC1CCl.[Na]I>[DMSO]>C1CCCCC1CI</chem>
PART - 3	
Reaction E	<chem>C1(I)CCCCC1.[Na]Cl>[DMSO]>C1(Cl)CCCCC1</chem>
Reaction F	<chem>C1(I)CCCCC1.[Na]Cl>[EtOH]>C1(Cl)CCCCC1</chem>
PART - 4	
Reaction G	<chem>C1(I)CCCCC1.[Na][N]=[N]=[N]>[DMSO]>C1([N]=[N]=[N])CCCCC1</chem>
Reaction H	<chem>C1(Br)CCCCC1.[Na][N]=[N]=[N]>[DMSO]>C1([N]=[N]=[N])CCCCC1</chem>

280

ORGANIC Chemistry for IIT-JEE







PART - 5	
Reaction I	
Reaction J	

14. Tick your answer in the given box.

Alkyl Halide		Expect S_N2 (at a reasonable rate)	
	2-D Structure		
(a)			Yes
			No
(b)			Yes
			No
(c)			Yes
			No
(d)			Yes
			No
(e)			Yes
			No

ALKYL HALIDES

281

(f)	Bromocyclohexane			Yes
				No
(g)	Bromobenzene			Yes
				No
(h)	Benzyl bromide			Yes
				No
(i)	1-Bromo-2,2-dimethyl propane			Yes
				No
(j)	Bicyclo compound			Yes
				No
(k)	1-bromotriptycene			Yes
				No


282

ORGANIC Chemistry for IIT-JEE

15. Match the column

Column-I		Column-II	
(a)		(p)	It will undergo Nucleophilic Substitution reaction
(b)		(q)	It will undergo E_2 reaction
(c)		(r)	It will undergo E_1 reaction
(d)		(s)	It will undergo S_N2 reaction
		(t)	It will undergo S_N1 reaction

16.



How many (x) moles of HI consumed?

ALKYL HALIDES








ANSWERS — LEVEL 2

283

ANSWERS — LEVEL 2




1. d

2. c The reaction is Finkelstein reaction.

3. (a) F^- ; (b) I^- ; (c) F^- ; (d) $\text{NH}_2\text{--NH}_2$; (e) CH_3SO_3^- 4. (a) $\text{CH}_3\text{--Br}$ 

6. a - p, w; b - q, x; c - r, y; d - s, z

7. a - r, s; b - p, q; c - r, s; d - r, s




10. (X) A- reactants, B-transition state, C-Inter mediate, D- transition state


(Y) (i) exothermic (ii) B (iii) F (iv) a

286

ORGANIC Chemistry for IIT-JEE




when (P) undergoes Hoffmann exhaustive methylation (twice) then the product obtained will be :











Product (A) & (B) of the above reaction is :

- (a) $A = P, B = P$ (b) $A = Q, B = Q$ (c) $A = P, B = Q$ (d) $A = Q, B = P$




6. Which of these dehydrates most easily ?




(d) No reaction

ALKYL HALIDES (ELIMINATION)


287





Major product A is :



Major product A is :



10. In which of the following reaction Saytzeff alkene is major product ?



12. The conversion of 2, 3-dibromobutane to 2-butene with Zn is :




- (a) Redox reaction (b) α -Elimination
(c) β -Elimination (d) Both α -elimination and redox reaction

288

ORGANIC Chemistry for IIT-JEE




13. 1, 3-Dibromopropane is heated with zinc dust in ether. The product formed is :

- (a) propene
(b) propane
(c) cyclopropane
(d) 3-bromopropane



Product obtained in above reactions (1), (2) & (3) is :

- (a) A = B but C is different
(b) A = C, but B is different
(c) B = C, but A is different
(d) A = B = C all product are identical




Product obtained in above reactions (1), (2) & (3) is :

- (a) A = B, C is different
(b) A = C, B is different
(c) B = C, A is different
(d) A = B = C is same

ALKYL HALIDES (ELIMINATION)

289




Find (x).

(a) 3





(b) 4

(c) 5





(d) 6




Major product (A) is :







18. Which one of the following compound will be least susceptible to elimination of hydrogen bromide ?








19. Two alkenes, X(91% yield) and Y(9% yield) are formed when the following compound is heated.



The structures of X and Y, respectively are :







20. In the dehydrohalogenation of 2-bromobutane; which conformation leads to the formation of cis-2-butene ?









290

ORGANIC Chemistry for IIT-JEE





Product (B) of given reaction is:



22. What product will be formed from Hoffmann exhaustive methylation of following compound?



Product (A) is :








(a) Racemic

(b) Diastereomers

(c) G.I


(d) Positional isomers















ALKYL HALIDES (ELIMINATION)

291






26.



Major product of the reaction is :

27. The E_2 product of the following reaction will be ?28. An alkyl halide $C_5H_{11}Br$ on treatment with alc. KOH give 2-pentene only. The halide will be






29. End product (D) in the given sequence is:








- (c) F is the SnH_3PO_4 Major product obtained by dehydration of given alcohol is :
- for react Δ with H_2SO_4 as the leaving group has the react soluble solid character

ALKYL HALIDES (ELIMINATION)




295








Product (B) of above reaction is :



Product of the above reaction is :








Product (A) is :








Major product of the above reaction is :

296








ORGANIC Chemistry for IIT-JEE



LDA = Lithium di-isopropyl amide





Major product of the reaction, when the given compound undergoes Hoffmann exhaustive methylation is :




ALKYL HALIDES (ELIMINATION)

297



49.



Stereochemistry of the product is :



(a) Meso compound

(b) Racemic mixture






(c) Diastereomer

(d) Optically pure enantiomers




50.

Which of the following reactant is used to obtain above compound (A). (Assume that EtO^- is used in all the reaction)

51.










52.

Rank the following in order of decreasing rate in an E_2 reaction:(a) $a > b > c$ (b) $c > a > b$ (c) $c > b > a$ (d) $b > a > c$

298

ORGANIC Chemistry for IIT-JEE








Relation between (A) and (B) is :

(a) G.I.


(b) Positional isomer

(c) Enantiomer

(d) Chain isomer



56. Which best describes the product of the following reaction ?




(a) Absolute configuration has been inverted

(b) Absolute configuration has been retained

(c) Racemization (loss of absolute configuration) has occurred





(d) Loss of chirality has occurred (the product is achiral)

57. What is the major product of the following reaction ?




ALKYL HALIDES (ELIMINATION)

299



58. What will be the major product of each of the two reaction shown below ?




(a) 1 - X, 2 - X

(b) 1 - Y, 2 - X

(c) 1 - X, 2 - Y

(d) 1 - Y, 2 - Y




(a) a primary amine






(b) a tertiary amine

(c) a secondary amine

(d) a quaternary ammonium salt







(A) on heating isomerizes to (B). What is the structure of (B) ?







300




ORGANIC Chemistry for IIT-JEE



62. Which of the following carbocation will undergo rearrangement ?






63. In which of the following reaction resonance stabilized product will form ?



(d) All of these


64. In which of following reaction rearrangement take place with change in carbon skeleton ?



ALKYL HALIDES (ELIMINATION)


301

65. Consider the following reaction :




Which response contains all the correct statement about this process ?

- (1) Dehydration (2) E₂ mechanism
 (3) Carbon skeleton migration (4) Most stable alkene will form
 (5) Single-step reaction
 (a) 1, 3 (b) 1, 2, 3 (c) 1, 2, 5 (d) 1, 3, 4







- (a) *cis*-2-butene (b) *trans*-2-butene
 (c) 1-butene (d) Iso-butene



Above reaction is an example of 1,4-elimination. Predict the product.






- (a) $\text{CH}_3 - \text{CH} = \text{C} = \text{CH}_2$ (b) $\text{CH}_3 - \text{C} \equiv \text{C} - \text{CH}_3$
 (c) $\text{CH}_3 - \text{CH}_2 - \text{C} \equiv \text{CH}$ (d) $\text{H}_2\text{C} = \text{CH} - \text{CH} = \text{CH}_2$



(d) None of these

302

ORGANIC Chemistry for IIT-JEE



When (A) and (B) reacts with H_2SO_4 products obtained are :





(a) p, q, r, s

(b) p, q

(c) p, q, r

(d) p, q, s

70. Which of the following compound gives even number of Hoffmann's exhaustive methylation and elimination?



ANSWERS — LEVEL 1

1.	(d)	2.	A - a B - d	3.	(a)	4.	(c)	5.	(b)	6.	(b)	7.	(d)	8.	(b)
9.	(c)	10.	(d)	11.	(c)	12.	(c)	13.	(c)	14.	(d)	15.	(d)	16.	(c)
17.	(b)	18.	(b)	19.	(c)	20.	(a)	21.	(b)	22.	(a)	23.	(b)	24.	(a)
25.	(b)	26.	(a)	27.	(d)	28.	(c)	29.	(b)	30.	(c)	31.	(d)	32.	(d)
33.	(a)	34.	(a)	35.	(a)	36.	(b)	37.	(c)	38.	(c)	39.	(c)	40.	(c)
41.	(c)	42.	(b)	43.	(b)	44.	(b)	45.	(c)	46.	(a)	47.	(d)	48.	(b)
49.	(b)	50.	(a)	51.	(c)	52.	(b)	53.	(b)	54.	(b)	55.	(c)	56.	(d)
57.	(d)	58.	(b)	59.	(d)	60.	(b)	61.	(a)	62.	(b)	63.	(d)	64.	(a)
65.	(d)	66.	(b)	67.	(d)	68.	(b)	69.	(b)	70.	(a,b)				

ALKYL HALIDES (ELIMINATION)


303




1. Comprehension

E_2 reaction $\rightarrow$ Elimination bimolecular


In the general mechanism of the E_2 reaction a strong base abstract a proton on a carbon atom adjacent to the one of the leaving group. As the base abstracts a proton, a double bond forms and the leaving group leaves.



Mechanism :







A. Identify the rate of reaction of given compounds in E_2 reaction:











(a) $a > b > c > d$ (b) $a > c > b > d$ (c) $b > a > c > d$ (d) $b > d > a > c$

B. In given pairs, which compound is more reactive toward E_2 reaction:








304






ORGANIC Chemistry for IIT-JEE







Product (A) and (B) are :







Select the products (A) and (B) from the compounds (P) and (Q) given below:



E. Which of the following compound is inert toward E_2 reaction.







ALKYL HALIDES (ELIMINATION)**305****2. Match the column :**

Column (I)		Column (II)	
E_2 reaction (elimination bimolecular)		No. of possible products. (including stereoisomerism)	
(a)		(p)	0
(b)		(q)	1
(c)		(r)	2
(d)		(s)	3

3. Match the Column :





HEM = Hoffmann exhaustive methylation followed by elimination.

Column (I)		Column (II)	
Reaction		Product	
(a)		(p)	$H_2C = CH - CH_2 - CH = CH_2$
(b)		(q)	$H_2C = CH - CH_2 - CH_2 - CH = CH_2$
(c)		(r)	$H_2C = CH - CH_2 - \overset{\overset{CH_3}{ }}{C} = CH_2$
(d)		(s)	$H_2C = CH - \overset{\overset{CH_3}{ }}{CH} - CH = CH_2$





306

ORGANIC Chemistry for IIT-JEE

4. Match the column :

Column (I)		Column (II)	
(a)		(p)	Product are Diastereomers
(b)		(q)	Carbocation is intermediate
(c)		(r)	2nd order reaction
(d)		(s)	1st order reaction

5. Match the column :

Column (I)		Column (II)	
(a)		(p)	Optically active product
(b)		(q)	Optically inactive product
(c)		(r)	2nd order reaction
(d)		(s)	unimolecular reaction

ALKYL HALIDES (ELIMINATION)

307

6. Match the column :

Column (I)		Column (II)	
E₂ reactions (elimination bimolecular)		Number of products (including stereoisomerism)	
(a)	$\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{Br} \xrightarrow{\text{alc. KOH}}$	(p)	1
(b)	$\text{CH}_3 - \underset{\text{Br}}{\text{CH}} - \text{CH}_2 - \text{CH}_3 \xrightarrow{\text{alc. KOH}}$	(q)	2
(c)	$\text{CH}_3 - \underset{\text{Br}}{\underset{\text{CH}_3}{\text{C}}} - \text{CH}_2 - \text{CH}_3 \xrightarrow{\text{alc. KOH}}$	(r)	3
(d)	$\text{Ph} - \text{CH}_2 - \underset{\text{Br}}{\text{CH}} - \text{CH}_2 - \text{CH}_3$	(s)	4


7. Match the column :

Column (I)		Column (II)	
(a)	$\text{CH}_3 - \text{CH}_2 - \underset{\text{OH}}{\text{CH}} - \text{CH}_3 \xrightarrow[\Delta]{\text{H}^+} \text{(A)}$	(p)	E ₁
(b)	$\text{CH}_3 - \text{CH}_2 - \underset{\text{Cl}}{\text{CH}} - \text{CH}_3 \xrightarrow[\Delta]{\text{NaNH}_2}$	(q)	E ₂
(c)	$\text{CH}_3 - \overset{\text{O}}{\parallel}{\text{C}} - \text{CH}_2 - \underset{\text{Br}}{\text{CH}} - \text{CH}_3 \xrightarrow[\Delta]{\text{EtONa}}$	(r)	Ei (elimination intramolecular)
(d)	$\text{Cyclohexyl} - \text{N}^+(\text{Me})_2 - \text{O}^- \xrightarrow{\Delta}$	(s)	E ₁ CB


310

ORGANIC Chemistry for IIT-JEE


13. Reaction-1







Reaction-2



Reaction-3

Sum of α -hydrogen (A + B + C) =

14.







Total number of products obtained in above reaction including minor products is (including stereoisomer)

15. Match the column (I) and (II).



Column (I)		Column (II)	
Reaction		Type of Reaction	
(a)	R - 2 - chlorobutane $\xrightarrow[\text{acetone}]{\text{KSH}}$	(p)	S_N1
(b)	R - 2 - chlorobutane $\xrightarrow[\text{EtOH}]{\text{EtO}^-\text{Na}^+}$	(q)	S_N2
(c)	2 - bromo - 2 - methyl propane $\xrightarrow{\text{H}_2\text{O}}$	(r)	E_1
(d)	2 - butanol $\xrightarrow[\Delta]{\text{H}_2\text{SO}_4}$	(s)	E_2

(C) + (E) + (A) + (B) + (D) + (F) + (G) + (H) + (I) + (J) + (K) + (L) + (M) + (N) + (O) + (P) + (Q) + (R) + (S) + (T) + (U) + (V) + (W) + (X) + (Y) + (Z)

ALKYL HALIDES (ELIMINATION)**311****16. Match the column (I) and (II).**


Column (I)		Column (II)	
	Reaction		Type of Reaction
(a)		(p)	S_N1
(b)		(q)	S_N2
(c)		(r)	E_1
(d)		(s)	E_2

17. Select whether the following reagent combination will result in elimination or substitution reactions leading to the major product.




	Reaction	Substitution	Elimination
(a)			
(b)			

312

ORGANIC Chemistry for IIT-JEE


(c)	$\text{CH}_3 - \overset{\text{Cl}}{\underset{ }{\text{CH}}} - \text{CH}_2 - \text{CH}_3 \xrightarrow{\text{alc-KOH}}$		
(d)	$\text{CH}_3 - \overset{\text{CH}_3}{\underset{\text{H}}{\underset{ }{\text{C}}}} - \text{I} \xrightarrow{\text{Na}^+ \text{N}_3^-}$		
(e)			
(f)	$\text{CH}_3 - \overset{\text{CH}_3}{\underset{\text{CH}_3}{\underset{ }{\text{C}}}} - \text{Cl} \xrightarrow{\text{H}_2\text{O}}$		

18. Match the Column (I) and (II) (Matrix).







Column (I)		Column (II)	
	Reaction		Comment on product
(a)		(p)	Racemic mixture
(b)		(q)	Major product consist of even number of α -hydrogen
(c)		(r)	Will not undergo dehydration

ALKYL HALIDES (ELIMINATION)


313

(d)		(s)	Major product consist of odd number of α -hydrogen


19. For each of the following amines (A through D), exhaustive methylation (treatment with excess methyl iodide), followed by Hoffmann elimination (heating with AgOH), repeated as necessary, removes the nitrogen atom in the form of trimethylamine. Indicate the number of repetitive Hoffmann eliminations required to remove the nitrogen by a number (1 to 4) in the designated answer sheet.

A.		B.		C.	
D.		E.		F.	
a.		b.		c.	
d.		e.		f.	

20.




$\longrightarrow$ x is total number of HEM (Hoffman Exhaustive Methylation and eliminations) to remove nitrogen from given compound.



$\xrightarrow{\text{alc. KOH}}$ y is total number of possible E_2 product (including stereoisomer)

Sum of $x+y = ?$

21.



$\xrightarrow{\text{EtOH}}$ (x) ($S_N1 + E_1$) products. (including stereoisomer)
consider all products


Total number of products are :

314

ORGANIC Chemistry for IIT-JEE






ANSWERS — LEVEL 2

1. A – a; B – a; C – b; D – c; E – c;
2. a – s; b – r; c – q; d – p
3. a – s; b – r; c – q; d – p
4. a – p, q, s; b – p, r; c – q, s; d – r
5. a – p, r; b – p, r; c – p, r; d – q, r
6. a – p; b – r; c – q; d – s
7. a – p; b – q; c – s; d – r
8. a – p; b – q; c – r; d – s
9. $X = 3, Y = 3, Z = 2, P = 0 \Rightarrow 3 + 3 + 2 + 0 = 8$
10. $x = 3, y = 2, z = 3 \Rightarrow 3 + 2 + 3 = 8$
11. 32
12. 33
13. 28
14. $x = 3, y = 1, z = 3, p = 2$
Sum = 9
15. (a – q), (b – s), (c – p), (d – r)
16. (a – q), (b – s), (c – p), (d – r)
17. Substitution – d, f
Elimination – a, b, c, e
18. a – p, q; b – p, q; c – q; d – r
19. a – 3; b – 2; c – 3; d – 1; e – 2; f – 3
20. 6
21. 6



5C



ALKYL HALIDES



316






ORGANIC Chemistry for IIT-JEE

2. In the reactions given below,




the compounds A and B are :





- (a) chain isomers (b) position isomers
(c) functional isomers (d) metamers
3. Which is the major product expected from the following S_N2 reaction ?



4. Consider the following E_1 , S_N1 reaction :



The missing product(s) is(are) :








- (a) 1 and 3 (b) 3 and 4 (c) 2 and 3 (d) 1, 2, 3 and 4






ALKYL HALIDES

317

5. What is the product of the following S_N2 reaction ?



6. Select the reagent that will yield the greater amount of substitution on reaction with $\text{CH}_3-\text{CH}_2-\text{Br}$:
- $\text{CH}_3\text{CH}_2\text{OK}$ in dimethyl sulfoxide (DMSO)
 - $(\text{CH}_3)_3\text{COK}$ in dimethyl sulfoxide (DMSO)
 - Both (a) and (b) will give comparable amounts of substitution
 - Neither (a) nor (b) will give any amount of substitution
7. Under the specified conditions, substrate X undergoes substitution and elimination reactions to give products $A-D$. A and B are stereoisomers, but not enantiomers. C and D are enantiomers. A is not an isomer of C . Which of the following could be the starting material X ?






(a) (I)

(b) (II)

(c) (III)





(d) (IV)

Compare rate of E_2 reaction :(a) $c > b > a$ (b) $a > b > c$ (c) $b > a > c$ (d) $c > a > b$


318

ORGANIC Chemistry for IIT-JEE

9. Which reaction results in the formation of a pair of enantiomers ?





10. Rate limiting S_N1 follows the sequence




True statement about sequence on the basis of assumption that R contains 3 different groups is :

- (a) more stable carbocation, greater is in the proportion of racemization
- (b) the more nucleophilic the solvent greater in the proportion of inversion
- (c) In above sequence (b) represent separately solvated, pair of ions
- (d) All of these

11. Compare the two methods shown for the preparation of carboxylic acids :



Which one of the following statements correctly describes this conversion ?



- (a) Both method 1 and method 2 are appropriate for carrying out this conversion
- (b) Neither method 1 nor method 2 is appropriate for carrying out this conversion
- (c) Method 1 will work well, but method 2 is not appropriate
- (d) Method 2 will work well, but method 1 is not appropriate





12. Which of the following statements is true ?

- (a) $CH_3CH_2S^-$ is both a stronger base and more nucleophilic than $CH_3CH_2O^-$
- (b) $CH_3CH_2S^-$ is a stronger base but is less nucleophilic than $CH_3CH_2O^-$
- (c) $CH_3CH_2S^-$ is a weaker base but is more nucleophilic than $CH_3CH_2O^-$
- (d) $CH_3CH_2S^-$ is both a weaker base and less nucleophilic than $CH_3CH_2O^-$


ALKYL HALIDES

319


13. In the given pair of alcohols, in which pair second alcohol is more reactive than first towards hydrogen bromide ?



14. Which product would be expected to predominate in the given reaction ?





15. Which is the major product of the following reaction ?




320




ORGANIC Chemistry for IIT-JEE



16.




$\xrightarrow{\text{Alc. KOH}}$ (A) ; Major product of this reaction is :




17. Rate of S_N2 reaction is :



(A)



(B)



(C)

(a) (B) > (A) > (C)

(b) (C) > (A) > (B)


(c) (A) > (B) > (C)

(d) (A) > (C) > (B)


18.

1,2-dichloro ethane $\xrightarrow{\text{Na}, \text{CH}_3\text{CH}_2\text{MgI}}$ $\text{C}_4\text{H}_8\text{S}_2$ (P)



Unknown product (P) of the above reaction is :




(a)



(b)




19.




$\xrightarrow{\text{Moist Ag}_2\text{O}}$ (A) product


Major product (A) is :




(a)



(b)




(c)



(d)

20.






(MOM chloride)
(Methoxy methyl chloride)





$\xrightarrow{\text{KCN}}$ Product of reaction is :

ALKYL HALIDES




321



21. In the given pair of compound, in which pair the second compound is more reactive than first toward S_N2 reaction ?






22. Which compound might be synthesized by the S_N2 displacement of an alkyl-halide ?




(d) All of these

23. Identify C in the following series $C_3H_7I \xrightarrow[\text{alc.}]{KOH} A \xrightarrow[\Delta]{NBS} B \xrightarrow[\text{alc.}]{KCN} C$.




24. What sequence of reagents is required to accomplish the following transformation ?



- (a) (1) NBS, ROOR (2) $CH_3CH_2O^-$ (3) 2HBr (4) NH_2^- (5) disiamyl borane (6) H_2O_2, OH^-
 (b) (1) $Cl_2, h\nu$ (2) OH^- , heat; (3) 2HCl (4) OH^- , heat (5) $HgSO_4, H_2SO_4$
 (c) (1) NBS, ROOR; OH^- , DMSO
 (d) (1) $Br_2, h\nu$ (2) *t*-butoxide (3) BH_3 , THF (4) H_2O_2, OH^-

25. Which of the reagents shown below would accomplish the following transformations?




A

- (a) H_3O^+
 (b) NaOH
 (c) HBr in ether
 (d) $NaNH_2$

B


- $BH_3 - THF; H_2O_2/NaOH$
 $BH_3 - THF; H_2O_2/NaOH$
 $Hg(OAc)_2/H_2O; NaBH_4$
 $Hg(OAc)_2/H_2O; NaBH_4$

26. What are the products obtained from the following reaction ?




322

ORGANIC Chemistry for IIT-JEE



27. The back-side attack on 2-bromobutane by methoxide (CH_3O^-) gives the product shown below. Which Fischer projection represents 2-bromobutane used as the reactant in this reaction ?




28. Consider the following statements :

- (1) Bridgehead halides are inert towards both $\text{S}_{\text{N}}1$ and $\text{S}_{\text{N}}2$ reactions (till one of the ring size is eight member ring)
- (2) The first step in both $\text{S}_{\text{N}}1$ and E_1 reactions is the same
- (3) $\text{S}_{\text{N}}2$ reactions proceed with total retention of configuration
- (4) E_2 eliminations are by the use of a solvent of low polarity and high concentration of a strong base

Which of the above statements are correct?

- (a) 1, 2 and 4 (b) 1 and 3
(c) 2, 3 and 4 (d) 1, 2, 3 and 4

29. Consider the following alcohols :



The order of decreasing reactivities of these alcohols towards substitution with HBr is :

- (a) III > I > IV > II (b) III > I > II > IV
(c) I > III > IV > II (d) I > III > II > IV

ALKYL HALIDES

323





30. In solvolysis of 1,2-dimethyl propyl *p*-toluene sulfonate in acetic acid at 75°C, how many (alkene + substitution) products will be formed ?

- (a) 2 (b) 3 (c) 4 (d) 5


31. Benzotrichloride reacts with milk of lime to form :

- (a) Benzal (b) Benzoic acid (c) Benzyl alcohol (d) Phenol

32. $\text{Br} - \text{CH}_2 - (\text{CH}_2)_2 - \text{CH}_2 - \text{Br} + \text{CH}_3\text{NH}_2 \longrightarrow$ Product of the reaction is :







33. The configurations of the reactant and the product in the following reaction, respectively, are:







- (a) R, R (b) R, S (c) S, R (d) S, S

34. 1-4-dichlorohexane (1 mole) + NaI (1 mole) $\xrightarrow{\text{Acetone}}$ Product of the reaction is :



35. Alkyl halides can be obtained by all methods except :



36. In order to prepare 1-chloropropane, which of the following reactants can be employed ?





(a) Propene and HCl in the presence of peroxide

(b) Propene and Cl_2 followed by treatment with aq. KOH




(c) Propanol-1 and SOCl_2 /pyridine

(d) Any of the above can be used

37. Which alkyl halide has maximum density ?






38. Which of the following molecules would have a carbon-halogen bond most susceptible to nucleophilic substitution ?




324

ORGANIC Chemistry for IIT-JEE

39. When benzyl chloride is treated with ethanolic KCN, the major product formed is :
 (a) benzyl ethyl ether (b) benzyl alcohol (c) benzyl cyanide (d) benzyl isocyanide
40. Which of the following is most reactive towards nucleophilic substitution reaction ?
 (a) $\text{CH}_2 = \text{CH} - \text{Cl}$ (b) $\text{C}_6\text{H}_5\text{Cl}$
 (c) $\text{CH}_3\text{CH} = \text{CHCl}$ (d) $\text{ClCH}_2 - \text{CH} = \text{CH}_2$
41. Which of the following reaction will not give ether as a major product ?
 (a) $\text{CH}_3\text{CH}_2\text{Cl} + \text{Ag}_2\text{O}(\text{dry}) \longrightarrow$ (b) $(\text{CH}_3)_3\text{CCl} + \text{CH}_3\text{CH}_2\text{O}^- \text{Na}^+ \longrightarrow$



Product (A) and (B) in above reaction is :



ANSWERS — LEVEL 1

1.	(c)	2.	(c)	3.	(b)	4.	(a)	5.	(a)	6.	(a)	7.	(c)	8.	(b)
9.	(b)	10.	(d)	11.	(c)	12.	(c)	13.	(d)	14.	(a)	15.	(c)	16.	(c)
17.	(c)	18.	(a)	19.	(c)	20.	(b)	21.	(d)	22.	(d)	23.	(b)	24.	(d)
25.	(d)	26.	(b)	27.	(d)	28.	(a)	29.	(a)	30.	(d)	31.	(b)	32.	(b)
33.	(d)	34.	(d)	35.	(c)	36.	(c)	37.	(a)	38.	(d)	39.	(c)	40.	(d)
41.	(b)	42.	(b)												

ALKYL HALIDES

325



1. The following organic halide derivatives (A to J) are reacted in ethanol solution with each of the nucleophiles : acetate, methylthiolate, cyanide and hydroxide anions. Six possible results from these combinations of reactants are designated (1) through (6) below :

Write the number corresponding to your best estimate of the outcome of each reaction in the appropriate answer box below.

A	B	C	D	E
F	G	H	I	J

Possible Outcome :

(1) No reaction	(2) Substitution
(3) Elimination	(4) Substitution and elimination
(5) No reaction or slow substitution	(6) No reaction or slow elimination

Compound		A	B	C	D	E	F	G	H	I	J
(i)	$\text{CH}_3\text{CO}_2\text{Na}$										
(ii)	CH_3SNa										
(iii)	NaCN										
(iv)	NaOH										

326

ORGANIC Chemistry for IIT-JEE

2. In each of the following sections three organic halogen compounds are listed. In the box given enter a number (1 to 3) indicating the order of reactivity of the designated (1 is most reactive and 3 is least).

(a) S_N2 substitution by NaOCOCH_3 in methanol:

1. $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$ ☐ 2. $(\text{CH}_3)_2\text{CHBr}$ ☐ 3. $\text{CH}_2=\text{CHCH}_2\text{Br}$ ☐

(b) S_N2 substitution by NaI in acetone:

1. $\text{C}_6\text{H}_5\text{Cl}$ ☐ 2. $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$ ☐ 3. $\text{C}_6\text{H}_5\text{CHClCH}_3$ ☐

(c) S_N2 substitution by NaCN in methanol:

1. $\text{CH}_3\text{CH}_2\text{Cl}$ ☐ 2. $\text{CH}_3\text{CH}_2\text{F}$ ☐ 3. $\text{CH}_3\text{CH}_2\text{I}$ ☐

(d) S_N2 substitution by NaSCH_3 in methanol:

1. $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{Br}$ ☐ 2. $\text{CH}_3\text{CH}_2\text{CHBrCH}_2\text{CH}_3$ ☐ 3. $(\text{CH}_3)_3\text{CCH}_2\text{Br}$ ☐

3. Isobutyl alcohol (2-methyl-1-propanol), $(\text{CH}_3)_2\text{CHCH}_2\text{OH}$, can be transformed to each of the compounds (a through l) listed in the left-hand column. In each case the number of steps needed to accomplish the change is noted, and an answer box is provided for your reagent selections. Fourteen reagents (designated A through N) are listed in the right-hand column.


Write letters designating the reagent or reagents you believe will achieve the desired transformation in the box to the right of the product formula. In the case of a multi-step sequence write the reagents in the order they are to be used. In some cases you may wish to use a previously prepared compound as a reactant. If so, write the number (a to l) corresponding to the desired compound.

Desired product		No. of steps	Write options	Reagent List	
a.	$(\text{CH}_3)_2\text{CHCH}_2\text{Br}$	one		A.	$\text{Hg}(\text{OAc})_2$ in H_2O
b.	$(\text{CH}_3)_2\text{C}=\text{CH}_2$	one		B.	PBr_3 & heat
c.	$(\text{CH}_3)_2\text{CHCH}=\text{O}$	one		C.	NaBH_4 in alcohol
d.	$(\text{CH}_3)_2\text{CHCO}_2\text{H}$	one		D.	LiAlH_4 in THF (aqueous workup)
e.	$(\text{CH}_3)_3\text{CBr}$	two		E.	NaCN in alcohol
f.	$(\text{CH}_3)_2\text{CHCH}_2\text{C}\equiv\text{N}$	two		F.	PCC in CH_2Cl_2
g.	$(\text{CH}_3)_2\text{CHCH}_2\text{OCOCH}_3$	one		G.	Jones' reagent (CrO_3 in H_3O^+)
h.	$(\text{CH}_3)_2\text{CHCO}_2\text{C}_2\text{H}_5$	two		H.	HBr in CH_2Cl_2
i.	$(\text{CH}_3)_2\text{CHCH}_2\text{OCH}_2(\text{CH}_3)_2$	two		I.	H_3PO_4 and heat
j.	$(\text{CH}_3)_2\text{COH}$	three		J.	$(\text{CH}_3\text{CO})_2\text{O}$ + pyridine
k.	$(\text{CH}_3)_2\text{CHCH}_2\text{NH}_2$	three		K.	NaN_3 in aqueous alcohol
l.	$(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{NH}_2$	two		L.	$\text{C}_6\text{H}_5\text{CO}_3\text{H}$ in CH_2Cl_2 (peracid)
				M.	NaH in ether and heat
				N.	$\text{C}_2\text{H}_5\text{OH}$ + acid catalyst & heat

ALKYL HALIDES**327****ANSWERS — LEVEL 2**

- 1.**
- | | A | B | C | D | E | F | G | H | I | J |
|-------|---|---|---|---|---|---|---|---|---|---|
| (i) | 2 | 2 | 2 | 1 | 1 | 1 | 6 | 2 | 2 | 6 |
| (ii) | 2 | 2 | 2 | 1 | 1 | 5 | 6 | 2 | 2 | 6 |
| (iii) | 2 | 2 | 2 | 1 | 1 | 1 | 3 | 3 | 2 | 3 |
| (iv) | 4 | 2 | 2 | 1 | 1 | 5 | 3 | 3 | 4 | 3 |
- 2.** $a - 3 > 1 > 2$; $b - 2 > 3 > 1$; $c - 3 > 1 > 2$; $d - 1 > 2 > 3$
- 3.** $a - B$; $b - I$; $c - F$; $d - G$; $e - I, H$ or $2 H$; $f - B, E$ or I, E ; $g - J$; $h - G, N$ or $4N$
 $i - N, I$; $j - I, A, C$ or $2AC$ or ILD or $2LD$; $k - B, K, D$ or $1KD$; $l - B, E, D$ or $1ED$ or $6D$

ATDB.uno




6

ALCOHOL, ETHERS AND EPOXIDES



ATDB.uno LEVEL - 1

1. The following transformation involves a carbocation rearrangement. The carbocation is generated by protonation of the hydroxyl group, followed by the loss of water. Which bond has to migrate in the carbocation to yield the product indicated (after the deprotonation) ?




(a) a

(b) b

(c) c






(d) d

2. Identify the major product.








ALCOHOL, ETHERS AND EPOXIDES

329








4. Predict the product when given compound reacts with LiAlH_4 .



330

ORGANIC Chemistry for IIT-JEE






5. Predict the product when given compound (A, in the above question 4) reacts with NaBH_4 .







Methyl benzoate

The labelled O^{18} will be in :

- (a) H_2O (b) Methyl benzoate
(c) Both (a) and (b) (d) Benzoic acid







8. Which is oxidized most easily ?




ALCOHOL, ETHERS AND EPOXIDES

331



- (a)  (b)  (c)  (d) 


10. Which of the following react with HBr at faster rate ?





- (a)  (b) 
 (c)  (d) 








Above conversion can be done by :

- (a) $NaBH_4$ (b) $LiAlH_4$ (c) PCC (d) $KMnO_4$



- (a)  (b) 
 (c)  (d) 








- (a)  (b) 
 (c)  (d) 






332

ORGANIC Chemistry for IIT-JEE





14. Predict the major product of the given reaction.







15. Identify the major product,



16. $\xrightarrow{H^+}$ (A); Product (A) is :









17. $\xrightarrow{LiAlH_4}$ (A) Major; product (A) is :
















ALCOHOL, ETHERS AND EPOXIDES

333







(a) No reaction



334

ORGANIC Chemistry for IIT-JEE



22. Consider the following alcohols,



The order of decreasing reactivities of these alcohols towards nucleophilic substitution with HBr is:

- (a) III > I > IV > II (b) III > I > II > IV (c) I > III > IV > II (d) I > III > II > IV






23.



Sum of number of alcoholic groups in product (P) and (Q) is:







- (a) 1 (b) 2 (c) 3 (d) 5

24. In presence of dil. HCl, compound A is converted to a constitutional isomer (B), compound B is:













ALCOHOL, ETHERS AND EPOXIDES

335








- (a) Racemic
(c) Meso

- (b) Diastereomer
(d) Optically pure




336

ORGANIC Chemistry for IIT-JEE




32. Which is the best reagent to convert isopropyl alcohol to isopropyl bromide ?





ALCOHOL, ETHERS AND EPOXIDES

337




Major product obtained in the above reaction is :



(c) Racemic

(d) Diastereomers













Total number of stereoisomers of product (B) will be:

(a) 2

(b) 3


(c) 4

(d) 5




338

ORGANIC Chemistry for IIT-JEE



(1)



(2)




(3)



(4)


For the above conversion the correct order of reagents used is :

(a) $1 \rightarrow 2 \rightarrow 3 \rightarrow 4$ (b) $1 \rightarrow 4 \rightarrow 3 \rightarrow 2$ (c) $1 \rightarrow 4 \rightarrow 2 \rightarrow 3$ (d) $2 \rightarrow 3 \rightarrow 4 \rightarrow 1$ 

Find missing reagents.

(a) $x = \text{LiAlH}_4, y = \text{NaBH}_4$ (b) $x = \text{LiAlH}_4/\text{AlCl}_3, y = \text{LiAlH}_4$ (c) $x = \text{LiAlH}_4, y = \text{LiAlH}_4/\text{AlCl}_3$ (d) $x = \text{H}_2/\text{Ni}, y = \text{H}_2/\text{Pt}$

39. In solvolysis of 1, 1-dimethylpropyl tosylate in acetic acid at 75°C , (alkene + substitution product) will be formed by mechanism ?

(a) $\text{S}_{\text{N}}2, \text{E}_2$ (b) $\text{S}_{\text{N}}2, \text{E}_1$ (c) $\text{S}_{\text{N}}1, \text{E}_2$ (d) $\text{S}_{\text{N}}1, \text{E}_1$  x = moles of anhydride consumed

(a) 1

(b) 2

(c) 3

(d) 4

41. Identify product when (*R*) – and (*S*) – 2 – butanol reacts with (*R,R*) tartaric acid in acidic medium.

(a) Racemic

(b) Diastereomer


(c) Meso

(d) Pure enantiomer


ALCOHOL, ETHERS AND EPOXIDES

339


42. An alcohol of formula $C_9H_{12}O$ reacts with $Na_2Cr_2O_7$ to form a compound having formula $C_9H_{10}O$. The original alcohol might be :




43. An optically active alcohol of formula $C_9H_{12}O_2$ produced the following compound when refluxed with $KMnO_4$.



The original compound showed these properties also :



What is structure of (A)?



44. Which are not cleaved by HIO_4 ?

I : glycerol

II : glycol

III : 1, 3-propenediol

IV : methoxy-2-propanol





(a) I, II, III, IV

(b) I, II


(c) II, III

(d) III, IV

45. Which of the following reactions require an oxidising agent ?







46. What is the major product of the following reaction ?







340


ORGANIC Chemistry for IIT-JEE



47. Which of the esters shown, after reduction with LiAlH_4 and aqueous workup, will yield two molecules of only a single alcohol ?



48. For the following reaction, select the statement that best describes the change.




(a) The alcohol is oxidized to an acid, and the Cr(VI) is reduced

(b) The alcohol is oxidized to an aldehyde, and the Cr(VI) is reduced

(c) The alcohol is reduced to an aldehyde, and the Cr(III) is oxidized

(d) The alcohol is oxidized to a ketone, and the Cr(VI) is reduced

49. What is the product of the following reaction ?







(a) Only 1

(b) 1 : 1 mixture of 2 and 3


(c) Only 2

(d) 1 : 1 : 1 mixture of 1, 2, and 3

50. An organic compound *B* is formed by the reaction of ethylmagnesium iodide ($\text{CH}_3\text{CH}_2\text{MgI}$) with a substance *A*, followed by treatment with dilute aqueous acid. Compound *B* does not react with PCC in dichloromethane. Identify *A* ?



51. Which of the following reagents would carry out the following transformation ? ($\text{D} = {}^2\text{H}$)




(a) NaBD_4 in CH_3OH

(b) LiAlH_4 , then D_2O

(c) NaBD_4 in CH_3OD

(d) LiAlD_4 , then D_2O

52. Which sequence of steps describes the best synthesis of 2-methyl-3-pentanone ?







ALCOHOL, ETHERS AND EPOXIDES

341








- (a) (1) 1-Propanol + $(\text{CH}_3)_2\text{CHMgBr}$, diethyl ether
 (2) H_3O^+
 (3) PCC, CH_2Cl_2
- (b) (1) 1-Propanol + $\text{Na}_2\text{Cr}_2\text{O}_7$, H_2SO_4 , H_2O , heat
 (2) SOCl_2
 (3) $(\text{CH}_3)_2\text{CHCl}$, AlCl_3
- (c) (1) 1-Propanol + PCC, CH_2Cl_2
 (2) $(\text{CH}_3)_2\text{CHLi}$, diethyl ether
 (3) H_3O^+
 (4) $\text{Na}_2\text{Cr}_2\text{O}_7$, H_2SO_4 , H_2O , heat
- (d) (1) 2-Propanol + $\text{Na}_2\text{Cr}_2\text{O}_7$, H_2SO_4 , H_2O , heat
 (2) $\text{CH}_3\text{CH}_2\text{CH}_2\text{Li}$, diethyl ether
 (3) H_3O^+
 (4) PCC, CH_2Cl_2

53. Diols (I-IV) which react with CrO_3 in aqueous H_2SO_4 and yield products that readily undergo decarboxylation on heating, are :




- (a) I and II (b) II and III (c) II and IV (d) I and IV

54. Which of the following compounds are not oxidized by HIO_4 ?




- (a) 5, 6, 7 (b) 4, 5, 6, 7 (c) 6, 7 (d) 3, 4, 5, 6, 7




342





ORGANIC Chemistry for IIT-JEE




56. Unknown (A) in the reaction (given in Q. 55) is :



57. In the given table, identify the incorrect option. The digit in box indicate the moles of that substance.

Reactant	HIO_4 consumed	HCO_2H formed	HCHO formed
(a) 	2	1	2
(b) 	3	2	1
(c) 	0	0	0
(d) 	1	1	1


58. Succinic acid $\xrightarrow{\Delta}$ (A) $\xrightarrow[\Delta]{\text{NH}_3}$ (B) $\xrightarrow[\text{KOH}]{\text{Br}_2}$ (C); Product (C) will be :




ALCOHOL, ETHERS AND EPOXIDES

343

- 59A. Given are the structures of cyclic D-glucoside. Moles of HIO_4 consumed with X and Y are respectively:




(X)








(Y)

- (a) 2, 2 (b) 3, 3
(c) 2, 3 (d) 3, 2
- B. Moles of formic acid formed in X and Y respectively are:
(a) 1, 2 (b) 2, 1
(c) 2, 3 (d) 3, 2
- C. Moles of HCHO formed are:
(a) 1, 1 (b) 2, 2
(c) 1, 2 (d) 2, 1
60. In which of the following group each member gives positive iodoform test ?
(a) methanol, ethanol, propanone (b) ethanol, propanal, methanal
(c) ethanol, ethanal, isopropyl alcohol (d) propanal, propanol-2, propanone
61. $\text{H}_2\text{O}^{18} + \text{Na} \xrightarrow{\text{(base)}} (\text{A}) + (\text{B})$







Product (C) of the reactions is:







Product (B) of the reaction is :

344

ORGANIC Chemistry for IIT-JEE








B. Product (C) of above reaction is :







63. Two unknown compounds X and Y, both having molecular formula C_4H_8O , give following results with four chemical tests.

	Bromine	Chromic acid	Lucas reagent
Compound X	decolourises	bubbles	Orange to Green
Compound Y	No reaction	No reaction	No reaction

Compound X and Y respectively are :









Product (N) is :








ALCOHOL, ETHERS AND EPOXIDES**345**





65. Assign the structure of major product (X) of the reaction given below.



Product (A) and (B) respectively in the above reaction are :




Product (A) of above reaction is :



346


ORGANIC Chemistry for IIT-JEE



In above reaction molecular formula of glycerol increases by :



- (a) $\text{C}_4\text{H}_4\text{O}_2$ (b) $\text{C}_6\text{H}_6\text{O}_6$ (c) $\text{C}_6\text{H}_6\text{O}_2$ (d) $\text{C}_6\text{H}_6\text{O}_3$

69. Give the best conditions for this transformation:



- (a) $\text{CH}_3\text{OH}, \text{H}^+$ (cat.), heat (b) $\text{H}_2\text{O}, \text{H}^+$ (cat.), heat
(c) Mg , ether, CH_3OH (d) $\text{SOCl}_2, \text{CH}_3\text{OH}$

70. Give the major organic product of the following reaction.




Product (D) in above reaction is :



ALCOHOL, ETHERS AND EPOXIDES


347

72. Select the best method for the preparation of the following compounds :



(MCPBA = Metachloro per benzoic acid)

- (a) reaction of cyclohexanone with CH_3Li
 (b) reaction of 1-methylcyclohexene with $\text{Hg}(\text{OAc})_2$ followed by NaBH_4
 (c) reaction of cyclohexene with BH_3 ; $\text{NaOH}/\text{H}_2\text{O}_2$, following by CH_3Br
 (d) reaction of cyclohexene with MCPBA, followed by CH_3MgBr
73. Identify the reagents (1-4), required for the transformations shown and arrange them in correct order.



(1) LAH (LiAlH_4)

(2) OsO_4

(3) NaIO_4

(4) NaBH_4


(a) $1 \rightarrow 3 \rightarrow 4 \rightarrow 2$

(b) $2 \rightarrow 3 \rightarrow 1 \rightarrow 4$

(c) $2 \rightarrow 1 \rightarrow 3 \rightarrow 4$


(d) $1 \rightarrow 2 \rightarrow 3 \rightarrow 4$

74. Which describes the best stereochemical aspects of the following reaction ?




- (a) Inversion of configuration occurs at the carbon undergoing substitution.
 (b) Retention of configuration occurs at the carbon undergoing substitution.
 (c) Racemization (loss of configuration) occurs at the carbon undergoing substitution.
 (d) The carbon undergoing substitution is not stereogenic

75. Which of following is an example of Pinacol-Diazotization ?



348

ORGANIC Chemistry for IIT-JEE




76. $(A) \xrightarrow{\text{H}_2\text{O}^+} B + C$; (B) and (C) both give +ve iodoform test. Compound (A) is :





- (a) $\text{CH}_3-\text{CH}=\text{CH}-\text{O}-\text{CH}_2-\text{CH}_3$ (b) $\text{CH}_3-\overset{\text{H}}{\underset{\text{CH}_3}{\text{C}}}-\text{O}-\text{CH}_2-\text{CH}_3$
- (c) $\text{CH}_3-\overset{\text{CH}_2}{\underset{\parallel}{\text{C}}}-\text{O}-\text{CH}_2-\text{CH}_3$ (d) both (b) and (c)

77. A solution of $\text{Ph}_3\text{CCO}_2\text{H}$ in conc. H_2SO_4 gives (X) when poured into methanol X is :





- (a) $\text{Ph}_3\text{C}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{CH}_3$ (b) $\text{Ph}_2\text{CH}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{CH}_3$
- (c) $\text{Ph}_3\text{C}-\text{OCH}_3$ (d) $\text{Ph}_3\text{C}-\text{CH}_3$



Product (B) of the above reaction is :




- (a)  (b)  (c)  (d) 

79. In the given pair of alcohol, in which pair second alcohol is more reactive than first towards hydrogen bromide?

- (a)  and 
- (b)  and 
- (c) $\text{CH}_3-\underset{\text{OH}}{\text{CH}}-\text{CH}_2-\text{CH}_3$ and $\text{CH}_3-\text{CH}_2-\underset{\text{CH}_3}{\text{CH}}-\text{CH}_2-\text{OH}$
- (d) $\text{CH}_3-\underset{\text{OH}}{\text{CH}}-\text{CH}_2-\text{CH}_3$ and $(\text{CH}_3)_2\underset{\text{OH}}{\text{C}}-\text{CH}_2-\text{CH}_3$





ALCOHOL, ETHERS AND EPOXIDES**349**

80. Rank the transition states that occur during the following reaction steps in order of increasing stability (least $\rightarrow$ most stable)







- (a) $1 < 2 < 3$ (b) $2 < 3 < 1$ (c) $1 < 3 < 2$ (d) $2 < 1 < 3$

81. (A), Product (A) is :






82. In which of the following reactions hydrogen gas will not be evolved ?






**83.**

When Grignard reagent reacts with (B) product (D) will be obtained.

Reactant (A) of the above reaction is :



84. (A) $\xrightarrow{\text{LiAlH}_4}$ 2(B) ; structure of (A) is :
(Chiral alcohol only)


**85.**

350

ORGANIC Chemistry for IIT-JEE

Above conversion can be achieved by :


- (a) LiAlH_4 (b) NaBH_4 (c) H_2/Ni (d) CrO_3



(Mandelic acid)

Identify product of above Fischer esterification reaction :


- (a) $\text{Ph}-\text{CH}(\text{O}-\text{Et})-\text{CO}_2\text{H}$ (b) $\text{Ph}-\text{CH}(\text{O})-\text{C}=\text{O}$
 (c) $\text{Ph}-\text{CH}(\text{OH})-\text{CO}_2\text{Et}$ (d) $\text{Ph}-\text{CH}(\text{OH})-\text{C}(=\text{O})-\text{Et}$



Product (C) of the above reaction is :

- (a) $\text{CH}_3-\text{C}(=\text{O})-\text{CH}_2-\text{CH}_2-\text{C}(\text{OH})(\text{CH}_3)-\text{CH}_3$ (b) $\text{CH}_3-\text{C}(\text{OH})(\text{CH}_3)-\text{CH}_2-\text{CH}_2-\text{C}(\text{OH})(\text{CH}_3)-\text{CH}_3$
 (c) $\text{CH}_3-\text{C}(=\text{O})-\text{CH}_2-\text{CH}_2-\text{C}(=\text{O})-\text{CH}_3$ (d) $\text{CH}_3-\text{C}(=\text{O})-\text{CH}_2-\text{C}(=\text{O})-\text{CH}_3$


88. What is the major product of the following reaction ?




- (a) $\text{CH}_3-\text{CH}(\text{OH})-\text{CH}_2-\text{CHO}$ (b) $\text{CH}_3-\text{C}(=\text{O})-\text{CH}_2-\text{C}(=\text{O})-\text{H}$
 (c) $\text{CH}_3-\text{C}(=\text{O})-\text{CH}_2-\text{COOH}$ (d) $\text{CH}_3-\text{CH}(\text{OH})-\text{CH}_2-\text{COOH}$

ALCOHOL, ETHERS AND EPOXIDES**351**





89. The major reason that phenol is a better Bronsted acid than cyclohexanol is that :
- it is a better proton donor.
 - the cyclohexyl group is an electron donating group by induction, which destabilizes the anion formed in the reaction by resonance.
 - phenol is able to stabilize the anion formed in the reaction.
 - the phenyl group is an electron withdrawing group by induction, which stabilizes the anion formed in the reaction.
90. Which of these reagents would accomplish the following reduction ?




- NaBH_4
- LiAlH_4
- 1 mole H_2 , poisoned catalyst, low pressure
- H_3O^+




Identify A and B :

- $A = \text{NaBH}_4$, $B =$  (C) $A = \text{NaBH}_4$, $B =$ 
- $A = \text{LiAlH}_4$, $B =$  (d) $A = \text{LiAlH}_4$, $B =$ 



Product (A) in above reaction is:





- $\text{Ph} - \text{CH}_2 - \underset{\text{OEt}}{\text{CH}} - \text{CH}_3$, (inversion)
- $\text{Ph} - \text{CH}_2 - \underset{\text{OEt}}{\text{CH}} - \text{CH}_3$, (retention)
- $\text{Ph} - \text{CH}_2 - \underset{\text{OEt}}{\text{CH}} - \text{CH}_3$, (racemic)
- $\text{Ph} - \text{CH} = \text{CH} - \text{CH}_3$




- $\text{X} = \text{Ph} - \overset{\text{O}}{\parallel} \text{C} - \text{O}^{18} - \text{CH}_3$ (Trans esterification)

352

ORGANIC Chemistry for IIT-JEE




Fastest rate of reaction will be when R is :




95. Select the correct statement.

- (a) Solvolysis of $(\text{CH}_3)_2\text{C} = \text{CH} - \text{CH}_2 - \text{Cl}$ in ethanol is over 6000 times greater than alkyl chloride (25°C)
 (b) $\text{CH}_3 - \text{CH} = \text{CH} - \text{CH}_2 - \text{OH}$ when reacts with HBr give a mixture of 1-bromo-2-butene and 3-bromo 1-butene
 (c) When solution of 3-buten-2-ol in aqueous sulphuric acid is allowed to stand for one week, it was found to contain 1-butene, 2-butene, 1-butanol and 2-butanol
 (d) All of these



- (a) H^+/Δ , Zn(Hg) , HCl (b) HIO_4 , LiAlH_4
 (c) HIO_4 , H^+/Δ (d) H^+/Δ , HIO_4



Compound (C) is :

- (a) $\text{CH}_3 - \underset{\text{OH}}{\text{CH}} - \text{CH}_2 - \underset{\text{Cl}}{\text{CH}} - \text{CH}_3$
 (b) $\text{HOCH}_2 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{Cl}$
 (c) $\text{HO} - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \underset{\text{Cl}}{\text{CH}} - \text{CH}_3$
 (d) $\text{HO} - \text{CH}_2 - \text{CH}_2 - \underset{\text{Cl}}{\text{CH}} - \text{CH}_2 - \text{CH}_3$


ALCOHOL, ETHERS AND EPOXIDES**353**





98. Iodoform can be obtained on warming NaOH and iodine with :

- (a) $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$ (b) $(\text{CH}_3)_2\text{CH}\overset{\text{O}}{\parallel}\text{C}\text{C}_2\text{H}_5$
 (c) $\text{CH}_3 - \overset{\text{O}}{\parallel}\text{C} - \text{OCH}_3$ (d) $(\text{CH}_3)_2\text{CHCH}_2\text{OH}$

99. Which of these is a reducing agent ?

- (a) CrO_3/H^+ (b) KMnO_4
 (c) LiAlH_4 (d) O_3

100.  $\xrightarrow[\text{(ii). H}_2\text{O}_2/\text{OH}^-]{\text{(i). (BH}_3)_2}$ (P); Product (P) in the reaction is:


- (a)  (b)  (c)  (d) 





101. $\text{CH}_3 - \overset{\text{CH}_3}{\underset{\text{OH}}{\text{C}}} - \text{CH}_3 \xrightarrow[\text{cool}]{\text{Na}_2\text{Cr}_2\text{O}_7}$ (P); Product (P) in the reaction is:

- (a) $\text{CH}_3 - \overset{\text{CH}_3}{\text{C}} = \text{CH}_2$ (b) $\text{CH}_3 - \overset{\text{CH}_3}{\underset{\text{Cl}}{\text{C}}} - \text{Cl}$
 (c) $\text{CH}_3 - \overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}} - \text{O} - \overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}} - \text{CH}_3$ (d) No reaction

102. 1, 2, 3 - butanetriol undergoes oxidative cleavage of HIO_4 . During this process


- (a) 1 equivalent of HIO_4 consumed & HCO_2H & $\text{H}_3\text{C} - \overset{\text{O}}{\parallel}\text{C} - \text{CO}_2\text{H}$ are formed
 (b) 2 equivalents of HIO_4 consumed & HCO_2H , $\text{HCH} = \text{O}$ & $\text{CH}_3 - \text{CH} = \text{O}$ are formed
 (c) 3 equivalents of HIO_4 consumed & HCO_2H (2 eq.) & 1 eq. of $\text{CH}_3\text{CO}_2\text{H}$ are formed
 (d) 2 equivalents of HIO_4 consumed & 2 eq. of HCO_2H & 1 eq. of $\text{CH}_3\text{CH} = \text{O}$ is formed

103.  $\xrightarrow[\text{(ii) H}_2\text{O}]{\text{(i) LiAlH}_4}$ (A); Product (A) of the reaction is : (96%)

- (a)  (b)  (c)  (d) 

354


ORGANIC Chemistry for IIT-JEE



Above reaction is/an example of :



- (a) esterification (b) saponification
(c) *trans*-esterification (d) hydrolysis

105. What is the major organic product of the following sequence of reactions ?




- (a) $(\text{CH}_3)_2\text{CHCH}(\text{OH})\text{CH}_2\text{CH}_3$ (b) $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{OH}$
(c) $(\text{CH}_3)_2\text{CHCH}_2\text{CH}(\text{OH})\text{CH}_3$ (d) $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{CH}_2\text{OH}$



106. The structure of the product formed in the reaction given below is :



- (a) $\text{HCHO}, \text{HCO}_2\text{H}$ (b) $\text{HCHO}, 2\text{HCO}_2\text{H}$
(c) $\text{CO}_2, 2\text{HCO}_2\text{H}$ (d) $\text{CO}_2, \text{HCHO}, \text{HCO}_2\text{H}$



- (a) $4\text{HCO}_2\text{H}, \text{HCHO}$ (b) $4\text{CH}_2\text{O}, \text{HCO}_2\text{H}$
(c) $\text{CO}_2, 4\text{HCHO}$ (d) $\text{CO}_2, 3\text{HCO}_2\text{H}, \text{HCHO}$



ALCOHOL, ETHERS AND EPOXIDES

355






Ratio of moles of formic acid obtained in reaction (i) and reaction (ii) is : $\text{H} + (\text{A})$

(a) $3/4$ (b) $4/5$ (c) $1/2$ (d) $5/4$

(a) $\frac{3}{4}$ (b) $\frac{4}{5}$

$$(c) -1 \text{ H}_2\text{O} \quad \leftarrow \quad (d) \frac{5}{4}$$

110. Which of the following compound gives 2HCHO , CO_2 , HCO_2H when oxidised by periodic acid?



111. Hydration of 3-phenylbut-1-ene in dil. H_2SO_4 will give mainly : (A)

(a) 3-Phenylbutan-1-ol  (b) 3-Phenylbutan-2-ol 
(c) 2-Phenylbutan-2-ol  (d) 2-Phenylbutan-1-ol 

112. Decarboxylation of sodium salicylate with soda lime forms :

(a) Salicylic acid (b) Phenol (c) Benzene (d) None of these

[illegible]

17. In the Libby case, the Supreme Court ruled that the president has the authority to remove executive branch officials without congressional approval.

113. $\text{CH}_3\text{CH}=\text{CH}-\text{O}-\text{CH}_3 \xrightarrow{\text{H}^+/\text{H}_2\text{O}}$ what is the maximum value of (x)?

the CH₂-OH group




[illegible]

(c) Addition of $\text{CH}_3\text{COOCH}_3$ will cause the formation of an equal number of moles of

 $\text{H}_2\text{C}(\text{OH})-\text{CH}_2\text{OH}$ (d) $\text{HO}-\text{CH}_2-\text{CH}_2-\text{O}-\text{CH}_2-\text{CH}_2-\text{OH}$ (e)

(a) $1. \text{H}_2\text{C}=\text{CH}_2$ (b) $2. \text{H}_2\text{C}=\text{CH}_2$ (c) $3. \text{H}_2\text{C}=\text{CH}_2$ (d) $4. \text{H}_2\text{C}=\text{CH}_2$

123. PCC (Pyridinium chlorochromate) is a good reagent for which of the following transformations?



124. How many primary alcohols (including stereoisomers) are possible with formula $C_4H_{10}O$?

(a) two CH_3

(c) (d)

125. (c)  prepared by the reaction of (d) benzaldehyde with the Grignard reagent, $\text{CH}_3\text{CH}_2\text{MgBr}$.

$$\text{CH}_3\text{C}(\text{OH})(\text{H})\text{CH}_2\text{CH}_2\text{CH}_3 \quad (\text{d})$$

(c) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ and $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$

(c) CH_3Br and AlCl_3


358

ORGANIC Chemistry for IIT-JEE

126. 0.092 g of a compound with the molecular formula $C_3H_8O_3$ on reaction with an excess of CH_3MgI gives 67.00 mL of methane at STP. The number of active hydrogen atoms present in a molecule of the compound is :

(a) one (b) two (c) three (d) four

127. Migratory aptitude of the following in decreasing order is :



128. The major product formed in the reaction is :

(a) (A) (b) (B) (c) (C) (d) (D)

117. In the following reaction, the major product is :



118. Ethylmagnesium iodide reacts with an aldehyde to form a secondary alcohol. The aldehyde is :

(a) CH_3CHO (b) CH_3CH_2CHO (c) $CH_3CH_2CH_2CHO$ (d) $CH_3CH_2CH_2CH_2CHO$

129. Reaction of R-2-butanol with p-toluenesulphonyl chloride in pyridine then LiBr gives :

(a) R-2-butyl bromide (b) S-2-butyl tosylate (c) R-2-butyl tosylate (d) S-2-butyl bromide

130. Optically active 2-octanol rapidly loses its optical activity when exposed to :

(a) dilute acid (b) dilute base (c) light (d) humidity

131. If (±) 2-methyl butanoic acid were esterified by reaction with (±) 2-butanol, how many optically active compounds would be present in the final equilibrium reaction mixture ?

(a) 2 (b) 3 (c) 4 (d) 6

124. How many primary alcohols (including stereoisomers) are possible with formula $C_5H_{12}O$?





- 120.

- 132.


Product Z of above reaction is : (a) CH_3COOH and CH_3Cl (b) CH_3COOH and CH_3I (c) CH_3COOH and CH_3Br (d) CH_3COOH and CH_3Cl

ALCOHOL, ETHERS AND EPOXIDES




359



133.







(a) No reaction





134.







Product obtained in the above reaction is .



135.






Product (B) and (C) respectively are :






360

ORGANIC Chemistry for IIT-JEE




- (a) O_3 , Zn, then LiAlH_4 (b) $\text{O}_3/\text{H}_2\text{O}_2$, then LiAlH_4
 (c) cold dil. KMnO_4 , HIO_4 , then LiAlH_4 (d) All of these

138. Which of the following alcohol on treatment with HCl give 3-chloro-3-methyl cyclohexene as a product ?



140. Esterification (shown below) is a reaction converting a carboxylic acid to its ester. It involves only the carbonyl carbon. Esterification of (–)-lactic acid with methanol yields (+)-methyl lactate. Assuming that there are no side reactions, what is true about this reaction ?




- (a) An $\text{S}_{\text{N}}2$ process has occurred, inverting the absolute configuration of the chiral center
 (b) An $\text{S}_{\text{N}}1$ reaction at the chiral center has inverted the optical rotation
 (c) A diastereomer has been produced; diastereomers have different physical properties including optical rotation
 (d) Optical rotation is not directly related to absolute configuration, so the change in sign of rotation is merely a coincidence

ALCOHOL, ETHERS AND EPOXIDES

361

141. Which of the following sets of reagents, used in the order shown, would successfully accomplish the conversion shown ?



- (a) $\text{CH}_3\text{CH}_2\text{CH}_2\text{MgBr}$; H_3O^+ ; PCC, CH_2Cl_2
 (b) $\text{CH}_3\text{CH}_2\text{CH}_2\text{MgBr}$; H_3O^+ ; H_2SO_4 , heat PCC, CH_2Cl_2
 (c) $(\text{C}_6\text{H}_5)_3\text{P}^+-\text{C}^-\text{HCH}_2\text{CH}_3$, B_2H_6 ; $\text{CH}_3\text{CO}_2\text{H}$
 (d) $(\text{C}_6\text{H}_5)_3\text{P}^+-\text{C}^-\text{HCH}_2\text{CH}_3$; H_2O


142. Product ; Product of the reaction is :

- (a) (b) (c) (d)

143. Which of the following compounds on hydrolysis followed by heating gives a product, which gives positive iodoform test ?

- (a) (b) (c) (d)

144. Treatment of a 2° OH with $\text{CrO}_3/\text{H}_2\text{SO}_4$ yields an/a :
 (a) aldehyde (b) carboxylic acid (c) ester (d) ketone
145. Esterification of the acid **P** with the alcohols **Q** will give :




- (a) only one enantiomer (b) a mixture of diastereomers
 (c) a mixture of enantiomers (d) only one diastereomer


362

ORGANIC Chemistry for IIT-JEE


146.



Identify major product of the reaction, when the given compound is hydrolysed and heated strongly:



147.







find out the reactivity order with LiAlH_4 :


(a) $A > B > C > D$ (b) $B > C > D > A$ (c) $D > C > B > A$ (d) $D > B > C > A$

148.

Find out the reaction in which obtained product give positive isocyanide test:



149.



In the above given compound how many functional group reduced by LAH (Lithium aluminium hydride) and SBH (sodium borohydride) respectively?

(a) 4, 4

(b) 4, 3

(c) 3, 4

(d) 4, 2

ALCOHOL, ETHERS AND EPOXIDES**363****ANSWERS — LEVEL 1**

1.	(c)	2.	(a)	3.	(b)	4.	(c)	5.	(b)	6.	(b)	7.	(b)	8.	(a)
9.	(b)	10.	(b)	11.	(b)	12.	(b)	13.	(a)	14.	(a)	15.	(a)	16.	(a)
17.	(d)	18.	(a)	19.	(b)	20.	(a)	21.	(a)	22.	(a)	23.	(d)	24.	(a)
25.	(b)	26.	(b)	27.	(b)	28.	(b)	29.	(c)	30.	(c)	31.	(b)	32.	(b)
33.	(a)	34.	(b)	35.	(b)	36.	(c)	37.	(c)	38.	(c)	39.	(d)	40.	(c)
41.	(b)	42.	(b)	43.	(d)	44.	(d)	45.	(b)	46.	(a)	47.	(b)	48.	(b)
49.	(a)	50.	(b)	51.	(a)	52.	(c)	53.	(c)	54.	(c)	55.	(b)	56.	(a)
57.	(d)	58.	(c)	59.	A-d	59.	B-b	59.	C-a	60.	(c)	61.	(c)	62.	A-b
62.	B-b	63.	(b)	64.	(c)	65.	(b)	66.	(a)	67.	(a)	68.	(d)	69.	(a)
70.	(c)	71.	(b)	72.	(d)	73.	(d)	74.	(c)	75.	(b)	76.	(d)	77.	(c)
78.	(c)	79.	(d)	80.	(c)	81.	(c)	82.	(d)	83.	(a)	84.	(b)	85.	(b)
86.	(c)	87.	(c)	88.	(b)	89.	(d)	90.	(a)	91.	(c)	92.	(b)	93.	(b)
94.	(a)	95.	(d)	96.	(b)	97.	(c)	98.	(a)	99.	(c)	100.	(a)	101.	(d)
102.	(b)	103.	(c)	104.	(c)	105.	(b)	106.	(b)	107.	(d)	108.	(a)	109.	(c)
110.	(d)	111.	(c)	112.	(b)	113.	(b)	114.	(b)	115.	(b)	116.	(d)	117.	(b)
118.	(a)	119.	(b)	120.	(a)	121.	(a)	122.	(b)	123.	(b)	124.	(d)	125.	(a)
126.	(c)	127.	(b)	128.	(c)	129.	(d)	130.	(a)	131.	(c)	132.	(b)	133.	(c)
134.	(b)	135.	(a)	136.	(c)	137.	(d)	138.	(d)	139.	(b)	140.	(d)	141.	(c)
142.	(b)	143.	(b)	144.	(d)	145.	(b)	146.	(c)	147.	(b)	148.	(a)	149.	(b)

364

ORGANIC Chemistry for IIT-JEE



1. Consider the pairs of ethers, numbered I through V, shown below. To the right of each pair is a description of reaction conditions to be applied to each. One compound of the pair will react more rapidly than the other.

Which ether of the two will be more rapidly cleaved?









Write your answer in box.

	(A)	Ether Pairs	(B)	Cleavage Conditions
I.				Treated with HBr in CH_3CN , 40°C
II.				Treated with H_2SO_4 in CH_3CN , 40°C
III.				Treated with H_2SO_4 in CH_3CN , 40°C
IV.				Treated with 5% aqueous H_2SO_4 , 25°C
V.				Treated with 5% aqueous H_2SO_4 , 25°C

ALCOHOL, ETHERS AND EPOXIDES

365

2. Comprehension



Above compounds (a) to (h) are isomers of $\text{C}_5\text{H}_{12}\text{O}$.


Based on the above isomer answer the following (A to F).

- Which isomer is most reactive towards dehydration by conc. H_2SO_4 ?
- Which isomer will undergo rearrangement when treated with conc. H_2SO_4 ?
- Which isomers on dehydration with conc. H_2SO_4 give alkene which is capable to show geometrical isomerism?
- Which isomer is least acidic?
- Which isomers on dehydration give most stable alkene?
- Which isomer on dehydration with conc. H_3PO_4 undergo maximum rearrangement?

366

ORGANIC Chemistry for IIT-JEE

3. Comprehension



[O] = Oxidation

[H] = Reduction

Consider the above sequence and answer A to F.

- A. Conversion $(CH_3 - CH_3 \longrightarrow CH_3 - CH_2 - OH)$ alkane $\longrightarrow$ alcohol is achieved by:
 (a) $Br_2/h\nu$, alc. KOH (b) $Br_2/h\nu$, aq. KOH
 (c) Br_2/CCl_4 , $LiAlH_4$ (d) Br_2/CCl_4 , $NaBH_4$
- B. Conversion $R - CH_2 - OH \longrightarrow R - CHO$ can be done by:
 (a) PCC/CH_2Cl_2 (b) Cu , $300^\circ C$
 (c) CrO_3 (d) All of these
- C. Conversion $R - CHO \longrightarrow R - CO_2H$ can be done by:
 (a) $KMnO_4$ (b) H_2CrO_4
 (c) $K_2Cr_2O_7$ (d) All of these
- D. Conversion $R - CO_2H \longrightarrow R - CHO$ can be done by:
 (a) $LiAlH_4$ (b) $NaBH_4$
 (c) DIBAL - H (d) All of these
- E. Conversion $R - CHO \longrightarrow R - CH_2 - OH$ can be done by:
 (a) $LiAlH_4$ (b) $NaBH_4$
 (c) H_2/Ni (d) All of these
- F. Reduction $R - CH_2 - OH \longrightarrow R - CH_3$ can be done by:
 (a) $LiAlH_4$ (b) $NaBH_4 - AlCl_3$
 (c) $H_2 - Ni$ (d) Red P + HI

4. Which of the following is true for 3-methylbutanal ?

a.	This compound may be classified as an aldehyde.	
b.	This compound may be classified as a ketone	
c.	An aldol reaction takes place on treatment with NaOH solution.	
d.	There is no reaction with $LiAlH_4$ in ether solution.	
e.	An excess of CH_3MgBr in ether reacts to give 4-methyl-2-pentanol.	
f.	Wolff-Kishner reduction gives butane.	
g.	This compound is an isomer of 3-pentanone.	

ALCOHOL, ETHERS AND EPOXIDES**367**

5. This problem is an introduction to the planning of multistep syntheses.

For use, you have six reactant compounds (**A** through **F**) ; and eight reagents (**1** through **8**), shown below.


Following these lists, five multistep syntheses are outlined. For each of these, certain reactants or reagents must be identified by writing an appropriate letter or number in designated answer boxes. Write a single letter or number, indicating your choice of the best reactant or reagent, in each answer box.

Reactant Compounds :

(A)




(B)




(C)



(D)








(E)



(F)

Reagents :(1) Jones' reagent [$\text{Na}_2\text{Cr}_2\text{O}_7$ in H_3O^+](2) PCC [CrO_3 in pyridine + HCl](3) Sodium hydride NaH (4) Sodium borohydride NaBH_4 (5) Thionyl chloride SOCl_2 (6) Phosphorus tribromide PBr_3 (7) Aluminium trichloride AlCl_3

(8) Magnesium turnings in ether



368

ORGANIC Chemistry for IIT-JEE


6. Which of the following is true for 3-methyl-2-butanone?

a.	It may be prepared by CrO_3 oxidation of 2-methyl-2-butanol.	
b.	Its reaction with NaBH_4 gives a secondary alcohol.	
c.	It may be prepared by acidic Hg^{2+} catalyzed hydration of 3-methyl-1-butyne.	
d.	It forms a silver mirror on treatment with $[\text{Ag}(\text{NH}_3)_2]^+$.	
e.	This compound is an isomer of 4-penten-1-ol.	

7. Which of these methods would serve to prepare 1-phenyl-2-propanol?

a.	Addition of benzyl Grignard reagent to acetaldehyde (ethanal).	
b.	Addition of phenyl lithium to propylene oxide (methyloxirane).	
c.	Addition of phenyl Grignard reagent to acetone (2-propanone).	
d.	Acid-catalyzed hydration (addition of water to) of 2-phenyl-1-propene.	
e.	Addition of methyl Grignard reagent to acetophenone (methyl phenyl ketone).	
f.	Addition of methyl Grignard reagent to phenylacetaldehyde.	

8. Match the Column (I) and (II).

Column (I)		Column (II)	
Reaction		Name of Reaction	
(a)		(p)	Pinacol-Pinacolone rearrangement

ALCOHOL, ETHERS AND EPOXIDES

369

(b)	$\text{C(CH}_3)_2\text{CH}_2\text{OH} \xrightarrow[2\text{HCl}]{\text{NaNO}_2}$	(q)	Semi-Pinacol reaction
(c)	$\text{Propylene oxide} \xrightarrow[\Delta]{\text{H}^+}$	(r)	Pinacolic-Diazotization
(d)	$\text{CH}_3\text{CH(OH)CH(OH)CH}_3 \xrightarrow[(2) \text{Et}_3\text{N}, \Delta]{(1) \text{TsCl}}$	(s)	Pinacol fashion reaction


9. Match the Column (I) and (II).

Column (I)		Column (II)	
Reactant		Products	
(a)	$\text{1-methylcyclohexanol} \xrightarrow[\text{H}_2\text{SO}_4 \text{ Conc.}]{\text{CH}_3\text{OH}}$	(p)	
(b)	$\text{1-methylcyclohexanol} \xrightarrow[(2) \text{CH}_3\text{I}]{(1) \text{NaH}}$	(q)	
(c)	$\text{1-methylcyclohexanol} \xrightarrow[(3) \text{CH}_3\text{I}]{(1) \text{HBr}, (2) \text{Mg}}$	(r)	
(d)	$\text{1-methylcyclohexanol} \xrightarrow[(2) {}^{14}\text{CH}_3\text{I}]{(1) \text{Na}}$	(s)	

370

ORGANIC Chemistry for IIT-JEE


10.



Ratio of moles of formaldehyde obtained in the reaction (1) and reaction (2) ?

11. Comprehension

Di-tert-glycols react with the presence of acid to give tertiary ketones. The trivial name of the simplest glycol of this type is pinacol, and this type of reaction therefore is named pinacol rearrangement (in this specific case, the reaction is called a pinacol-pinacolone rearrangement). The rearrangement involves 4 steps. one of the hydroxyl groups is protonated in the first step. A molecule of water is eliminated in the second step and a tertiary carbocation is formed. The carbocation rearranges in the third step into a more stable carboxonium ion via a [1, 2] rearrangement. In the last step, the carboxonium ion is deprotonated and the product ketone is obtained.




A. What is R.D.S. of pinacol-pinacolone rearrangement ?

- (a) I step (b) II step
(c) III step (d) IV step


ALCOHOL, ETHERS AND EPOXIDES

371





How many products obtained in above reaction ?


- (a) 1 (b) 2
(c) 3 (d) 4



Product 'P' is :



Product (A) is :







372

ORGANIC Chemistry for IIT-JEE

ANSWERS — LEVEL 2

- I – B; II – A; III – A; IV – B; V – B
- A – d; B – a, c, e, f, g, h; C – a, b, c; D – d; E – d, e, f, g, h; F – e
- A – b; B – d, C – d, D – c, E – d; F – d
- a, c, e, g



- b, c, e
- a – p, b – r, c – s, d – q
- Ratio of reaction I and II = 2
- a, b, f
- a – s, b – r, c – q, d – p
- A – b; B – b; C – c; D – a








7

ALDEHYDES AND KETONES




ATDB.uno




374

ORGANIC Chemistry for IIT-JEE




Above conversion can be achieved by :





- (a) Wolff-Kishner reduction (b) Clemmensen reduction
(c) LiAlH_4 (d) NaBH_4




Above conversion can be achieved by :

- (a) Wolff-Kishner reduction
(b) Clemmensen reduction
(c) $\text{HS}-\text{CH}_2-\text{CH}_2-\text{SH}$, following by Raney Ni
(d) None of these



- (a)  (b)  (c)  (d) 




Above conversion can be achieved by :

- (a) Wolff-Kishner reduction (b) Clemmensen reduction
(c) LiAlH_4 (d) NaBH_4



ALDEHYDES AND KETONES

375






Identify product (A) & (B) from the given product P, Q, R:

- (a) $A = P, B = Q$ (b) $A = Q, B = R$ (c) $A = Q, B = P$ (d) $A = R, B = P$



376


ORGANIC Chemistry for IIT-JEE




Above conversion can be carried out by :





- (a) Clemmensen reduction (b) Wolff-Kishner reduction
(c) LiAlH_4 (d) NaBH_4

10. Increasing order of equilibrium constants for the formation of a hydrate:




- (a) $\text{IV} < \text{III} < \text{II} < \text{I}$ (b) $\text{IV} < \text{III} < \text{I} < \text{II}$
(c) $\text{I} < \text{II} < \text{III} < \text{IV}$ (d) $\text{II} < \text{III} < \text{I} < \text{IV}$





















- (a)  (b) 
(c)  (d) 

ALDEHYDES AND KETONES

377




Predict the product of hydrolysis of the above molecule.




378

ORGANIC Chemistry for IIT-JEE





Reagents to carry out above conversion, P, Q, R respectively are :

- (a) $\text{H}_2\text{C}=\text{CH}-\text{CH}_2-\text{Br}$, $(\text{HO}^\ominus)$, $[\text{HO}^\ominus, \Delta]$, Wacker-process
 (b) $\text{H}_2\text{C}=\text{CH}-\text{CH}_2-\text{Br}$, $(\text{HO}^\ominus)$, Wacker-process, $\text{HO}^\ominus, \Delta$
 (c) Wacker process, $\text{H}_2\text{C}=\text{CH}-\text{CH}_2-\text{Br}$, $(\text{HO}^\ominus)$, $\text{HO}^\ominus (\Delta)$
 (d) Wacker process, $\text{HO}^\ominus (\Delta)$, $\text{H}_2\text{C}=\text{CH}-\text{CH}_2-\text{Br}$, $(\text{HO}^\ominus)$




Above reaction is a Baeyer Villiger rearrangement of an asymmetric ketone with magnesium mono peroxo phthalate hexahydrate (in the drawing, Mg^{+2} is omitted for clarity) Identify major product.




Above compounds can be differentiated by following reagent:

- (a) 2-4 DNP (Brady reagent) (b) Tollen's reagent
 (c) Lucas reagent (d) NaHSO_3




Product (D) will be :
















ALDEHYDES AND KETONES

379







Relationship between products (Q) and (S) is:

- (a) Positional isomer (b) Chain isomer
(c) Stereoisomer (d) Functional isomer




Above conversion can be achieved by:




380

ORGANIC Chemistry for IIT-JEE


24. Which sequence represents the best synthesis of hexanal ?



- (a) 1. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Br} + \text{NaC}\equiv\text{CH}$
2. H_2O , H_2SO_4 , HgSO_4
- (b) 1. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2 + \text{CH}_3\text{C}(=\text{O})\text{OOH}$
2. CH_3MgBr , diethyl ether
3. H_3O^+
4. PCC , CH_2Cl_2
- (c) 1. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{C}(=\text{O})\text{CH}_3$
2. $\text{CH}_3\text{C}(=\text{O})\text{OOH}$
3. LiAlH_4
4. H_2O
5. PCC , CH_2Cl_2
- (d) 1. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{MgBr} + \text{H}_2\text{C}(\text{O})\text{CH}_3$
2. H_3O^+
3. PCC , CH_2Cl_2

25.  $\xrightarrow{\text{HNO}_3}$ (A) $\xrightarrow{\text{Ca(OH)}_2}$ (B), Product (B) in this reaction is :

26.




Compound (x) in the above reaction is :











- (a) $\text{Ph} - \text{C}(=\text{O}) - \text{CH}_3$ (b) $\text{Ph} - \text{C}(=\text{O}) - \text{H}$
- (c) $\text{Ph} - \text{CH}_2 - \text{C}(=\text{O}) - \text{H}$ (d) $\text{Ph} - \text{CH}_2 - \text{C}(=\text{O}) - \text{CH}_3$

ALDEHYDES AND KETONES






381





Product (C) of the above reaction is :



Product (A) of the above reaction is (bromination occur not in the benzene ring) :









Compound (A) is :








382

ORGANIC Chemistry for IIT-JEE






(d) both (b) and (c)



ALDEHYDES AND KETONES**383**

33. Correct order of reactivity of following compounds towards Grignard reagent?









(a) I > II > III

(b) II > I > III

(c) II > III > I

(d) I > III > II



Total number of products obtained in above reaction is :

(a) 2

(b) 3


(c) 4

(d) 5

384



ORGANIC Chemistry for IIT-JEE

36. What reagent and/or reaction conditions would you choose to bring about the following conversion?





- (a) 1. LiAlH_4 , 2. H_2O (b) H_2O , H_2SO_4 , heat
(c) H_2O , NaOH , heat (d) PCC , CH_2Cl_2

37. $\xrightarrow[\text{KH excess}]{\text{MeI excess}}$ 81% yield ; Product of the reaction is :



The above reduction can be best carried out by :

- (a) Clemmensen reduction (b) Wolff-Kishner reduction
(c) NaBH_4 (d) None of these




Product (A) and (B) is differentiated by:


- (a) 2,4-DNP (b) NaOI (c) Na-metal (d) NaHSO_3

ALDEHYDES AND KETONES


385



End product (C) in above reaction is :




41. Compound (X) $\text{C}_4\text{H}_8\text{O}$, which reacts with 2, 4-DNP derivative and gives negative haloform test is :




42. When a nucleophile encounters a ketone, the site of attack is :

- (a) the carbon atom of the carbonyl
 (b) the oxygen atom of the carbonyl
 (c) both the carbon and oxygen atoms with equal probability
 (d) no attack occurs as ketones do not react with nucleophiles


43. The correct order of rate of reaction toward nucleophilic addition reaction:



44. The structure  would be best classified as a(an) :


- (a) Acetal (b) Hemiacetal (c) Hydrate (d) Cyanohydrin

45. Which of the following pairs of reactants is most effective in forming an enamine ?







386

ORGANIC Chemistry for IIT-JEE







(d) None of these form an enamine.




46. The reaction of $\text{C}_6\text{H}_5\text{CH}=\text{CHCHO}$ with LiAlH_4 gives :



47. Product (B) of the reaction is :








48. Which of following compound is hemiacetal ?







(d) all of these



49. $\text{Ph}-\text{CH}_2-\text{C}\equiv\text{N} \xrightarrow[\text{THF}]{\text{LDA}} \xrightarrow{\text{CH}_3\text{I}} 71\%$; End product of the reaction will be :



Above conversion can be achieved by :



51. Product of the reaction is/are :




(d) Both (a) and (c)


ALDEHYDES AND KETONES

387




52. Product (A) of the reaction is:



53. Product (C) of the reaction is:




54. Product (A) of the reaction will be:



55. $\text{R}-\text{CHO} + \text{R}'-\text{NH}_2 \xrightarrow{\text{R}-\text{NH}_2} \text{R}-\text{CH}=\text{N}-\text{R}'$. This reaction gives best yield at pH:

- (a) pH 1-2 (b) pH 4-5 (c) pH 10-11 (d) pH 13-14


56. An aromatic compound A of the molecular formula $\text{C}_8\text{H}_{10}\text{O}$ on reaction with iodine and dilute NaOH gives a yellow precipitate. The structure of the compound is expected to be:



390

ORGANIC Chemistry for IIT-JEE



69. Consider the following sequence of reactions.



The ketone (A) is :



70. In the reaction,





71. The conversion of acetophenone into benzoic acid can be achieved by its reaction with :

- (a) sodium hydroxide followed by acidification
 (b) iodine and sodium hydroxide, followed by acidification
 (c) hydroxylamine followed by reaction with H_2SO_4
 (d) *m*-chloroperoxobenzoic acid

72. In which of the following compounds the methylenic hydrogens are the most acidic ?

- (a) $\text{CH}_3\text{COCH}_2\text{CH}_3$ (b) $\text{CH}_3\text{CH}_2\text{COOC}_2\text{H}_5$
 (c) $\text{CH}_3\text{CH}_2\text{CH}(\text{COOC}_2\text{H}_5)_2$ (d) $\text{CH}_3\text{COCH}_2\text{CN}$





73. Which is the major product of the following reaction ?



Product (B) is :

ALDEHYDES AND KETONES

391





75. The presence of unsaturation in organic compounds can be tested with :

(a) Schiff's reagent (b) Tollens' reagent (c) Fehling's reagent (d) Baeyer's reagent


76. Which of the following gives iodoform test ?

(a) $\text{CH}_3\text{CH}_2\text{OH}$ (b) $\text{C}_2\text{H}_5\text{CHO}$ (c) $(\text{CH}_2\text{OH})_2$ (d) None of these


77. Which of the following β -keto carboxylic acid does not undergo decarboxylation on heating ?







(d) None of these



ATDB.uno




Product (D) is :




392

ORGANIC Chemistry for IIT-JEE



79. The compound shown in the below undergoes racemization on reaction with aqueous acid.





Which of the following structures best represents the intermediate responsible for this process?



80. The final product of the following sequence of reaction is :







81. The amino ketone shown below undergoes a spontaneous cyclization on standing. What is the major product of this intramolecular reaction ?








ALDEHYDES AND KETONES

393







82. Compound (A) $C_6H_{12}O$ is optically active. Compound (A) give negative Tollens test and positive test with 2-4-di-nitro phenyl hydrazine. Identify A.







83.



84.




Product (B) of above the reaction is :



394

ORGANIC Chemistry for IIT-JEE









(a) Diastereomer






(b) Racemic

(c) Meso





(d) Optically pure enantiomer



Product (A) and (C) are :




Product (B) is :




ALDEHYDES AND KETONES





395





Product (P) is :

- (a) Hemiacetal (b) Acetal (c) Alcohol (d) Alkane



- (a)  (b) 
(c)  (d) 



Reactant	$K_{eq.}$
PhCHO	a
	b
Ph-C(=O)-CH ₃	c
CH ₃ -C(=O)-H	d


The correct order of decreasing value of $K_{eq.}$ is :

- (a) $a > b > c > d$ (b) $d > a > b > c$
(c) $d > b > a > c$ (d) $d > a > c > d$


396

ORGANIC Chemistry for IIT-JEE

93. Product (B) of the given reaction is :



94. End product (C) of the reaction is :







95. (A) $\xrightarrow{\text{O}_3}$ does not undergo self aldol condensation
 $\text{C}_{11}\text{H}_{8}\text{O} \xrightarrow{\text{O}_3} \text{Ph}-\text{CHO} + 2\text{b} \xrightarrow{\text{Ag}^+} \text{oxalic acid}$

Compound (A) will be :












(a) $\text{Ph}-\text{C}\equiv\text{C}-\text{C}\equiv\text{C}-\text{CHO}$ (b) $\text{Ph}-\text{C}\equiv\text{C}-\text{CH}=\text{CH}-\text{CHO}$ (c) $\text{Ph}-\text{CH}=\text{CH}-\text{C}\equiv\text{C}-\text{CHO}$ (d) $\text{Ph}-\text{CH}=\text{CH}-\text{C}\equiv\text{CH}-\text{CO}_2\text{H}$

96. Product ; Product of the reaction is :



ALDEHYDES AND KETONES**397**

97. Which pair of reactants compounds may be used to make given acetal ?



(A) and (B) is differentiated by :

(a) NaH

(b) 2,4-DNA



(c) Tollen's reagent

(d) NaHSO₃



398

ORGANIC Chemistry for IIT-JEE


- 100.** Which of the following pairs cannot be differentiated by Tollens' reagent ?
 (a) Benzaldehyde and benzyl alcohol (b) Hexanal and 2-hexanone
 (c) 2-Hexanol and 2-hexanone (d) Pentanal and diethyl ether
- 101.** An optically active compound $C_6H_{12}O$ gives positive test with 2, 4-dinitrophenyl hydrazine, but negative with Tollens' reagent, what is the structure of the compound ?



- 102.** Compound (A) $C_6H_{12}O_3$, when treated with I_2 in aqueous sodium hydroxide gives yellow precipitate. When A is treated with Tollens reagent no reaction occur. When A is hydrolysed and then treated with Tollens reagent, a silver mirror is formed in test tube. Compound (A) will be :







- 103.** $\begin{array}{c} CH_2 - CH_2 - CO_2H \\ | \\ CH_2 - CH_2 - CO_2H \end{array} \xrightarrow[BaCO_3]{\Delta} A \xrightarrow{NH_2-NH_2} B \xrightarrow[KOH]{heat} (C)$, Product (C) obtained is :








ALDEHYDES AND KETONES


399





104. Which of following does not react with NaHSO_3 (sodium bisulphite)?








105.  $\xrightarrow[\text{Raney Ni}]{\text{H}_2}$ (A) ; Product (A) is :



106.  + $\text{CH}_2 = \text{CH} - \text{Br} \xrightarrow{\text{KO}} (A)$; Product (A) is : (75%)




107.  $\xrightarrow[(2) \text{HCl/H}_2\text{O}]{(1) \text{Me-Li(excess)}} (A) \xrightarrow[\text{NaOH}]{\text{I}_2} (B) + \text{CHI}_3$; Product (B) in this reaction is :



400

ORGANIC Chemistry for IIT-JEE


108.



Arrange the following reagent in the correct order in which above transformation is carried out :

- (a) $\text{KOD}/\text{D}_2\text{O}, \text{H}^+/\Delta, \text{LiAlH}_4$ (b) $\text{H}^+/\Delta, \text{KOD}/\text{D}_2\text{O}, \text{LiAlH}_4$
 (c) $\text{KOD}/\text{D}_2\text{O}, \text{LiAlH}_4, \text{H}^+/\Delta$ (d) $\text{LiAlH}_4, \text{H}^+/\Delta, \text{KOD}/\text{D}_2\text{O}$


109.



Compound (C) can show geometrical isomerism. Product (E) of the reaction will be :


- (a) $\text{CH}_3 - \text{C}(=\text{O}) - \text{CH}_3$ (b) $\text{CH}_3 - \text{CH}_2 - \text{C}(=\text{O}) - \text{H}$
 (c) $\text{CH}_3 - \text{CHO}$ (d) HCHO

110. Arrange in their increasing order of equilibrium constants for hydration ?



- (a) $A < B < C < D < E$ (b) $A < C < B < E < D$
 (c) $A < C < E < B < D$ (d) $C < A < B < E < D$







111. End products of the following sequence of reactions are :



- (a) yellow ppt. of CHI_3 , cyclohexanecarboxylic acid (b) yellow ppt. of CHI_3 , cyclohexanecarbaldehyde
 (c) yellow ppt. of CHI_3 , cyclohexanone (d) yellow ppt. of CHI_3 , cyclohexane-1,2-dicarboxylic acid

ALDEHYDES AND KETONES

401




(a) Racemic mixture









(b) Diastereomers

(c) Meso

(d) Mixture of meso compound and optically active compound




$\therefore$ reactant (A) is :



402

ORGANIC Chemistry for IIT-JEE


116.



- (a) & (b)

 (c) & (d)


117.



- (a) & (b)

 (c) & (d)







118.



- (a) (b) (c) (d)

ALDEHYDES AND KETONES

403




Which of following can be used as a catalyst in the above reaction?

- (a) Cl^- (b) $\text{CH}_3-\overset{\overset{\text{O}}{\parallel}}{\text{C}}-\text{O}^-$ (c) $\text{Et}-\text{O}^-$ (d) HSO_4^-

121. Arrange the following carbonyl compounds in decreasing order of their reactivity in nucleophilic addition reaction.



(i)



(ii)




(iii)





(iv)

- (a) ii > iii > i > iv (b) ii > i > iv > iii
(c) iii > ii > i > iv (d) iii > i > iv > ii

122. The following reaction were carried out.






The final product formed in the above reaction sequence is :



404

ORGANIC Chemistry for IIT-JEE



A. Yield of each step as actually carried out in the laboratory is given above. What is overall yield of reaction?

(a) 42%

(b) 31%

(c) 21%

(d) 60%


B. What is the appropriate reagent to carry out above synthesis, i.e., A, B, C respectively are :

(a) Br_2/H^+ , LiAlH_4 , H^+

(b) Br_2/H^+ , NaPH_4 , HO^-

(c) NBS, AlCl_3 , HO^-

(d) P_2/HO^- , P_2 , HO^-



Product (E) is :




(a) Nylon 66

(b) Nylon 6

(c) Styrene

(d) Polystyrene






125. Methyl vinyl ketone on reaction with LiCuMe_2 gives a major product, whose structure is :







ALDEHYDES AND KETONES

405




126. Which of following is in capable to show iodoform test ?



Product (B) of above reaction is :







128. $\xrightarrow[\Delta]{\text{H}_3\text{O}^+} \text{A}$; Product obtained is :



(d) None of these


129. (A) $\xrightarrow{\text{LiAlH}_4} (\text{B}) \xrightarrow[\Delta]{\text{H}^+} \text{Diastereomers}$
Symmetrical Ketone

Reactant (A) is :




406

ORGANIC Chemistry for IIT-JEE


Value of x in above reaction is :

- (a) 1 (b) 2 (c) 3 (d) 4



Molecular weight of compound (A) is :


- (a) 58 (b) 120
(c) 60 (d) 182




Product (A) of above reaction will be

- (a) $\text{Ph} - \overset{\text{Ph}}{\underset{|}{\text{C}}} = \text{CH} - \text{O}$ (b) $\text{Ph}_2\text{CH} - \overset{\text{H}}{\underset{|}{\text{CH}_2}}$
- (c) $\text{Ph}_2\text{CH} - \overset{\text{OH}}{\underset{|}{\text{CH}}} - \text{OH}$ (d) $\text{Ph}_2\text{CH} - \overset{\text{O}}{\parallel} \text{C} - \text{CH}_3$

133. Which of the following will form stable hydrate ?

- (a) CCl_3CHO (Chloral) (b)  (Ninhydrin)
- (c) $(\text{CF}_3)_2\text{CO}$ (d) All of these

134. The pH at which maximum hydrate is present in an solution of oxaloacetic acid:




- (a) pH = 0 (b) pH = 12
(c) pH = 4 (d) pH = 6

ALDEHYDES AND KETONES

407

135. Arrange their stabilities of given gem- diols in decreasing order.







(a) I > II > III

(b) III > II > I

(c) I > III > II

(d) III > I > II

136. Maximum hydration takes place of :




137. The conversion, $\text{PhCN} \rightarrow \text{PhCOCH}_3$, can be achieved most conveniently by reaction with:

(a) CH_3MgBr followed by hydrolysis






(b) $\text{I}_2 - \text{NaOH}$, CH_3I

(c) dil. H_2SO_4 followed by reaction with CH_2N_2





(d) LAH followed by reaction with CH_3I



In the above reaction, product (B) is:




Structure of A is :







408


ORGANIC Chemistry for IIT-JEE








Product (B) is :




141.







To carry out above conversion, arrange the following reagents in correct order.



142.




Product (B) is:





ALDEHYDES AND KETONES

409




Identify appropriate reagents for the above reaction:


- (a) $a = \text{Br}_2/\text{CCl}_4$, $b = \text{aq. KOH}$
 (b) $a = \text{Br}_2/\text{H}^+$, $b = \text{aq. KOH}$
 (c) $a = \text{Br}_2/\text{H}^+$, $b = \text{alc. KOH}$
 (d) $a = \text{Br}_2/\text{HO}^-$, $b = \text{aq. KOH}$




145. The K_{eq} values in HCN addition to following aldehydes are in the order :



(I)





(II)



(III)

- (a) $\text{I} > \text{II} > \text{III}$ (b) $\text{II} > \text{III} > \text{I}$ (c) $\text{III} > \text{I} > \text{II}$ (d) $\text{II} > \text{I} > \text{III}$



relation between K_1 and K_2 is :

- (a) $K_1 = K_2$ (b) $K_1 > K_2$ (c) $K_2 > K_1$ (d) $K_1 = K_2 = 1$

410

ORGANIC Chemistry for IIT-JEE

ANSWERS — LEVEL 1

1.	(a)	2.	(b)	3.	(d)	4.	(c)	5.	(a)	6.	(c)	7.	(b)	8.	(b)
9.	(b)	10.	(c)	11.	(c)	12.	(c)	13.	(b)	14.	(b)	15.	(b)	16.	(b)
17.	(b)	18.	(b)	19.	(b)	20.	(d)	21.	(b)	22.	(b)	23.	(b)	24.	(d)
25.	(c)	26.	(b)	27.	(b)	28.	(a)	29.	(b)	30.	(c)	31.	(d)	32.	(c)
33.	(b)	34.	(b)	35.	(a)	36.	(b)	37.	(c)	38.	(d)	39.	(b)	40.	(a)
41.	(b)	42.	(a)	43.	(a)	44.	(b)	45.	(c)	46.	(c)	47.	(b)	48.	(d)
49.	(c)	50.	(b)	51.	(d)	52.	(b)	53.	(b)	54.	(c)	55.	(b)	56.	(b)
57.	(a)	58.	(d)	59.	(c)	60.	(c)	61.	(c)	62.	(a)	63.	(b)	64.	(b)
65.	(b)	66.	(d)	67.	(b)	68.	(d)	69.	(b)	70.	(b)	71.	(b)	72.	(d)
73.	(d)	74.	(a)	75.	(d)	76.	(a)	77.	(a)	78.	(c)	79.	(c)	80.	(c)
81.	(d)	82.	(b)	83.	(b)	84.	(b)	85.	(b)	86.	(a)	87.	(b)	88.	(c)
89.	(b)	90.	(b)	91.	(c)	92.	(b)	93.	(b)	94.	(b)	95.	(c)	96.	(c)
97.	(d)	98.	(b)	99.	(c)	100.	(c)	101.	(c)	102.	(c)	103.	(c)	104.	(c)
105.	(c)	106.	(b)	107.	(c)	108.	(c)	109.	(c)	110.	(b)	111.	(c)	112.	(b)
113.	(b)	114.	(c)	115.	(c)	116.	(a)	117.	(b)	118.	(a)	119.	(b)	120.	(c)
121.	(b)	122.	(b)	123.	A-c	123.	B-b	124.	(b)	125.	(a)	126.	(c)	127.	(a)
128.	(b)	129.	(c)	130.	(a)	131.	(a)	132.	(c)	133.	(d)	134.	(a)	135.	(a)
136.	(a)	137.	(a)	138.	(b)	139.	(d)	140.	(b)	141.	(d)	142.	(a)	143.	(c)
144.	(b)	145.	(d)	146.	(b)										










ALDEHYDES AND KETONES

411




1. Select the best choice for example (A to L) from the examples (a to n) given below. Write your choice in the box given.

A.	An acetal derivative of a ketone.	
B.	A chiral ketone.	
C.	An aldehyde that gives a aldol condensation with itself.	
D.	An oxime derivative	
E.	A reagent that reduces aldehydes to 1° alcohols.	
F.	An α , β -unsaturated ketone.	
G.	A reagent that oxidizes aldehydes to carboxylic acids.	
H.	A reagent that reduces ketones to alkanes.	
I.	An enamine derivative of a ketone.	
J.	An intermediate in imine formation.	
K.	A cyclic hemiacetal.	
L.	A cyanohydrin derivative.	

(a)		(b)		(c)	
(d)		(e)		(f)	$\text{Zn(Hg)H}_3\text{O}^{(+)}$
(g)		(h)	NaBH_4 aq. alcohol	(i)	
(j)	$\text{Ag(NH}_3)_2^{(+)}\text{OH}^{(-)}$	(k)		(l)	

412









ORGANIC Chemistry for IIT-JEE

(m)		(n)	$\text{CH}_3-\text{CH}_2-\text{C}(=\text{O})\text{H}$




2. The following questions refer to the compounds (A to G) shown below :

i.	Which compounds are reduced by sodium borohydride ?	ii.	Which compounds are hydrolyzed by hot aqueous acid ?	iii.	Which compound are oxidized by $\text{CrO}_3/\text{pyridine}$?






A		E		A		E	
B		F		B		F	
C		G		C		G	
D		H		D		H	

A.		B.		C.		D.	
E.		F.		G.		H.	

3. Match the column:


Column (I)		Column (II)	
(a)		(p)	racemic mixture
(b)		(q)	Diastereomers
(c)	$\text{Ph}-\text{CH}_2-\text{Cl} \xrightarrow{\text{KCN}}$	(r)	Nu-addition reaction
(d)		(s)	Nu-Substitutions reaction

ALDEHYDES AND KETONES**413****4. Complete the following table.**

	REACTANT	REAGENT(S)/ CONDITIONS	MAJOR ORGANIC PRODUCTS
a.		$\text{H}_2/\text{Pd} - \text{C}$ in ethanol (solvent)	A
b.		$\text{H}^+/\text{H}_2\text{O}/\Delta$	B
c.		$(\text{CH}_3)_2\text{C}^- - \text{P}^+(\text{C}_6\text{H}_5)_3$	C
d.		1. $\text{Li}^+[(\text{CH}_3)_2\text{Cu}]^-$ in dry ether 2. $\text{H}^+/\text{H}_2\text{O}$	D
e.	E	$\text{OH}^-/\text{ethanol}/\Delta$	

5. Comprehension

Consider the following reactions and answer A and B.



A. Suggest a reagent appropriate step (a) the synthesis.

(a) HO^-/Br_2 (1 mole)

(b) H^+/Br_2 (1 mole)

(c) both (a) and (b)

(d) None of these

414

ORGANIC Chemistry for IIT-JEE

B. Yield of each step as actually carried out in laboratory is given above each arrow. What is overall yield of the reaction ?




(a) 60%

(b) 21%

(c) 40%











(d) 68%

6.

Reaction 1.	
Reaction 2.	
Reaction 3.	

Degree of unsaturation present in compound (A + B + C) is ?

7. Within each set, which compound would be more reactive toward carbonyl addition reaction ?



	A	B
Set (1)		
Set (2)		
Set (3)		
Set (4)		
Set (5)		

ALDEHYDES AND KETONES

415


Set (6)		
Set (7)		
Set (8)		
Set (9)		
Set (10)		

8. Match the Column (I) and Column (II). (Matrix)

Column (I)			Column (II)	
(A)	 $\xrightarrow[\text{traces of KOH}]{\text{HCN}}$ (A) $\xrightarrow{\text{LiAlH}_4}$ (B) $\xrightarrow[\text{HCl}]{\text{NaNO}_2}$ (C)	(p)	Formation of six member ring takes place	
(B)	 $\xrightarrow{\text{NH}_2\text{OH}}$ (A) $\xrightarrow{\text{H}^+}$ (B) $\xrightarrow{\text{LAH}}$ (C)	(q)	Final product is Ketone	

416

ORGANIC Chemistry for IIT-JEE

(C)	$\text{CH}_3 - \overset{\text{O}}{\parallel} \text{C} - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \overset{\text{O}}{\parallel} \text{C} - \text{H} \xrightarrow[\Delta]{\text{HO}^-} (\text{A})$	(r)	Final product formed will give positive Tollens test
(D)	 $\xrightarrow[\Delta]{\text{H}^+} (\text{A})$	(s)	Final product formed will react with 2,4-DNP. (2,4-di-nitrophenyl hydrazine)

9. Consider reactions A through F. Those carbon atoms undergoing change, as part of a functional group, are marked as C¹², C¹⁴ or starred. In the cases shown, each carbon atom has either been reduced or oxidized. Your job is to identify the change in oxidation state that has occurred for each of the marked carbon.









Reaction		C ¹²	C ¹⁴
A.	$\text{CH}_3\text{CH}=\text{CH}_2 \xrightarrow{\text{Br}_2} \text{CH}_3\text{CHBrCH}_2\text{Br}$	Reduced	Reduced
		Oxidized	Oxidized
B.	$\text{CH}_3\text{CH}=\text{CH}_2 \xrightarrow[\text{(ii) H}_2\text{O}_2, \text{NaOH}]{\text{(i) B}_2\text{H}_6} \text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$	Reduced	Reduced
		Oxidized	Oxidized
C.	$\text{CH}_3\text{CH}_2\overset{*}{\text{C}}\text{H}=\text{C} \xrightarrow{\text{N}_2\text{H}_4} \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$	Reduced	
		Oxidized	
D.	$\text{CH}_3\text{CH}_2\overset{*}{\text{C}}\text{H}=\text{O} \xrightarrow[\text{H}_2\text{O, pH} > 8]{\text{Ag}^{(+)}} \text{CH}_3\text{CH}_2\text{CO}_2\text{H}$	Reduced	
		Oxidized	
E.	$\text{CH}_3\text{COCH}_2\text{CH}_2\text{CO}_2\text{H} \xrightarrow{\text{Heat}} \text{CH}_3\text{COCH}_3 + \text{O}=\text{C}=\text{O}$	Reduced	Reduced
		Oxidized	Oxidized
F.	$\text{H}_2\text{C}=\text{C}(\text{OH})\text{C}_2\text{H}_5 \xrightarrow{\text{tautomerization}} \text{H}_3\text{CCOC}_2\text{H}_5$	Reduced	Reduced
		Oxidized	Oxidized

10. Consider the possible formation of an aldehyde or ketone product when each of the ten compounds in the column on the left is treated with each of the reagents shown in the top row. Check the designated answer box if you believe an aldehyde or ketone will be formed.

Assume that the reagents may be present in excess. For each checked reaction, try to draw the structure of the major product (s).

ALDEHYDES AND KETONES

417

Starting	PCC $\text{C}_6\text{H}_5\text{NHCrO}_3\text{Cl}$	Jone's Reagent CrO_3 in aq. acid	$\text{Pb}(\text{OAc})_4$ [or HIO_4]	(i) O_3 , (ii) Zn dust	H_3O^+	(i) BH_3 in THF (ii) $\text{H}_2\text{O}_2 + \text{NaOH}$
						
						
						
						
						
						
						
						
						
						



418

ORGANIC Chemistry for IIT-JEE












11. Comprehension

Wittig reaction :

The reaction of a phosphorus ylide with an aldehyde (or) ketone introduces a carbon-carbon double bond in place of the carbonyl bond.

Mechanism :






Driving force of the reaction is high bond energy of (P=O). ($\Delta H = -ve$)

(a) *cis*-2-butene(b) *trans*-2-butene(c) *iso*-butene



(d) 1-butene

ALDEHYDES AND KETONES













419



E. Identify major product in given intramolecular Wittig reaction :







Product (A) and (B) respectively are :



420

ORGANIC Chemistry for IIT-JEE


12. Match the column :

Column (I)		Column (II)	
Conversion		Reagent	
(a)		(p)	$\text{NH}_2/\text{NH}_2/\text{HO}^\ominus, \Delta$ (Wolff-Kishner reduction)
(b)		(q)	$\text{Zn(Hg)}, \text{HCl}$ (Clemmensen reduction)
(c)		(r)	LiAlH_4
(d)		(s)	None





ALDEHYDES AND KETONES

421





13. Comprehension



A. Reactant (A) is :



B. Product (B) is :



422

ORGANIC Chemistry for IIT-JEE



ANSWERS — LEVEL 2

1. A – l; B – g; C – n; D – k; E – h; F – c; G – j; H – f; I – m; J – d; K – e; L – b

2. i – A, B, C, E, F; ii – D, G, H; iii – B, E, F

3. a – p, r; b – r; c – s; d – p, r

4. A : $\text{Ph}-\overset{\text{OH}}{\underset{|}{\text{CH}}}-\text{CH}_3$; B : $\text{Ph}-\text{CH}_2-\text{COOH}$; C : $\text{Ph}-\text{CH}=\text{C}\begin{matrix} \text{CH}_3 \\ \text{CH}_3 \end{matrix}$;



5. A – c; B – b

6. $A + B + C = 17$

7. set 1 – A; set 2 – ; set 3 – B; set 4 – ; set 5 – A; set 6 – B; set 7 – B;
set 8 – B; set 9 – A; set 10 – B









8. A – p, q, s; B – p; C – p, q, s; D – p, q, s

9. A : both are oxidized; B : C^{12} is reduced, C^{14} is oxidized; C : reduced; D : oxidized
E : C^{12} is reduced, C^{14} is oxidized; F : C^{12} is reduced, C^{14} is oxidized

ALDEHYDES AND KETONES

423


10.

Compound	PCC $\text{C}_5\text{H}_5\text{NHCrO}_3\text{Cl}$	Jone's Reagent CrO_3 in aq. acid	$\text{Pb}(\text{OAc})_4$ [or HIO_4]	(i) O_3 , (ii) Zn dust	H_3O^+	(i) BH_3 in THF (ii) $\text{H}_2\text{O}_2 + \text{NaOH}$
	✓	✓	✗	✗	✗	✗
	✗	✗	✗	✓	✓	✓
	✓	✓	✗	✓	✗	✗
	✗	✗	✗	✓	✗	✗
	✓	✓	✗	✓	✓	✓
	✗	✗	✗	✗	✗	✗
	✗	✗	✗	✓	✓	✓
	✗	✗	✗	✓	✓	✓
	✓	✓	✗	✗	✗	✗
	✓	✓	✓	✗	✗	✗

11. A – a; B – a; C – b; D – b; E – a; F – b

12. a – q; b – s; c – r; d – p

13. A. (c) B. (d)









8

ALDOL AND CANNIZARO REACTION







1. Compound A and B, both were treated with NaOH, producing a single compound C.




ALDOL AND CANNIZARO REACTION**425**

- (a) Dehydration, Hydrolysis
 (b) Retro aldol and further condensation
 (c) Perkin condensation & Clemmensen reduction
 (d) Clemmensen and Perkin condensation

3. This is an example of an intramolecular aldol reaction :


















- (a) $\text{Ph} - (\text{CH} = \text{CH})_2 - \text{CHO}$ (b) $\text{Ph} - (\text{CH} = \text{CH})_3\text{CHO}$
 (c) $\text{Ph} - (\text{CH} = \text{CH})_4\text{CHO}$ (d) $\text{Ph} - \text{CH} = \text{CH} - \text{CH} = \text{CH} - \text{CH}_3$








- (a) propanol (b) ethanol (c) butanol (d) pentanol

426

ORGANIC Chemistry for IIT-JEE




10. The reaction ,




ALDOL AND CANNIZARO REACTION

427

11. Compare enolate A with enolate B.







A









B

Which of the following statements is true ?





- (a) A is more stable than B
(b) A and B have the same stability
(c) B is more stable than A
(d) No comparison of stability can be made
12. Benzalacetone is the product of mixed aldol condensation between benzaldehyde ($\text{C}_6\text{H}_5\text{CH}=\text{O}$) and acetone [$(\text{CH}_3)_2\text{C}=\text{O}$]. What is its structure ?



13. Identify the major product P in the following reaction:





Product (B) is:




428


ORGANIC Chemistry for IIT-JEE




16. Identify the principal product of the following reaction?







17. Which one of the following compounds is the best choice for being prepared by an efficient mixed aldol addition reaction?









18. Identify the major product P in the following reaction:












ALDOL AND CANNIZARO REACTION**429**

19. The enolate ion that reacts with 3-buten-2-one to form (Y) is :








Product (B) in the above reaction is :



430


ORGANIC Chemistry for IIT-JEE

22. An organic compound with the molecular formula $C_9H_{10}O$ forms a 2,4-DNP derivative, reduces Tollen's reagent and undergoes Cannizzaro reaction, on vigorous oxidation it gives 1,2-benzenedicarboxylic acid. Structure of organic compound is:







Number of intramolecular aldol condensation product is :

- (a) 1 (b) 2 (c) 3 (d) 4





Compound (A) exist in geometrical isomers and (B) gives Cannizzaro reaction.

(A) will be :







ALDOL AND CANNIZARO REACTION**431**

25. Which of the following compounds will not undergo Cannizzaro reaction, when treated with 50% aqueous alkali?





Above reaction is known as :

- (a) Cannizzaro reaction, Disproportionation reaction
 (b) Tischenko reaction, Disproportionation reaction
 (c) Cross Cannizzaro reaction, Redox reaction
 (d) Tischenko reaction, Redox reaction







432

ORGANIC Chemistry for IIT-JEE











29. Choose the most reasonable reaction intermediate for the following reaction.



(d) None of these


















Product (B) of the above reaction is :



(d) none is correct

ALDOL AND CANNIZARO REACTION
















433








434

ORGANIC Chemistry for IIT-JEE

35. Which of the following reactant on reaction with conc. NaOH followed by acidification gives the following lactone as the product ?



ALDOL AND CANNIZARO REACTION**435****38.** The following reaction gives:

ATDB.uno

ANSWERS — LEVEL 1

1.	(a)	2.	(b)	3.	(a)	4.	(b)	5.	(b)	6.	(c)	7.	(b)	8.	(b)
9.	(b)	10.	(c)	11.	(a)	12.	(a)	13.	(a)	14.	(b)	15.	(a)	16.	(b)
17.	(b)	18.	(a)	19.	(c)	20.	(a)	21.	(c)	22.	(b)	23.	(c)	24.	(c)
25.	(d)	26.	(a)	27.	(c)	28.	(b)	29.	(c)	30.	(c)	31.	(a)	32.	(a)
33.	(c)	34.	(a)	35.	(c)	36.	(b)	37.	(b)	38.	(b)				


436

ORGANIC Chemistry for IIT-JEE






1. Comprehension

Mechanism of Cannizzaro's reaction of benzaldehyde is



A. Which of the following reactants can undergo Cannizzaro's reaction.?



(d) All of these

B. Order of the above reaction is.





(a) 1

(b) 2

(c) 3

(d) 4

C. Which of the following is best hydride donor in Cannizzaro's reaction?



D. Cannizzaro's reaction is:





(a) Reduction

(b) Disproportionation reaction

(c) Oxidation








(d) Ion - exchange reaction

E. Which of the following cannot undergo intramolecular Cannizzaro's reaction?

ALDOL AND CANNIZARO REACTION**437**

2. Aldol condensation proceeds by carbon-carbon bond formation between an enolate donor and a carbonyl acceptor. For each of the following aldol products (a through e) select a donor and an acceptor compound from the list at the bottom of the page (compounds A through H). Write the letter corresponding to your selection in the appropriate answer box.

	Aldol Product	Donor	Acceptor
a.			
b.			
c.			
d.	$(\text{CH}_3)_2\text{C}(\text{OH})\text{CH}_2\text{COCH}_3$		
e.			



438

ORGANIC Chemistry for IIT-JEE

3. Comprehension

During an experimental workup procedure, a chemist treated a starting material with NaOH in the solvent acetone $[(\text{CH}_3)_2\text{C}=\text{O}]$; however, the starting material was recovered unreacted. Instead, the chemist isolated a small amount of Product A (shown below).

Product A





The chemist determined that Product A resulted from the aldol self-condensation of acetone. Product A was identified based on the following observations.

Observations about Product A

1. Elemental analysis of Product A indicated that it consisted only of carbon, hydrogen, and oxygen.
2. product A had a molecular weight of 116 g/mol.
3. Product A was a methyl ketone because it gave a positive iodoform test.
4. When product A was treated with Br_2 in CCl_4 , the red bromine colour persisted, because no carbon-carbon double bonds were present to react with the bromine.


The structure of Product A was further confirmed when treatment with hot sulfuric acid resulted in the corresponding dehydration product, Product B.

- A. What is the molecular weight of a compound that undergoes an aldol self-condensation reaction to result in a β -hydroxy ketone with a molecular weight of 144?
- (a) 70 g/mol (b) 72 g/mol
(c) 74 g/mol (d) 76 g/mol
- B. The aldol self-condensation of acetone is an equilibrium that favours acetone over its condensation product. Which of the following experimental modifications is most likely to shift the position of equilibrium toward Product A?
- (a) Using only a catalytic amount of NaOH
(b) Using only a catalytic amount of acetone
(c) Removing Product A as it is formed
(d) Increasing the reaction temperature to the boiling point of acetone
- C. Based only on observation 1 and 2, which of the following compounds could have been Product A?




ALDOL AND CANNIZARO REACTION**439**


- D.** When a drop of Br_2 in CCl_4 is added to Product B, the resulting solution will be :
 (a) colourless, because Product B does not contain a carbon-carbon double bond
 (b) colourless, because Product B contains a carbon-carbon double bond
 (c) red, because Product B does not contain a carbon-carbon double bond
 (d) red, because Product B contains a carbon-carbon double bond
- E.** Which of the following compounds from the passage will give a positive iodoform test ?
 (a) Product A only (b) Product A and Product B
 (c) Product A and acetone only (d) Product A, Product B, and acetone

4. Comprehension

- A.** Structure of A is :
 (a) $\text{H}_2\text{C} = \text{CH} - \text{CHO}$ (b) $\text{Ph} - \text{CH} = \text{CH} - \text{CH}_3$
 (c) $\text{Ph} - \underset{\text{CH}_3}{\text{C}} = \text{CH}_2$ (d) $\text{Ph} - \underset{\text{CH}_3}{\text{CH}} = \text{C} - \text{CH}_3$
- B.** Structure of (B) and (C) differentiated by :
 (a) Tollen's reagent (b) Fehling solution (c) 2,4-DNP (d) NaHSO_3
- C.** Structure of E is :

**ANSWERS — LEVEL 2**

- A - d; B - c; C - a; D - b; E - c
- a - Donor = C, Acceptor = C; b - Donor = E, Acceptor = D;
c - Donor = B, Acceptor = A; d - Donor = G, Acceptor = G; e - Donor = F, Acceptor = B
- A - b; B - c; C - d; D - b; E - d
- A - b, B - b, C - c








9

CARBOXYLIC ACID AND THEIR DERIVATIVES







1. Identify C in the following sequence of reactions :







2. Saponification (basic hydrolysis) of $\text{C}_6\text{H}_5\text{C}(^\ominus)\text{CH}_3$ will yield :

[$^\ominus$ = mass - 18 isotope of oxygen]







CARBOXYLIC ACID AND THEIR DERIVATIVES**441**

Product (X) of the reaction is :



4. Which of the following is the correct order of decarboxylation of β -keto carboxylate anion ?








(a) $a > b > c > d$





(b) $c > d > a > b$

(c) $c > d > b > a$

(d) $d > c > a > b$




6. Which β -keto acid shown will not undergo decarboxylation ?



442

ORGANIC Chemistry for IIT-JEE

7. Choose the response that matches the correct functional group classification with the following group of structural formulas.




(a) Anhydride

(b) Lactam

(c) Imide

(d) Imide




Lactam

Imide

Lactone

Lactam







Lactone

Lactone





Anhydride

Lactone





8. ; Product (P) of the reaction is :



9. (Lactic acid) ; Product (X) of the reaction is :






10. ; Product (A) obtained is :




CARBOXYLIC ACID AND THEIR DERIVATIVES





443


11. Which of the following acid on heating gives geometrical isomers as a product ?








(d) All of these





12.  $\xrightarrow{\text{MeOH}}$ A $\xrightarrow{\text{PCl}_3}$ B $\xrightarrow{\text{MeNH}_2}$ C ; Product (C) of the reaction is :



13.  $\xrightarrow{\text{LiAlH}_4}$ A $\xrightarrow{\text{PCC}}$ B $\xrightarrow{\text{NH}_2-\text{NH}_2}$ C , Product (C) is :




14.  $\xrightarrow{-\text{CO}_2}$ A $\xrightarrow{\text{HCl}}$ B $\xrightarrow{\text{aq KOH}}$ C. Product (C) is :







444




ORGANIC Chemistry for IIT-JEE








In above reaction identify major product (A) of the reaction:



16. An optically active compound 'X' has molecular formula $C_4H_8O_3$. It evolves CO_2 with $NaHCO_3$. 'X' reacts with $LiAlH_4$ to give an achiral compound. 'X' is :







17. $CH_3 - \overset{O}{\parallel} C - O - CH_2 - CH_3 + H - \bullet^- \longrightarrow (\bullet = O^{18})$ One of the product of the reaction is :








Product (X) is :

CARBOXYLIC ACID AND THEIR DERIVATIVES






445










19. Identify final product in the following reaction;










20. Select the final product from this sequence of reactions.



21.




Which of the following reactants will complete the above reaction ?

23. Arrange the following in order of increasing reactivity (least $\longrightarrow$ most) towards nucleophile

446

ORGANIC Chemistry for IIT-JEE

24. Choose the best sequence of reactions for transformation given. Semicolons indicate separate reaction steps to be used in the order shown.




- (a) H_3O^+ ; SOCl_2 ; CH_3NH_2
 (b) $\text{HO}^-/\text{H}_2\text{O}$; PBr_3 ; Mg ; CO_2 ; H_3O^+ ; SOCl_2 ; CH_3NH_2
 (c) LiAlH_4 ; H_2O ; HBr ; Mg ; CO_2 ; H_3O^+ ; SOCl_2 ; CH_3NH_2
 (d) None of these would yield the desired product
25. A key step in the hydrolysis of acetamide in aqueous acid proceeds by nucleophilic addition of :

- (a) H_3O^+ to $\text{CH}_3\text{C}(=\text{O})\text{NH}_2$ (b) H_2O to $\text{CH}_3\text{C}(\text{OH})(\text{NH}_2)^+$
 (c) H_3O^+ to $\text{CH}_3\text{C}(\text{OH})(\text{NH}_2)^+$ (d) HO^- to $\text{CH}_3\text{C}(=\text{O})\text{NH}_2$

26. Which reaction is not possible for acetic anhydride ?

- (a) $(\text{CH}_3\text{C}(=\text{O}))_2\text{O} + 2\text{HN}(\text{CH}_3)_2 \longrightarrow \text{CH}_3\text{C}(=\text{O})\text{N}(\text{CH}_3)_2 + \text{CH}_3\text{CO}_2^- + \text{H}_2\text{N}^+(\text{CH}_3)_2$
 (b) $(\text{CH}_3\text{C}(=\text{O}))_2\text{O} + \text{H}_2\text{NCH}_2\text{CH}_2\text{OH} \longrightarrow \text{H}_2\text{NCH}_2\text{CH}_2\text{OC}(=\text{O})\text{CH}_3 + \text{H}^+\text{CH}_2\text{CH}_2\text{O}^-$
 (c) $(\text{CH}_3\text{C}(=\text{O}))_2\text{O} + \text{C}_6\text{H}_6 \xrightarrow{\text{AlCl}_3} \text{CH}_3\text{C}(=\text{O})\text{C}_6\text{H}_5 + \text{CH}_3\text{CO}_2\text{H}$
 (d) $(\text{CH}_3\text{C}(=\text{O}))_2\text{O} + \text{NaCl} \longrightarrow \text{CH}_3\text{C}(=\text{O})\text{Cl} + \text{CH}_3\text{CO}_2^-\text{Na}^+$

27. All but one of the following compounds react with aniline to give acetanilide. Which one does not?







- (a) $\text{CH}_3\text{C}(=\text{O})\text{Cl}$ (b) $\text{H}_3\text{C}-\text{C}(=\text{O})-\text{O}-\text{C}(=\text{O})-\text{CH}_3$
 (c) $\text{CH}_3\text{C}(=\text{O})\text{H}$ (d) $\text{C}_6\text{H}_5-\text{OC}(=\text{O})\text{CH}_3$





28. Which of the following best describes the nucleophilic addition step in the acid-catalyzed hydrolysis of acetonitrile (CH_3CN) ?

CARBOXYLIC ACID AND THEIR DERIVATIVES

447



29. The major product expected, when Phthalamide is treated with NaOH, is :



30. Which of following acid remains unaffected on heating ?

(a) malonic acid

(b) maleic acid

(c) Fumaric acid

(d) Succinic acid

31. $\text{Br}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{Br}$ reacts with $\text{C}_2\text{H}_5\text{MgBr}$ to give a cyclic product.







At which value of n the formation of six membered ring takes place ?

(a) $n = 2$

(b) $n = 3$

(c) $n = 5$

(d) $n = 6$




448






ORGANIC Chemistry for IIT-JEE

- (a) *cis*-anhydride
(c) both (a) & (b)

- (b) *trans*-anhydride
(d) mono-basic acid













Product (A) of the reaction is :



Product (A) of the reaction is :







- (a) Ethylene glycol
(c) Glyceryltrinitrate (explosive)

- (b) Glycerol
(d) Cumene hydrogen peroxide



CARBOXYLIC ACID AND THEIR DERIVATIVES

449



Above reaction is an example of :






(a) Esterification

(b) Saponification

(c) Hydrolysis



(d) Trans Esterification

40. Which of the following is an intermediate formed in the reaction shown below ?





(Principal component of coconut oil.)

Product is obtained in the above reaction is :





(c) Both (a) and (b) (d) None of these

42. The reaction of sodium acetate with acetyl chloride proceeds through which of the following mechanisms ?








450






ORGANIC Chemistry for IIT-JEE








43. Which is the major product of the following reaction ?



44. Which is the major product of the following reaction ?




45. Ethanoic acid + 3-methyl-1-butanol $\xrightleftharpoons[\text{H}_2\text{SO}_4]{\text{traces}}$ (A); Compound (A) is :




CARBOXYLIC ACID AND THEIR DERIVATIVES

451





47. Which of the following compounds will undergo decarboxylation on heating ?



(a) 2 and 3 (b) 3 and 4 (c) 3 only (d) 1 and 4


48. Which one of the following is not an intermediate in the generally accepted mechanism for the reaction shown below ?



49. $\text{ROCH}_2\text{CO}_2\text{H} \xrightarrow[\text{(major product)}]{\text{3 eq. EtOH, dry HCl gas}} \text{(A)}$; Product A is :







50. Identify the compound C in the following sequence :








452





ORGANIC Chemistry for IIT-JEE











51. What is the final product (B) of this sequence ?



52. Which of the following undergoes decarboxylation most readily on being heated ?





53. What is compound Z ?






CARBOXYLIC ACID AND THEIR DERIVATIVES





453





55. $\text{CH}_3 - \text{CH} = \text{CH} - \text{CH}_2 - \text{CO}_2\text{H} \xrightarrow{\Delta} (\text{X})$ (major); Product (X) is :





56. $\text{H} - \text{O} - \overset{\text{O}}{\parallel} \text{C} - (\text{CH}_2)_n - \overset{\text{O}}{\parallel} \text{C} - \text{O} - \text{H} \xrightarrow{\Delta} \text{product}$, At what value of (n) given compound will not evolve CO_2 gas ?



57. CO_2H
|
 $(\text{CH}_2)_n$; If ($n = 4$) then di-carboxylic acid would be known as :






Product (A) of the above reaction is :








454

ORGANIC Chemistry for IIT-JEE



Identify (A).






How many product will be formed when above compound undergo de-carboxylation?

(a) 0

(b) 1

(c) 2

(d) 3




(c) Both (a) and (b)

(d) none of these

CARBOXYLIC ACID AND THEIR DERIVATIVES**455**


62. Products obtained in the given reactions are shown below.



The number of possible products for x and y are :


- (a) 1, 1 (b) 1, 2 (c) 2, 1 (d) 2, 2

63.



Product (A) of the above reaction is :

- (a) $\text{Ph}-\text{NH}_2$ (b) $\text{Ph}-\text{CH}_2-\text{NH}_2$
 (c) $\text{Ph}-\text{CH}_2-\text{NH}-\text{CO}_2\text{H}$ (d) $\text{Ph}-\text{CH}_2-\text{NH}-\text{CHO}$
64. Which of the following pair is C_2 -epimer ?
 (a) *D*-Glucose, *D*-Maltose (b) *D*-Glucose, *D*-Mannose
 (c) *D*-Allose, *D*-Ribose (d) *D*-Glucose, *D*-Arabinose
65. Total number of enol possible for the compound formed during given reaction will be (including stereoisomer):



- (a) 2 (b) 3 (c) 4 (d) 5

ANSWERS — LEVEL 1

1.	(b)	2.	(b)	3.	(a)	4.	(c)	5.	(a)	6.	(b)	7.	(d)	8.	(b)
9.	(b)	10.	(b)	11.	(d)	12.	(b)	13.	(b)	14.	(a)	15.	(b)	16.	(c)
17.	(c)	18.	(b)	19.	(a)	20.	(a)	21.	(b)	22.	(c)	23.	(b)	24.	(a)
25.	(b)	26.	(d)	27.	(c)	28.	(d)	29.	(c)	30.	(c)	31.	(b)	32.	(c)
33.	(a)	34.	(a)	35.	(b)	36.	(a)	37.	(b)	38.	(b)	39.	(d)	40.	(d)
41.	(c)	42.	(c)	43.	(b)	44.	(c)	45.	(b)	46.	(a)	47.	(c)	48.	(b)
49.	(b)	50.	(c)	51.	(d)	52.	(d)	53.	(d)	54.	(d)	55.	(c)	56.	(c)
57.	(c)	58.	(c)	59.	(b)	60.	(b)	61.	(c)	62.	(c)	63.	(b)	64.	(b)
65.	(b)														

456

ORGANIC Chemistry for IIT-JEE




1. Match the Column (I) and (II). (Matrix)

Column (I)		Column (II)	
Reaction		Products formed	
(a)	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{HO}_2\text{C} - \text{C} - \text{CO}_2\text{H} \\ \quad \\ \text{H} \quad \text{D} \\ \\ \text{Ph} \end{array} \xrightarrow{\Delta} $	(p)	Diastereomers
(b)	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{HO}_2\text{C} - \text{C} - \text{CO}_2\text{H} \\ \\ \text{Et} \end{array} \longrightarrow $	(q)	Racemic mixture
(c)	$\xrightarrow{\Delta}$	(r)	Meso compound
(d)	$\xrightarrow{\Delta}$	(s)	CO ₂ gas will evolve

ANSWERS — LEVEL 2

1. a – p, s; b – q, s; c – p, s; d – r








10 AMINES







ATDB.uno

1. In which of the following reaction cyanide will be obtained as a major product ?








Product (A) is :







458

ORGANIC Chemistry for IIT-JEE

3. Which of the following alkene cannot be prepared by de-amination of $n\text{-Bu}-\text{NH}_2$ with NaNO_2/HCl ? (n-Butyl)
- (a) 1-butene (b) *cis*-2-butene (c) *trans*-2-butene (d) Iso-butene
4. Predict the major product P in the following reaction.







5. Product of this reaction is :



6. + $\text{CH}_2 = \text{CH}-\text{CH}_2-\text{CH}_2-\text{OH}$
- (A) 48% + (B) 47%


A will be :



AMINES






459

7. Which of the following isomers of C_8H_9NO is the weakest base ?
- (a) *o*-Aminoacetophenone (b) *p*-Aminoacetophenone
(c) *m*-Aminoacetophenone (d) Acetanilide
8. Rank the following compounds in order of increasing basic strength. (weakest $\rightarrow$ strongest) :








- (a) $4 < 2 < 1 < 3$ (b) $4 < 3 < 1 < 2$
(c) $4 < 1 < 3 < 2$ (d) $2 < 1 < 3 < 4$

9. Which of the following arylamines will not form a diazonium salt on reaction with sodium nitrite in hydrochloric acid ?
- (a) *m*-Ethylaniline (b) *p*-Aminoacetophenone
(c) 4-Chloro-2-nitroaniline (d) *N*-Ethyl-2-methylaniline
10. Identify product *D* in the following reaction sequence :




11. Which one of the following is best catalyst for the reaction shown below ?



460

ORGANIC Chemistry for IIT-JEE





12. The major products obtained from the following sequence of reactions are :



- (a) $(\text{CH}_3)_2\text{CHCH}_2\text{NH}_2 + \text{H}_2\text{C} = \text{CH}_2$ (b) $(\text{CH}_3)_2\text{NCH}_2\text{CH}_3 + \text{H}_2\text{C} = \text{C}(\text{CH}_3)_2$

- (c) $(\text{CH}_3)_2\text{CHCH}_2\text{N}(\text{CH}_3)\text{CH}_2\text{CH}_3 + \text{H}_2\text{C} = \text{CH}_2$ (d) $(\text{CH}_3)_3\text{N}^+\text{CH}_2\text{CH}_3\text{I}^- + \text{H}_2\text{C} = \text{CH}_2$



13. Which amine yields *N*-nitroso amine after treatment with nitrous acid ($\text{NaNO}_2, \text{HCl}$) ?




14. $\xrightarrow{\text{HNO}_2}$ (A); Product (A) is :

- (a) cyclopentane carboxy aldehyde (b) cyclohexane-1,2-diol
(c) 2-aminocyclohexene (d) cyclohex-2-enol

15. Choose the appropriate product for this reaction.






16. Which of the following product will be obtained in the given (consider minor product also) Beckmann-type rearrangement ?



AMINES

461



(d) all of these

17. Deamination (or) diazotization of $n\text{-Bu-NH}_2$ with NaNO_2/HCl gives isomeric butene.

(a) 2

(b) 3

(c) 4

(d) 5

ATDB.uno

ANSWERS — LEVEL 1











1	(c)	2	(a)	3	(d)	4	(a)	5	(b)	6	(b)	7	(d)	8	(b)
9	(d)	10	(b)	11	(c)	12	(c)	13	(d)	14	(a)	15	(b)	16	(d)
17	(b)														

462

ORGANIC Chemistry for IIT-JEE

LEVEL-2


1. Five amine syntheses are outlined below. In each reaction box enter a single letter designating the best reagent and conditions selected from the list at the bottom of the page.

A.		First Step Second Step	
B.		First Step Second Step Third Step	
C.		First Step Second Step	
D.		First Step Second Step	
E.		First Step Second Step Third Step Fourth Step	

(a)	(i) LiAlH_4 in ether (ii) H_2O & base	
(b)	$\text{C}_2\text{H}_5\text{NH}_2$ (cat. $\text{H}^{(+)}$)	
(c)	NaCN in alcohol	
(d)	H_2 & Ni catalyst or H_2 & Pd catalyst	
(e)	NaN_3 in alcohol	
(f)	$(\text{CH}_3\text{CO})_2\text{O}$ & pyridine	
(g)	$\text{C}_2\text{H}_5\text{Br}$	

AMINES


463

(h)	 , $\text{H}^{\oplus}$	
(i)	$2\text{CH}_3\text{I}$ & pyridine	
(j)	KOH in H_2O	


ATDB.uno

ANSWERS — LEVEL 2

1. A – c, a or c, d; B – b, d, f; C – h, d; D – d, i or a, i; E – e, a, h, a

































11 CARBENE AND NITRENE



(α -hydroxy amide)








Product of this Hoffmann bromamide reaction is :

















CARBENE AND NITRENE**465**

466

ORGANIC Chemistry for IIT-JEE




Product (A) & (B) respectively in the above reaction are :






CARBENE AND NITRENE

467



Product (A) is :




when $\gamma = CH_3CO_2^-$, $Cl-C(=O)-CH_2-CO_2^-$, $n-C_4H_9CO_2^-$
(i) (ii) (iii)

- (a) (i) > (ii) > (iii) (b) (iii) > (i) > (ii)
(c) (iii) > (ii) > (i) (d) (iii) > (i) > (ii)



15. When primary amine reacts with chloroform in ethanolic KOH, then product is :

- (a) an isocyanide (b) an aldehyde
(c) a cyanide (d) an alcohol

16. The reaction of chloroform with alcoholic KOH and *p*-toluidine forms :








17. What is the product (Q) of the following reaction ?











468

ORGANIC Chemistry for IIT-JEE



19. Which of the following reaction, does not give chloro benzene as a product ?



x = moles of KOH consumed is :

(a) 1

(b) 2


(c) 3

(d) 4



22. Heating the acyl azide in dry toluene under reflux for 3-hours give a 90% yield for a heterocyclic product. Identify the product (A).

CARBENE AND NITRENE

469




23.




470



ORGANIC Chemistry for IIT-JEE





Product (A) of the above reaction is :



25. A rather interesting example of the Wolff rearrangement with 2-diazocyclohexanone in methanol is given below. Identify the major product :




26. The orbital picture of a singlet carbene (:CH₂) can be drawn as :






CARBENE AND NITRENE

471









(d) none of these

27. The orbital picture of a triplet carbene can be drawn as :




(d) none of these

28.



29.




Select the suitable reagent for above conversion.


(a) $\text{CH}_2\text{N}_2 / \Delta$ (b) $\text{CBr}_4 / \text{RLi}$ (c) $\text{H}_2\text{C} = \text{CH}_2$ (d) $t\text{-BuOK}$

472




ORGANIC Chemistry for IIT-JEE




Product (A) will be :




31. The major product formed in the following reaction is



(c) 50 : 50 mixture of above two compounds (d)




32.







To carry out above conversion reagent used in decreasing order.












(a) Na/liq. NH_3 , $\text{CHBr}_3/\text{NaOH}(\Delta)$ (b) $\text{H}_2/\text{Pd} - \text{CaCO}_3$, $\text{CHBr}_3/\text{NaOH}(\Delta)$ (c) Na/liq. NH_3 , $\text{CHCl}_3/\text{NaOH}$ (d) $\text{H}_2/\text{Pd} - \text{CaCO}_3$, $\text{CHCl}_3/\text{NaOH}$

33.



Product (A) of the reaction is :



CARBENE AND NITRENE**473**

Which of the following product(s) is/are can be obtained in the above reaction.

- (a) Isopentane (b) 3-Methyl hexane (c) *n*-Pentane (d) 3-Methyl pentane

ANSWERS — LEVEL 1

1.	(b)	2.	(b)	3.	(c)	4.	(b)	5.	(a)	6.	(d)	7.	(c)	8.	(b)
9.	(c)	10.	(b)	11.	(a)	12.	(a)	13.	(b)	14.	(c)	15.	(a)	16.	(b)
17.	(d)	18.	(a)	19.	(d)	20.	(a)	21.	(c)	22.	(a)	23.	(b)	24.	(b)
25.	(a)	26.	(a)	27.	(c)	28.	(c)	29.	(a)	30.	(b)	31.	(a)	32.	(b)
33.	(c)	34.	(c)	35.	(b)	36.	(d)								

474


ORGANIC Chemistry for IIT-JEE




LEVEL-2

1. Comprehension

Hoffmann bromamide reaction involves conversion of a carboxylic acid amide into an amine with a loss of a carbon atom on treatment with aqueous sodium hypobromite. Thus Hoffmann result in shortening of a carbon chain.



Mechanism of the reaction is :




A. Number of moles of NaOH consumed in above reaction.

(a) 1





(b) 2





(c) 3

(d) 4

(a) $\text{Ph}-\text{NH}_2$ (b) $\text{Ph}-\text{CH}_2-\text{NH}_2$ (c) $\text{Ph}-\text{NH}-\text{CH}_3$ (d) $\text{Ph}-\text{N} \begin{array}{l} \nearrow \text{CH}_3 \\ \searrow \text{CH}_3 \end{array}$

C. Which of the following will not give Hoffmann bromamide reaction.




CARBENE AND NITRENE**475**

(d) None of these

2. Comprehension

Given is mechanism of Beckmann rearrangement.




A. Rate determining step in Beckmann rearrangement is :

(a) I

(b) II

(c) III


(d) IV



On treatment H_2SO_4 followed by hydrolysis in acidic medium above compound gives.

(a) $\text{CH}_3 - \text{CO}_2\text{H}$, $\text{Ph} - \text{NH}_2$ (b) $\text{CH}_3 - \text{NH}_2$, $\text{Ph} - \text{CO}_2\text{H}$ (c) $\text{Ph} - \text{CH}_2 - \text{NH}_2$ + $\text{Ph} - \text{CO}_2\text{H}$ (d) $\text{Ph} - \text{CO}_2\text{H}$ + $\text{CH}_3 - \text{CO}_2\text{H}$





C. Which of the following reagent cannot used in Beckmann rearrangement ?

(a) TsOH (b) $\text{R} - \text{SO}_2\text{Cl}$ (c) BF_3 (d) $\text{Ph} - \text{Li}$ 

Product (A) of the above reaction is :

476




ORGANIC Chemistry for IIT-JEE







3. Match the column I and II.

Column (I)		Column (II)	
(a)		(p)	D.B.E. = even for product (Double bond equivalent)
(b)		(q)	D.B.E. = odd for product
(c)		(r)	Ring expansion takes place
(d)		(s)	Carbene will be formed

CARBENE & NITRENE**477****4. Match the column I and II.**

Column (I)		Column (II)	
(a)		(p)	Reimer Tiemann reaction
(b)		(q)	Reimer Tiemann expansion (or) Abnormal RNT reaction
(c)		(r)	Simman-smith reaction.
(d)		(s)	Increase in carbon takes place



5. Match the column I and II.

Column (I)		Column (II)	
(a)		(p)	Aromatic compound will formed
(b)		(q)	Migration take place from carbon to electron deficient nitrogen
(c)		(r)	Carbene will formed in this reaction
(d)		(s)	N ₂ will evolve.




478

ORGANIC Chemistry for IIT-JEE

6. Match the column I and II:

Column (I)		Column (II)	
Reaction		Intermediate	
(a)	$\text{CHCl}_3 + \text{KOH} \xrightarrow{\Delta}$	(p)	Carbocation
(b)	 $\xrightarrow[\Delta]{\text{Ph-Li}}$	(q)	Carbanion
(c)	 $\xrightarrow[\Delta]{\text{Na}}$	(r)	Free radical
(d)	 $\xrightarrow[\Delta]{\text{H}^+}$	(s)	Carbene


7. Matrix :

Column (I)		Column (II)	
Reaction		Product	
(a)	 $\xrightarrow[\text{KOH}, \Delta]{\text{CHCl}_3}$	(p)	
(b)	 $\xrightarrow[\text{KOH}, \Delta]{\text{CHFCIBr}}$	(q)	
(c)	 $\xrightarrow[\text{KOH}, \Delta]{\text{CHCl}_2\text{Br}}$	(r)	


CARBENE & NITRENE

479

(d)




(s)






8. Comprehension




1. Consider the given reaction for preparation of alkyne. (Fritsch reaction).



Anti group will migrate because of less steric hindrance.







C. Rate of reaction when the halide ion:



480

ORGANIC Chemistry for IIT-JEE














9. Comprehension

Wolff rearrangement





When α -Diazoketones are photo-irradiated or heated at high temperature or reacted with silver oxide or silver salts at room temperature, they lose nitrogen and rearrange to form ketene.

The ketenes react rapidly with water, alcohol and amines. Therefore, the reactions called Wolff-rearrangement.













CARBENE & NITRENE

481












(d) None of these



482

ORGANIC Chemistry for IIT-JEE




ANSWERS — LEVEL 2

1. A – d ; B – a ; C – d ; D – a,
2. A – b; B – b; C – d; D – b
3. a – p, r, s; b – q, r, s; c – q, r, s; d – p, r, s
4. a – q, s; b – p, s; c – s; d – r, s
5. a – p, q, s; b – r; c – p, r; d – q
6. a – q, s; b – q, s; c – q, s; d – p
7. a – q; b – p; c – q; d – q
8. A – a; B – c; C – b; D – b
9. A – b; B – c; C – d; D – b; E – b; F – b; G – d

12 AROMATIC COMPOUNDS

LEVEL - 1



Identify the position where electrophilic aromatic substitution (EAS) is most favourable.

- (a) A
(b) B
(c) C
(d) A and C




Correct order of rate of EAS (electrophilic aromatic substitution) is :

- (a) $c > b > a > d$
(b) $c > d > a > b$
(c) $a > b > c > d$
(d) $c > d > b > a$


484

ORGANIC Chemistry for IIT-JEE








Above (C—N) coupling reaction take place at :


- (a) low pH (b) Intermediate pH
(c) high pH (d) any pH
4. Which of the following has the lowest heat of combustion ?







5. The product obtained from the reaction is :



- (a)  (b) 
(c)  (d) 




The end product (C) is :

- (a)  (b) 
(c)  (d) 


AROMATIC COMPOUNDS**485**

7. How many benzylic hydrogens are present in the hydrocarbon shown below ?



- (a) 3 (b) 4 (c) 5 (d) 6 (e) 8

8. The major product formed in the reaction is :




- (a) (b) (c) (d)

9. The major product formed in the reaction is :



- (a) (b) (c) (d)

10. Increasing order of rate of reaction with $\text{HNO}_3/\text{H}_2\text{SO}_4$ is :




- (a) $\text{iii} < \text{ii} < \text{i}$ (b) $\text{ii} < \text{iii} < \text{i}$ (c) $\text{i} < \text{iii} < \text{ii}$ (d) $\text{i} < \text{ii} < \text{iii}$

486


ORGANIC Chemistry for IIT-JEE

11. Increasing order of rate of reaction with $\text{Br}_2/\text{AlCl}_3$ is :




- (a) $\text{iii} < \text{i} < \text{ii} < \text{iv}$ (b) $\text{iv} < \text{ii} < \text{i} < \text{iii}$ (c) $\text{ii} < \text{iv} < \text{iii} < \text{i}$ (d) $\text{iv} < \text{ii} < \text{iii} < \text{i}$

12. Increasing order of equilibrium constant for the formation of a hydrate is :




- (a) $\text{i} < \text{ii} < \text{iii} < \text{iv}$ (b) $\text{iv} < \text{ii} < \text{i} < \text{iii}$ (c) $\text{ii} < \text{iv} < \text{iii} < \text{i}$ (d) $\text{iv} < \text{ii} < \text{iii} < \text{i}$

13. Rank the following reactions A, B and C in order of increasing rate,




- (a) $\text{B} > \text{A} > \text{C}$ (b) $\text{B} > \text{C} > \text{A}$ (c) $\text{A} > \text{B} > \text{C}$ (d) $\text{A} > \text{C} > \text{B}$

14. Rank in order of increasing rate of reaction towards EAS with bromine in the presence of FeBr_3 .



- (a) $\text{B} < \text{A} < \text{C}$
(c) $\text{A} < \text{B} < \text{C}$

- (b) $\text{B} < \text{C} < \text{A}$
(d) $\text{A} < \text{C} < \text{B}$




Identify the position where E.A.S. can take place.






- (a) 1 (b) 2 (c) 3 (d) 4

AROMATIC COMPOUNDS

487



Product (B) in the above reactions is:




(a) 1

(b) 2


(c) 4

(d) 6


18. Arrange the following in decreasing order of reactivity towards EAS (electrophilic aromatic substitution)



(a)



(b)




(c)

(a) $a > b > c$


(b) $c > b > a$

(c) $a > c > b$


(d) $c > a > b$




(a)



(b)



(c)



(d)

Decreasing order of rate of electrophilic aromatic substitution is :

(a) $a > b > c > d$

(b) $a > c > b > d$


(c) $b > a > c > d$

(d) $b > c > a > d$


20. Arrange the following in increasing order of rate of Nitration:

488


ORGANIC Chemistry for IIT-JEE




(a)




(b)



(c)




(d)




(e)

(a) $b < c < a < d < e$ (b) $d < e < a = c < b$ (c) $d < a = c < e < b$ (d) $a < c < b < e < d$


21.



(a)



(b)






(c)


The rate of nitration will be:

(a) $a > b > c$ (b) $a > c > b$ (c) $a = b = c$ (d) $c > a > b$


22. The major product of the reaction is



(c)




(b)




(d)


23. Arrange in their decreasing order of rate of electrophilic aromatic substitution :



(i)



(ii)








(iii)







(a) $i > ii > iii$ (b) $iii > ii > i$ (c) $iii > i > ii$ (d) $i > iii > ii$

AROMATIC COMPOUNDS





489



25. In which of the following compound electrophilic aromatic substitution take place in phenyl ring present in left hand side ?




27. Which of the following compounds is the slowest to react with nitrosonium ion (NO^+)?








490





ORGANIC Chemistry for IIT-JEE




Ibuprofen is :



What is the major product of above Friedel-Craft reaction ?







30. What combination of acid chloride or anhydride and arene would you choose to prepare given compound ?




AROMATIC COMPOUNDS

491



31. In the given conversion best yield will obtained with :



(a) $A = \text{CH}_3 - \text{C}(=\text{O}) - \text{Cl}, \text{AlCl}_3, B = \text{Zn}(\text{Hg}), \text{HCl}$

(b) $A = \text{Zn}(\text{Hg}), \text{HCl}, B = \text{CH}_3 - \text{C}(=\text{O}) - \text{Cl}, \text{AlCl}_3$


(c) $A = \text{CH}_3 - \text{CH}_2 - \text{Cl}, \text{AlCl}_3, B = \text{Zn}(\text{Hg}), \text{HCl}$

(d) $A = \text{NH}_2 - \text{NH}_2 / \text{HO}^-, D, B = \text{CH}_3 - \text{CH}_2 - \text{Cl}, \text{AlCl}_3$

492



ORGANIC Chemistry for IIT-JEE

32. Rank the following in order of decreasing rate of reaction with alkoxide ion ($\text{CH}_3\text{CH}_2\text{O}^-$) in a nucleophilic aromatic substitution reaction :




- (a) $3 > 4 > 1 > 2$ (b) $3 > 4 > 2 > 1$ (c) $2 > 1 > 4 > 3$ (d) $4 > 3 > 2 > 1$



33. Identify the principal organic product of the following reaction.



34. Which position will be attacked most rapidly by the nitronium ion ($-\text{NO}_2^+$) when the compound undergoes nitration with $\text{HNO}_3/\text{H}_2\text{SO}_4$:





- (a) A (b) B (c) C (d) D








AROMATIC COMPOUNDS





493







36. All the hydrocarbons shown are very weak acids. One, however, is far more acidic than the others. Which one is the strongest acid ?



Product (D) in above sequence is :



38. The action of bromine water (excess) on salicylic acid results in the formation of :



494

ORGANIC Chemistry for IIT-JEE

39. What is the correct order of *o/p* ratio when E^+ attacks the following system ?

PhF
A


PhCl
B

PhBr
C

PhI
D

(a) $A < B < C < D$ (b) $A = B = C = D$ (c) $D < C < B < A$ (d) $D < B < A < C$

40. How many products are capable of being formed from toluene in each of following reaction ?




(a) $A = 3, B = 6, C = 8$

(b) $A = 3, B = 6, C = 6$

(c) $A = 3, B = 6, C = 10$

(d) $A = 3, B = 4, C = 6$

41. Nitration takes place at the which position of the given compound ?








(a) A

(b) B










(c) C

(d) D

















AROMATIC COMPOUNDS

495





45. The reaction of toluene with chlorine in the presence of light gives :
















496

ORGANIC Chemistry for IIT-JEE








(b) no reaction














AROMATIC COMPOUNDS

497




Product (A) of the above reaction is :



498

ORGANIC Chemistry for IIT-JEE




above reaction is an example of Nucleophilic aromatic substitution. Which of the following halide ($-X$) is most readily replaced.

- (a) $-F$ (b) $-Cl$ (c) $-Br$ (d) $-I$

57. When comparing the hydrogenation of benzene with that of a hypothetical 1, 3, 5-cyclohexatriene, benzene _____ than the cyclohexatriene.

- (a) absorbs 152 kJ/mol more heat (b) gives off 152 kJ/mol more heat
(c) absorbs 152 kJ/mol less heat (d) gives off 152 kJ/mol less heat

58. Which of the following hydrogens is most easily abstracted on reaction with bromine free radicals, $Br\cdot$?



- (a) a (b) b (c) c (d) d

59. The electrophilic aromatic substitution proceeds through:

- (a) free radical (b) sigma complex (c) benzene (d) carbene

60. Which of the following substitution of benzene is ortho-para in electrophilic substitution and ortho-para in nucleophilic substitution?

- (a) $-NO_2$ (b) $-NO$
(c) $-SO_3H$ (d) $-SO_2Me$


61. The number of possible isomers of dichloronitrobenzene is:

- (a) 3 (b) 4 (c) 6 (d) 8


62. Which of the following is not an aromatic compound?



63. Consider the following sequence of reactions.



The end product (B) is:





AROMATIC COMPOUNDS

499



64. $\text{Ph} - \text{NO}_2 + \text{Et} - \text{Cl} \xrightarrow{\text{AlCl}_3} (A)$, Product (A) of the given reaction is :

(a) $\text{Ph} - \text{NH} - \text{Et}$

(b) no-reaction



65. In nitration of benzene by mixed acid the rate of reaction will be :

(a) $\text{C}_6\text{H}_6 = \text{C}_6\text{D}_6 = \text{C}_6\text{T}_6$ (b) $\text{C}_6\text{H}_6 > \text{C}_6\text{D}_6 > \text{C}_6\text{T}_6$ (c) $\text{C}_6\text{H}_6 = \text{C}_6\text{D}_6 > \text{C}_6\text{T}_6$ (d) $\text{C}_6\text{H}_6 < \text{C}_6\text{D}_6 < \text{C}_6\text{T}_6$ (b) $\text{Ph} - \text{N} = \text{C} - \text{Ph}$ (c) $\text{Ph} - \text{N} = \text{N} - \text{Ph}$ (d) $\text{Ph} - \text{CH} = \text{CH} - \text{Ph}$

67. Which of the following ring compounds obeys Huckel's rule ?

(a) $\text{C}_4\text{H}_4^{-1}$ (b) $\text{C}_4\text{H}_4^{+1}$ (c) $\text{C}_4\text{H}_4^{-2}$ (d) C_4H_4







68. Nitration of which of the following reactant gives maximum of meta product (using $\text{HNO}_3/\text{H}_2\text{SO}_4$) ?

(a) Toluene

(b) Aniline

(c) Benzene

(d) Isopropyl benzene



Which of the following is true statement about the reaction ?

(a) Ortho isomer is major if PhONa is used(b) Para isomer is major if PhOK is used


(c) Product formed is further used for preparation of drug aspirin

(d) All of these


500

ORGANIC Chemistry for IIT-JEE

71. Two benzyne intermediates are likely to be formed equally. Reaction with amide ion can occur in two different directions with each benzyne, giving three possible products. They are formed in a 1 : 2 : 1 ratio. Asterisk (*) refers to ^{14}C .



72. Which one of the following undergoes nucleophilic aromatic substitution at the fastest rate ?



73. For the reaction ; ; the best combination of reactants is :




74. The action of AlCl_3 in Friedel Craft's reaction is:




75. *n*-Butylbenzene on oxidation with hot alkaline KMnO_4 gives :



76. Which sequence of steps describes the best synthesis of 2-phenylpropene ?








77. What are the products of the following reaction ?







AROMATIC COMPOUNDS

501

78. What is the product obtained by heating the following allylic ether of phenol ?







79. When you ingest aspirin, it passes through your stomach, which has an acidic pH, before traveling through the basic environment of your intestine. Provide the structure form as it exists in the intestine.







80. Which of the following sets of reagents, used in the order shown, would be enable for the preparation of *p*-chlorophenol from *p*-chloronitrobenzene ?

- (a) 1. Fe, HCl ; 2. NaOH ; 3. NaNO₂, H₂SO₄ ; 4. H₃PO₂
 (b) 1. Fe, HCl ; 2. NaOH ; 3. NaNO₂, H₂SO₄ ; 4. H₂O, heat
 (c) 1. Fe, HCl ; 2. NaOH ; 3. NaNO₂, H₂SO₄ ; 4. ethanol
 (d) 1. NaOH, heat ; 2. HCl

81. Which one of the following compounds undergoes bromination of its aromatic ring (electrophilic aromatic substitution) at the fastest rate ?














82. Product (Q) in this reaction is :



502

ORGANIC Chemistry for IIT-JEE



This reaction is example of :

(a) Intermolecular C - N coupling

(b) Intramolecular C - N coupling

(c) Intermolecular N - N coupling

(d) Intramolecular N - N coupling

86. The total number of isomeric trimethylbenzene is :





(a) 2

(b) 3

(c) 4

(d) 6


87. Caliene, C_7H_6 , is expected to be a fairly polar aromatic molecule. Which of the following resonance forms contributes to the greatest extent towards the real structure (resonance hybrid) of the molecule ?




AROMATIC COMPOUNDS

503


88. Which of the following molecules is expected to have the greatest resonance stabilization ?




89. In the reaction given below, the major product formed is :




90. *p*-aminophenol reacts with one equivalent of acetyl chloride in the presence of pyridine to give mainly :



91. Which of the following reactions can be used to prepare acetophenone ?



92. Consider the following sequence of reactions.



The products (B) and (C) are :



- (a) benzaldehyde and acetaldehyde (b) benzoic acid and acetic acid
(c) phenol and propionaldehyde (d) phenol and acetone

93. An organic compound having the molecular formula $\text{C}_8\text{H}_{10}\text{O}$ on being heated with I_2 and dilute NaOH gives a yellow precipitate. The expected compound is :








504






ORGANIC Chemistry for IIT-JEE








94. The product (B) of the reaction sequence is :



95. Consider the following sequence of reactions.




96. For the reaction, the product expected is :




AROMATIC COMPOUNDS

505

97. Hydrogenation of naphthalene (*P*) with excess hydrogen gas stops cleanly at 1, 2, 3, 4-tetrahydronaphthalene (*Q*). What conclusion can be drawn from this experiment ?




- (a) the hydrogenation of *P* is exothermic
 (b) one aromatic ring of *P* is more reactive than the aromatic ring of *Q*
 (c) one aromatic ring of *P* is less reactive than the other ring of *Q*
 (d) reduction of the first C = C of *P* is faster than reduction of the second or third C = C
98. Suggest the best reaction conditions for the synthesis shown below.




- (a) (1) HNO_3 , H_2SO_2 ; then (2) Br_2
 (b) (1) Br_2 ; then (2) HNO_3 , H_2SO_2
 (c) (1) CH_3Br , AlBr_3 ; then (2) HNO_3 , H_2SO_3
 (d) HNO_3 , H_2SO_2 , then (2) Br_2 , FeBr_3

99.





In the above compound Cl will be liberated easily in the form of :

- (a) $\text{Cl}^\oplus$ (b) Cl^- (c) $\text{Cl}^\bullet$ (d) Cl^{2+}
100. Consider the following sequence of reactions:





- (a) benzonitrile (b) benzylamine (c) aniline (d) benzamide
101. The major product of the acetylation of salicylic acid with $\text{Ac}_2\text{O}/\text{H}^+$ followed by heating with anhydrous AlCl_3 is :





506

ORGANIC Chemistry for IIT-JEE







102. Which one of the following statements is **True**:







- (a) PhLi adds to both compounds with equal ease
 (b) PhLi does not add to either of the compounds
 (c) PhLi reacts readily with 1 but does not add to 2
 (d) PhLi reacts readily with 2 but does not add to 1

103. The major product expected from the mono-bromination of phenyl benzoate is :





104. The Birch reduction of benzoic acid gives :



105. The decreasing order of reactivity of meta-nitrobromobenzene (I), 2,4,6-trinitrobromobenzene (II), para-nitrobromobenzene (III), and 2,4-dinitrobromobenzene (IV) towards HO^- ions is :

- (a) $\text{I} > \text{II} > \text{III} > \text{IV}$ (b) $\text{II} > \text{IV} > \text{III} > \text{I}$ (c) $\text{IV} > \text{II} > \text{III} > \text{I}$ (d) $\text{II} > \text{IV} > \text{I} > \text{III}$

106. Which of the following tetracarboxylic acid form di-anhydride :








(c) neither (a) nor (b)

(d) both (a) and (b)

AROMATIC COMPOUNDS




507








- (a) $LiAlH_4$ (b) $NaBH_4$
 (c) $LiAlH_4(t-BuO)_3$ (d) PCC/CH_2Cl_2





512






ORGANIC Chemistry for IIT-JEE


127. (a)  (b)  (c)  (d) 

128. $\text{Ph-NH}_2 \xrightarrow{\text{CH}_3\text{-Cl (2mole)}} \text{(A)} \xrightarrow{\text{Ph-N}_2\text{Cl}} \text{(B)}$ (major)  (B)  (C)  (D) 

Product of the above reaction is : 
















129. (a)  (b)  (c)  (d) 

130. $\text{NO}_2\text{-C}_6\text{H}_4\text{-Cl} \xrightarrow{\text{aq. NH}_3} \text{(A)} \xrightarrow{\text{Br}_2} \text{(B)} \xrightarrow{\text{(i) NaNO}_2 + \text{HCl}} \text{(C)}$  (a)  (b)  (c)  (d) 

Product (C) of the above reaction is : 







AROMATIC COMPOUNDS

513




514

ORGANIC Chemistry for IIT-JEE








AROMATIC COMPOUNDS










515



Which of the following compound on hydrolysis gives reactant (A) :






Product (A) of the above reaction is :





516


ORGANIC Chemistry for IIT-JEE



Product (C) of the above reaction is :




Product of the above Friedel-Craft reaction is :






AROMATIC COMPOUNDS

517



143. Which of the following 2-halo nitrobenzene is most reactive towards nucleophilic aromatic substitution ?



144. Choose the best method to prepare given compound :




Benzocaine has been used as a component of appetite suppressants, burn and sunburn remedies. Benzocaine is :






518


ORGANIC Chemistry for IIT-JEE



Predict major product of the above reaction is :



149. Arrange in their decreasing order of rate in $\text{S}_\text{N}\text{Ar}$.







(a) $i > ii > iv > iii > v$

(b) $ii > i > iii > v > iv$

(c) $v > iii > i > ii > iv$

(d) $v > iii > ii > i > iv$


150. Which one of the following compounds undergoes bromination of its aromatic ring (electrophilic aromatic substitution) at the fastest rate ?



AROMATIC COMPOUNDS


519

151. What is the product of the following reaction ?




- (a) (b)
 (c) (d)

152. Which sequence represents the best synthesis of 4-isopropylbenzonitrile ?



4-Isopropylbenzonitrile

- (a) 1. Benzene + $(\text{CH}_3)_2\text{CHCl}$, AlCl_3 ; 2. Br_2 , FeBr_3 ; 3. KCN
 (b) 1. Benzene + $(\text{CH}_3)_2\text{CHCl}$, AlCl_3 ; 2. HNO_3 , H_2SO_4 ; 3. Fe , HCl ,
 4. NaOH 5. Na_2CO_3 , $\text{C}_6\text{H}_5\text{F}_2\text{O}$
 (c) 1. Benzene + $(\text{CH}_3)_2\text{CHCl}$, AlCl_3 ; 2. HNO_3 , H_2SO_4 ; 3. Fe , HCl ;
 4. NaNO_2/HCl 5. KCN
 (d) 1. Benzene + HNO_3 , H_2SO_4 ; 2. $(\text{CH}_3)_2\text{CHCl}$, AlCl_3 ; 3. Fe , HCl ;
 4. NaNO_2 , HCl , H_2O ; 5. CuCN









- (a) (b)
 (c) (d)





520

ORGANIC Chemistry for IIT-JEE











154.



155. What is correct order of rate of nitration of the following compounds ?




156.

157. Compound A ($\text{C}_7\text{H}_8\text{O}$) is insoluble in water, dilute HCl & aqueous NaHCO_3 , but it dissolves in dilute NaOH. When A is treated with Br_2 water it is converted into a compound $\text{C}_7\text{H}_5\text{OBr}_3$ rapidly. The structure of A is :


AROMATIC COMPOUNDS

521

158. Give the product of the following reaction sequence :



159. Give the product of the following reaction sequence:




160. Which represents an intermediate formed in the reaction of toluene and chlorine at elevated temperature in sunlight ?



161. The decreasing order of reactivity of *m*-nitrobromobenzene (I), 2, 4, 6- trinitrobromo-benzene (II), *p*-nitrobromobenzene (III), and 2,4-dinitrobromobenzene (IV), towards OH^- ions is :

- (a) $\text{I} > \text{II} > \text{III} > \text{IV}$
 (b) $\text{II} > \text{IV} > \text{III} > \text{I}$
- (c) $\text{IV} > \text{II} > \text{III} > \text{I}$
 (d) $\text{II} > \text{IV} > \text{I} > \text{III}$





162. Which one of the following compounds is most reactive for $\text{ArS}_{\text{N}}2$ reaction ?



522

ORGANIC Chemistry for IIT-JEE


163. Which one amongst the following carbocations is most stable ?



164. Cyclopentadiene is much more acidic than cyclopentane. The reason is that :

- (a) cyclopentadiene has conjugated double bonds
- (b) cyclopentadiene has both sp^2 and sp^3 hybridized carbon atoms
- (c) cyclopentadiene is a strain-free cyclic system
- (d) cyclopentadienide ion, the conjugate base of cyclopentadiene, is an aromatic species and hence has higher stability






165.



Friedel-Crafts acylation reaction can be used to obtain the compounds

- (a) II, III and IV
- (b) I, III and IV
- (c) I and II
- (d) II and III


166. The major product of the reaction is :




AROMATIC COMPOUNDS

523


167. The decreasing order of reactivity of given compound towards nucleophilic substitution with aqueous NaOH is :




(I)



(II)



(III)



(IV)






(a) I > II > III > IV

(b) II > IV > III > I


(c) IV > II > III > I

(d) II > IV > I > III





168. Identify the end product (B) of the following sequence of reactions.



169. Consider the following sequence of reactions :




The end product (C) is :



524


ORGANIC Chemistry for IIT-JEE

170. For the diazonium ions the order of reactivity towards diazo-coupling with phenol in the presence of dilute NaOH is :




- (a) $I < IV < II < III$
(b) $I < III < IV < II$
(c) $III < I < II < IV$
(d) $III < I < IV < II$

171. Major product obtained in given reaction is :



172.  $\xrightarrow[\text{or AlCl}_3]{\text{H}^+}$ (B) ; (A) & (B) are isomers. Product (B) is :
(A) (major)


Dewar's Benzene




AROMATIC COMPOUNDS

525


173. The step shown below is a recent synthesis of corannulene.



Product (A) is :




174.



Identify the position where E.A.S. will take place :

- (a) a (b) b
 (c) c (d) all the position are identical



The labelled carbon goes with :


- (a) $\text{Ph}-^{14}\text{CO}_2\text{H}$ (b) $^{14}\text{CO}_2$
 (c) $\text{Ph}-^{14}\text{CH}_2-\text{CO}_2\text{H}$ (d) $^{14}\text{CH}_4$

526


ORGANIC Chemistry for IIT-JEE

176. What is the expected order of reactivity of the following compounds in electrophilic chlorination ($\text{Cl}_2 + \text{FeCl}_3$) ?


(more reactive > less reactive)




I



II



III



IV






(a) I > II > III > IV

(b) IV > III > II > I

(c) III > I > IV > II

(d) II > III > I > IV

177. Which of the following is the major product from sulfonation of α -tetralone ?



178. Which of the following procedures would be best for the preparation of phenyl benzyl ether ?



(a) $\text{C}_6\text{H}_5\text{Cl} + \text{C}_6\text{H}_5\text{CH}_2\text{O}^{(-)}\text{Na}^{(+)}$

(b) $\text{C}_6\text{H}_5\text{O}^{(-)}\text{Na}^{(+)} + \text{C}_6\text{H}_5\text{CH}_2\text{Cl}$


(c) $2\text{C}_6\text{H}_5\text{Cl} + \text{Na}_2\text{O}$

(d) $2\text{C}_6\text{H}_5\text{MgBr} + \text{CH}_2\text{O}$

AROMATIC COMPOUNDS


527

179. Which of the following procedures would be best for achieving the following reaction ?







- (a) (i) KOH and heat (ii) CH₃C≡C—Br
 (b) (i) KMnO₄ and heat (ii) CH₃C≡C⁽⁻⁾Na⁽⁺⁾ (iii) excess H₂O
 (c) (i) NBS in CCl₄ and heat (ii) CH₃C≡C⁽⁻⁾Na⁽⁺⁾
 (d) (i) Mg in ether (ii) CH₃C≡CBr (iii) excess H₃PO₄

180. Which of the following procedures would be best for achieving the following reaction ?



- (a) (i) Br₂ + FeBr₃ (ii) KMnO₄ and heat (iii) HNO₃ and H₂SO₄
 (b) (i) KMnO₄ and heat (ii) Br₂ + FeBr₃ (iii) HNO₃ and H₂SO₄
 (c) (i) NBS in CCl₄ and heat (ii) KMnO₄ and heat (iii) HNO₃ and H₂SO₄
 (d) (i) NBS in CCl₄ and heat (ii) NaNO₂ and heat

181. Phenol reacts with acetone in the presence of conc. sulphuric acid to form a C₁₅H₁₆O₂ product. Which of the following compounds is this product ?




182. Heating benzene in a large excess of 80% D₂SO₄ in D₂O results in what product ?

- (a) C₆H₅SO₃D (b) C₆H₅OD (c) C₆H₅D (d) C₆D₆

183. A solution of cyclohexene in benzene is stirred at 0°C while concentrated sulphuric acid is added. After washing away the acid and removing the excess benzene, what product is isolated?

- (a) cyclohexylbenzene (b) 1-cyclohexylcyclohexene
 (c) *trans*-1,2-diphenylcyclohexane (d) 1,1-diphenylcyclohexane







184. Identify the reagents S and µ in the scheme below in which R is converted to the nitrite V via the benzylic halide T.




528

ORGANIC Chemistry for IIT-JEE





R, S and μ respectively are :

R	S	μ
(a) 	 (AlCl ₃)	HCN
(b) 	 (AlCl ₃)	HCN
(c) 	 (AlCl ₃)	KCN
(d) 	 (AlCl ₃)	KCN

185. Two aromatic compounds P and Q give product R.



Reactant P, Q and reagent used in above reaction are :

P	Q	Reagent
(a) 		AlCl ₃
(b) 		AlCl ₃
(c) 		AlCl ₃
(d) 		ZnCl ₂

AROMATIC COMPOUNDS

529


186. Which of the following C_6H_6 compounds has a single set of structurally equivalent hydrogen atoms ?



I




II



III







IV







V

- (a) I and II
(b) I and IV
(c) I and V
(d) I, II and III
187. Which of the following compounds would not be considered aromatic in its behaviour?
- (a)
- (b)
- (c)
- (d)
188. A C_8H_{10} hydrocarbon is nitrated by HNO_3 and sulphuric acid. Two, and only two, $C_8H_9NO_2$ isomers are obtained. Which of the following fits this evidence ?
- (a) ethyl benzene
(b) *ortho*-xylene
(c) *meta*-xylene
(d) *para*-xylene
189. Which of the following benzene ring substituents is deactivating but *ortho-para* directing ?
- (a) $-N=O$
(b) $-OCH_3$
(c) $-COCH_3$
(d) $-NO_2$
190. Which of the following compounds forms *ortho*-benzoic acid when oxidized by hot aqueous potassium permanganate ?



191. Which of the following organic chlorides will not give a Friedel-Craft alkylation product when heated with benzene and $AlCl_3$?
- (a) $(CH_3)_3CCl$
(b) $CH_2=CHCH_2Cl$
(c) CH_3CH_2Cl
(d) $CH_2=CHCl$
192. Which of the following is aromatic ?




530

ORGANIC Chemistry for IIT-JEE

193. Which of the following substance will increase the acidity of phenol ?


- (a) Dil. H_2SO_4 (b) Dil. HCl (c) Conc. H_2SO_4 (d) Conc. CH_3COOH

194.  + $\text{PhMgBr} \longrightarrow E + F$



Pyrrole

$E + \text{MeCl} \longrightarrow G + H$


$F + \text{MeCl} \longrightarrow$ no reaction without a catalyst




1




2




3




4



5




6






7

The structure of products $E - H$, respectively are

- (a) 3, 2, 6, 7 (b) 4, 5, 6, 1
(c) 3, 4, 5, 2 (d) 3, 2, 4, 5


195.  $\xrightarrow{\text{H}^+} (A)$; Product A is :




(d) none of these

AROMATIC COMPOUNDS


531




Identify the product of the above rearrangement reaction.




197. Product obtained in the following transformation is :




198. The compound X in the reaction.



532

ORGANIC Chemistry for IIT-JEE

- (a) Nucleophilic addition
(b) Nucleophilic substitution
(c) Electrophilic addition
(d) Electrophilic substitution
(e) Free radical substitution



- (a) Nucleophilic addition
(b) Nucleophilic substitution
(c) Electrophilic addition
(d) Electrophilic substitution
(e) Free radical substitution

AROMATIC COMPOUNDS**533****ANSWERS — LEVEL 1**

1.	(b)	2.	(d)	3.	(b)	4.	(c)	5.	(a)	6.	(b)	7.	(c)	8.	(b)
9.	(d)	10.	(d)	11.	(a)	12.	(a)	13.	(a)	14.	(a)	15.	(a)	16.	(b)
17.	(b)	18.	(a)	19.	(b)	20.	(b)	21.	(c)	22.	(c)	23.	(b)	24.	(b)
25.	(d)	26.	(c)	27.	(c)	28.	(b)	29.	(b)	30.	(b)	31.	(b)	32.	(a)
33.	(b)	34.	(d)	35.	(c)	36.	(c)	37.	(a)	38.	(c)	39.	(a)	40.	(b)
41.	(b)	42.	(c)	43.	(c)	44.	(b)	45.	(c)	46.	(b)	47.	(b)	48.	(b)
49.	(b)	50.	(b)	51.	(b)	52.	(a)	53.	(c)	54.	(c)	55.	(a)	56.	(a)
57.	(d)	58.	(b)	59.	(b)	60.	(b)	61.	(c)	62.	(b)	63.	(b)	64.	(b)
65.	(a)	66.	(a)	67.	(c)	68.	(b)	69.	(a)	70.	(d)	71.	(b)	72.	(a)
73.	(a)	74.	(c)	75.	(a)	76.	(d)	77.	(c)	78.	(b)	79.	(c)	80.	(b)
81.	(b)	82.	(c)	83.	(a)	84.	(c)	85.	(d)	86.	(b)	87.	(d)	88.	(b)
89.	(d)	90.	(d)	91.	(d)	92.	(d)	93.	(d)	94.	(b)	95.	(c)	96.	(d)
97.	(b)	98.	(d)	99.	(b)	100.	(b)	101.	(b)	102.	(c)	103.	(d)	104.	(a)
105.	(b)	106.	(d)	107.	(c)	108.	(b)	109.	(b)	110.	(c)	111.	(c)	112.	(b)
113.	(c)	114.	(a)	115.	(b)	116.	(b)	117.	(c)	118.	(c)	119.	(b)	120.	(c)
121.	(b)	122.	(b)	123.	(c)	124.	(b)	125.	(b)	126.	(b)	127.	(c)	128.	(c)
129.	(c)	130.	(b)	131.	(d)	132.	(b)	133.	(b)	134.	(a)	135.	(b)	136.	(a)
137.	(b)	138.	(c)	139.	(d)	140.	(b)	141.	(b)	142.	(c)	143.	(a)	144.	(b)
145.	(b)	146.	(c)	147.	(a)	148.	(c)	149.	(c)	150.	(b)	151.	(d)	152.	(c)
153.	(c)	154.	(b)	155.	(c)	156.	(c)	157.	(c)	158.	(b)	159.	(c)	160.	(c)
161.	(b)	162.	(c)	163.	(d)	164.	(d)	165.	(c)	166.	(c)	167.	(b)	168.	(d)
169.	(d)	170.	(b)	171.	(b)	172.	(a)	173.	(a)	174.	(b)	175.	(b)	176.	(d)
177.	(b)	178.	(b)	179.	(c)	180.	(a)	181.	(b)	182.	(d)	183.	(a)	184.	(c)
185.	(c)	186.	(b)	187.	(b)	188.	(b)	189.	(a)	190.	(b)	191.	(d)	192.	(b)
193.	(c)	194.	(a)	195.	(c)	196.	(b)	197.	(b)	198.	(b)	199.	(d)	200.	(e)

534

ORGANIC Chemistry for IIT-JEE



1. Each of the six compounds shown at the bottom of the page has two aromatic (benzene) rings. In each case the two rings are different and are labeled A & B. If an **electrophilic substitution, such as nitration or bromination**, is carried out on each compound, then identify which ring (A or B) will be preferentially attacked, and indicate the orientation of the substitution (ortho/para, meta or all sites).

Compound	Reactivity	Substitution	Compound	Reactivity	Substitution
1.	A	ortho/para	2.	A	ortho/para
	B	meta		B	meta
		all sites			all sites
3.	A	ortho/para	4.	A	ortho/para
	B	meta		B	meta
		all sites			all sites
5.	A	ortho/para	6.	A	ortho/para
	B	meta		B	meta
		all sites			all sites

Compound		Compound	
1.		2.	
3.		4.	
5.		6.	

AROMATIC COMPOUNDS**535**

2. When given substituents on a benzene ring, as activating or de-activating and as ortho-para or meta directing for electrophilic aromatic substitution fill the following by appropriate (✓) right or (X) wrong.

	Substituent	Activating	De-activating	Ortho/para	Meta
1.	$-\text{OCH}_3$				
2.	$\begin{array}{c} \text{O} \\ \\ -\text{C}-\text{O}-\text{CH}_3 \end{array}$				
3.	$\begin{array}{c} -\text{O}-\text{C}-\text{CH}_3 \\ \\ \text{O} \end{array}$				
4.	$-\text{CH}_3$				
5.	$-\text{F}$				
6.	$-\text{Ph}$				
7.	$\begin{array}{c} \text{O} \\ \\ -\text{NH}-\text{C}-\text{CH}_3 \end{array}$				
8.	$\begin{array}{c} \text{O} \\ \\ -\text{C}-\text{NH}-\text{CH}_3 \end{array}$				
9.	$-\text{Br}$				
10.	$-\text{CN}$				
11.	$-\text{CF}_3$				
12.	$\begin{array}{c} \text{O} \\ \\ -\text{C}-\text{NH}_2 \end{array}$				
13.	$\begin{array}{c} \text{O} \\ \\ -\text{C}-\text{OH} \end{array}$				
14.	$-\text{CH}=\text{CH}_2$				

536

ORGANIC Chemistry for IIT-JEE

15.	$\begin{array}{c} \text{O} \\ \\ -\text{CH} = \text{CH} - \text{C} - \text{OH} \end{array}$				
16.	$\begin{array}{c} \text{O} \\ \\ -\text{CH} = \text{CH} - \text{C} - \text{H} \end{array}$				
17.	$-\text{S} - \text{Et}$				
18.	$\begin{array}{c} -\text{S} - \text{Et} \\ \\ \text{O} \end{array}$				
19.	$\begin{array}{c} \text{O} \\ \\ -\text{S} - \text{Et} \\ \\ \text{O} \end{array}$				
20.	$-\text{N} = \text{O}$				
21.	$-\text{CH}_2\text{NH}_2$				
22.	$-\text{CHX}_2$				

3. Devise a series of reactions to convert benzene into *meta*-chlorobromobenzene.

Select reagents and conditions from the following table, listing them in the order of use.

Compound		Compound		Compound	
1.	sulphuric acid (conc.) heat	5.	Mg in ether	9.	$\text{Cu}_2\text{Br}_2 + \text{HBr}$
2.	$\text{Cl}_2 + \text{FeCl}_3$ and heat	6.	PBr_3	10.	$(\text{CH}_3\text{CO})_2\text{O} + \text{Pyridine}$
3.	$\text{NaNO}_2 + \text{H}_3\text{O}^{(+)} \text{ } 0^\circ\text{C}$	7.	H_3PO_2		
4.	H_2 Pt catalyst	8.	$\text{HNO}_3(\text{conc.}) + \text{H}_2\text{SO}_4(\text{conc.})$ and heat		



(a) 1 then 2 then 6

(b) 2 then 8 then 4 then 3 then 9

(c) 8 then 4 then 10 then 2 then 3 then 9

(d) 8 then 2 then 4 then 3 then 9

AROMATIC COMPOUNDS**537****4. Match the Column (I) and Column (II). (Matrix)**

Column (I)		Column (II)	
(a)		(p)	Aromatic
(b)		(q)	$(4n + 2)\pi$ electron in a single ring
(c)	$\text{Fe}(\text{C}_5\text{H}_5)_2$	(r)	4π electron in a single ring
(d)	$\text{Cr}(\text{C}_6\text{H}_6)_2$	(s)	Effective atomic number of metal = 36


5. Match the Column (I) and Column (II).

Column (I)		Column (II)	
Compound (Monocyclic)		Number of π - electron	
(a)	$\text{C}_4\text{H}_4^{-2}$	(p)	$2\pi e$
(b)	$\text{C}_4\text{H}_4^{+2}$	(q)	$6\pi e$
(c)	$\text{C}_9\text{H}_9^{+1}$	(r)	$8\pi e$
(d)	$\text{C}_9\text{H}_9^{-1}$	(s)	$10\pi e$

538

ORGANIC Chemistry for IIT-JEE

6. Match the Column (I), Column (II) and Column (III). (Matrix)

Column I		Column II		Column III	
(a)		(p)	Aromatic	(w)	$(4n + 2)\pi$ electron. $n = 0, 1, 2, 3$
(b)		(q)	Non-aromatic	(x)	$4n\pi$ electron $n = 1, 2, 3$
(c)		(r)	Anti-aromatic	(y)	Non-planar compound
(d)		(s)	Planar compound	(z)	Readily reacts with active metal






7. Match the Column (I), Column (II) and Column (III). (Matrix)

Column I		Column II		Column III	
(a)		(p)	Readily react with active metal	(w)	Aromatic
(b)		(q)	Readily undergo Dimerization at room temperature	(x)	Anti-aromatic
(c)		(r)	$(4n + 2)\pi$ electron $n = 0, 1, 2, 3$	(y)	Non-aromatic
(d)		(s)	$4n\pi$ electron	(z)	High dipole

AROMATIC COMPOUNDS

539

8. Among the following compound.

Compound		Compound		Compound	
(a)		(b)		(c)	
(d)		(e)		(f)	
(g)	$C_8H_8^{-2}$	(h)	$C_3H_3^{+}$	(i)	
(j)		(k)		(l)	

- (a) Number of compounds which are aromatic = P
 (b) Number of compounds which are anti-aromatic = Q
 (c) Number of compounds which are non-aromatic = R
 (d) Number of compounds which readily = S
 Undergo dimerization at room temperature
 (e) Number of compound which reacts with active metal = T
 Sum of $P + Q + R + S + T =$





9. Of the following compounds which will react with Br_2 at room temperature in dark.

(a)	Benzene (C_6H_6)
(b)	Cyclohexene (C_6H_{10})
(c)	Cyclohexane (C_6H_{12})
(d)	Propanoic Acid ($C_2H_5CO_2H$)
(e)	Phenol (C_6H_5OH)
(f)	Nitrobenzene ($C_6H_5NO_2$)
(g)	Hexyne (C_6H_{10})
(h)	2,2-dichloropropane ($C_3H_6Cl_2$)

540

ORGANIC Chemistry for IIT-JEE

10. Among the following compound.







Compound		Compound		Compound	
(a)		(b)	$C_8H_8^{-2}$	(c)	
(d)		(e)		(f)	
(g)		(h)		(i)	$C_3H_3^{+1}$
(j)		(k)		(l)	

- (a) Number of compounds which are aromatic = w
 (b) Number of compounds which are non-aromatic = x
 (c) Number of compounds which are anti-aromatic = y
 (d) Number of compounds which readily undergo Dimerization at room temperature = z
 Sum of w + x + y + z = ...

AROMATIC COMPOUNDS

541

11. Complete the following table.


Reactant	Reagents(s)/Conditions	Major Organic Products
(a) 	(A)	
(b) 	1. NaNO ₂ in dilute H ₂ SO ₄ /0 – 5°C 2. heat or boiling	(B)
(c) 	SO ₃ /conc. H ₂ SO ₄	(C)
(d) (D)	1. NaOH heated at 330°C 2. dilute H ₃ O ⁺	 and 
(e) 	1. aqueous NaOH heated at 60°C 2. dilute H ₃ O ⁺	(E)

542


ORGANIC Chemistry for IIT-JEE

12. Comprehension


Given is the energy profile diagram of nitration of benzene using mixed acid. ($\text{HNO}_3 + \text{H}_2\text{SO}_4$)




A. Identify (x) in above reaction :



B. Identify T.S₁ in the above reaction :













C. Identify T.S₂ in the above reaction :



AROMATIC COMPOUNDS

543

13. Examine the ten structural formulas shown below and select those that satisfy each of the following conditions. Enter one or more letters (a through j) in each answer box, reflecting your choice for each.







Compound		Compound	
a.		b.	
c.		d.	
e.		f.	
g.		h.	
i.		j.	

A.	Which compounds undergo electrophilic nitration more rapidly than benzene ?	
B.	Which compounds give meta substitution under electrophilic bromination conditions ?	

544

ORGANIC Chemistry for IIT-JEE

14. Nitrobenzene is a versatile compound that may be converted into a wide variety of substituted benzenes. Five such synthesis are shown below. In each reaction box above an arrow write letters designating the reagents and conditions, selected from the list at the bottom of the page, that would effect the transformation. The reagents must be written in the answer box in the correct order of their use. You may assume appropriate heating or cooling takes place, and more than one equivalent of the reagent may be used if needed.



Reactant	Reagent		Product
 nitrobenzene	a.	→	v.
	b.	→	w.
	c.	→	x.
	d.	→	y.
	e.	→	z.
			
			
			
			
			

Reagents		Reagents	
A.	H ₂ , Ni catalyst	F.	Cl ₂ & FeCl ₃
B.	KBr & Cu ₂ Br ₂	G.	NaOH 10% solution
C.	KCN & Cu ₂ (CN) ₂	H.	(CH ₃ CO) ₂ O, pyridine
D.	HNO ₂ 0°C	I.	HNO ₃ /H ₂ SO ₄
E.	CH ₃ I & pyridine		

AROMATIC COMPOUNDS**545****15. Match the column I and II.**

Column (I)		Column (II)	
Group		Effect on phenyl ring	
(a)	$-\text{CH}=\text{CH}-\text{CO}_2\text{H}$	(p)	<i>o/p</i> -directors
(b)	$-\text{O}-\overset{\text{O}}{\parallel}{\text{S}}-\text{CH}_3$	(q)	meta-directors
(c)	$-\text{NH}-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_3$	(r)	Activating group
(d)	$-\overset{\text{O}}{\parallel}{\text{S}}-\text{CH}_3$	(s)	De-activating group

16. Match the column I and II.


Column (I)		Column (II)	
Group		Effect on phenyl group	
(a)		(p)	Activating group
(b)		(q)	De-activating group

546


ORGANIC Chemistry for IIT-JEE

(c)		(r)	<i>o/p</i> -director
(d)		(s)	meta-director

17. Comprehension

Nucleophilic Aromatic substitution ($\text{S}_{\text{N}}\text{Ar}$) :A substituted benzene derivative containing- NO_2 and Cl group at *p*-position is subjected to Nu-substitution.

A. Match the column I and II :



Column (I)		Column (II)	
$\text{X} = \text{halogen}$		relative reactivity toward ($\text{S}_{\text{N}}\text{Ar}$).	
(a)	- F	(p)	312
(b)	- Cl	(q)	1
(c)	- Br	(r)	0.8
(d)	- I	(s)	0.6















AROMATIC COMPOUNDS

547

B. If step-2 were rate determining step, which halogen of aryl halide is most reactive toward SN_{Ar} .

- (a) Fluoride (b) Chloride (c) Bromide (D) Iodide






C. Which of the following is most reactive toward SN_{Ar} .







548

ORGANIC Chemistry for IIT-JEE


F. The cumulative effect of their fluorine activate the rings of penta and hexa fluorobenzene toward nucleophilic aromatic substitution. What is compound X in the following synthesis ?



G. Which is the best route for the synthesis of starting from benzene of?



18. Identify product (A) and write its structure.




AROMATIC COMPOUNDS

549


SUBJECTIVE PROBLEMS

1.



Double bond equivalent of D is :

2. How many isomers 'x' of C_8H_{10} when reacts with hot alkaline KMnO_4 give only aromatic dicarboxylic acid ? How many isomers 'y' of C_4H_8 when reacts with hot alkaline KMnO_4 give carbondioxide ?
Sum of $x+y=?$
3. How many groups are *o/p* director in the electrophilic aromatic substitution ?



550

ORGANIC Chemistry for IIT-JEE

ANSWERS — LEVEL 2

1.

Compound	Reactivity	Substitution
1	B	ortho/para
2	A	ortho/para
3	B	ortho/para
4	A	ortho/para
5	B	meta
6	B	ortho/para

2.

	Substituent	Activating	De-activating	Ortho/para	Meta
1.	$-\text{OCH}_3$	✓	✗	✓	✗
2.	$-\text{C}(=\text{O})\text{OCH}_3$	✗	✓	✗	✓
3.	$-\text{O}-\text{C}(=\text{O})\text{CH}_3$	✓	✗	✓	✗
4.	$-\text{CH}_3$	✓	✗	✓	✗
5.	$-\text{F}$	✗	✓	✓	✗
6.	$-\text{Ph}$	✓	✗	✓	✗
7.	$-\text{NH}-\text{C}(=\text{O})\text{CH}_3$	✓	✗	✓	✗
8.	$-\text{C}(=\text{O})\text{NH}-\text{CH}_3$	✗	✓	✗	✓
9.	$-\text{Br}$	✗	✓	✓	✗
10.	$-\text{CN}$	✗	✓	✗	✓

AROMATIC COMPOUNDS

551

11.	$-\text{CF}_3$	X	✓	X	✓
12.	$\begin{array}{c} \text{O} \\ \\ -\text{C}-\text{NH}_2 \end{array}$	X	✓	X	✓
13.	$\begin{array}{c} \text{O} \\ \\ -\text{C}-\text{OH} \end{array}$	X	✓	X	✓
14.	$-\text{CH}=\text{CH}_2$	✓	X	✓	X
15.	$\begin{array}{c} \text{O} \\ \\ -\text{CH}=\text{CH}-\text{C}-\text{OH} \end{array}$	X	✓	✓	X
16.	$\begin{array}{c} \text{O} \\ \\ -\text{CH}=\text{CH}-\text{C}-\text{H} \end{array}$	X	✓	✓	X
17.	$-\text{S}-\text{Et}$	✓	X	✓	X
18.	$\begin{array}{c} -\text{S}-\text{Et} \\ \\ \text{O} \end{array}$	X	✓	✓	X
19.	$\begin{array}{c} \text{O} \\ \\ -\text{S}-\text{Et} \\ \\ \text{O} \end{array}$	X	✓	X	✓
20.	$-\text{N}=\text{O}$	X	✓	✓	X
21.	$-\text{CH}_2\text{X}$	X	✓	X	✓
22.	$-\text{CHX}_2$	X	✓	X	✓

3. d

4. a - p, q; b - p, q; c - p, q, s; d - p, q, s

5. a - q; b - p; c - r; d - s

6. a - p, s - w; b - p, s - w; c - q - x, y, z; d - q - w - y

7. a - p, q, s - x; b - p - s - y; c - r - w, z; d - r - w, z


8. $P + Q + R + S + T = 19$

9. b, e, g

10. $w + x + y + z = 14$

552

ORGANIC Chemistry for IIT-JEE



12. A - a; B - b; C - c

13. A - a, b, e, f, h; B - c, d, g, i (Note : yet C_6H_5Br is less reactive than C_6H_6 but *o/p* directing)


14. v - F, A, D, B; w - A, H, I, G, D, C; x - A, E, D; y - F, I, A or I, F, A;

z - A, H, I, F, G, A or A, H, I, F, A, G

15. a - p, s; b - p, r; c - p, r; d - p, s

16. a - p, r; b - p, r; c - p, r; d - p, r

17. A - a - p, b - q, c - r, d - s; B - d; C - d; D - a; E - b; F - c; G - a



Subjective Problems



1. 9

2. 5

3. 6

13 PRACTICAL ORGANIC CHEMISTRY



LEVEL-1

1.  and ; Compounds (X) and (Y) can be differentiated by :
- (X) (Y)

- (a) $\text{H}_3\text{O}^{\oplus}$, NaOI (b) $\text{H}_3\text{O}^{\oplus}$, then Fehling test
(c) $\text{H}_3\text{O}^{\oplus}$, then Na (d) Both (b) and (c)

2. Compound $\text{CH}_3 - \text{CH} \begin{matrix} \text{OEt} \\ \text{OEt} \end{matrix}$ and $\text{CH}_3 - \text{CH}_2 - \text{O} - \text{CH}_2 - \text{CH}_3$ can be differentiated by :
(P) (Acetal) (Q)

- (a) $\text{H}_3\text{O}^{\oplus}$, Na (b) $\text{H}_3\text{O}^{\oplus}$, Tollens' test
(c) $\text{H}_3\text{O}^{\oplus}$, Fehling test (d) All of these


3.  and  can be differentiated by :
- (aniline) (cyclohexyl amine)

- (a) Hinsberg test (b) Iso-cyanide test
(c) NaNO_2 , HCl, then β -Naphthol (d) NaOH


554

ORGANIC Chemistry for IIT-JEE


4.



(p-ethyl phenol)



(p-methyl anisole)




(p-ethyl benzyl alcohol)


Above compounds can be differentiated by using the reagent:

(a) NaOH, Tollen's reagent, FeCl_3 (b) CrO_3 , Tollen's reagent, FeCl_3 (c) Tollen's reagent, CrO_3 , FeCl_3 (d) Na, Tollen's reagent, FeCl_3


5.




Salicyclic acid



Ethyl salicylate



Ethyl acetyl




acetyl salicyclic acid


Above compounds can be differentiated by the salicylate. Which of the following chemical test? (used in decreasing order)

(a) NaOH, FeCl_3 , NaHCO_3 (b) aq. NaHCO_3 , FeCl_3 , NaOH(c) NaOI, NaOH, NaHCO_3 (d) NaOH, Na, NaHCO_3


6.




(m-nitrophenol)



(m-nitro benzoic acid)



(m-nitro aniline)




(m-dinitro benzene)


Above compounds can be differentiated by which of the following chemical test ? (used in decreasing order)

(a) NaOH, NaHCO_3 , HCl(b) HCl, NaOH, NaHCO_3 (c) NaHCO_3 , NaOH, HCl(d) NaOH, HCl, NaHCO_3


7.




Product (A) in the above reaction is :




(a)



(b)




(c)



(d)

PRACTICAL ORGANIC CHEMISTRY




555




Above compounds (P) & (Q) can be differentiated by :

- (a) amm. AgNO_3 (b) NaOH
(c) FeCl_3 (d) Both (a) & (b)

9. Which of following acid give positive Tollen's reagent test.

- (a) $(\text{CH}_2)_2$  (b) CH_2  (c)  (d) HCO_2H

10. Which of following compounds give positive Tollen's test?



11. Give a simple test to differentiate cyclohexane and cyclohexene

- (a) $\text{Br}_2/\text{H}_2\text{O}$ (b) Bayer's reagent
(c) Tollen's reagent (d) Both (a) and (b)

12. Give test to differentiate Bromobenzene and Benzyl bromide (PhCH_2Br).

- (a) (i) aq. KOH (ii) Na (c) KMnO_4 (b) AgNO_3 (d) All these

13. Give test to differentiate 1,1-dichloroethane and 1, 2-dichloroethane :

- (a) 2,4 -DNP then aq. KOH (b) aq. KOH then 2, 4-DNP
(c) NaHSO_3 (d) Lucas reagent

14. Test to differentiate between (CH_3OH) and $(\text{Ph} - \text{OH})$ is/are :
(methanol) (Phenol)

- (a) Litmus test (b) FeCl_3
(c) $\text{Br}_2/\text{H}_2\text{O}$ (d) All of these

15. Acetaldehyde and benzaldehyde can be differentiated by :

- (a) Fehling test (b) Iodoform test
(c) Tollen's reagent (d) both (a) and (b)

16. Ethylamine and diethylamine cannot be differentiated by :

- (a) Hinsberg test (b) carbylamine test
(c) Iodoform test (d) both (a) and (b)

17. Lassaigne's test for the detection of nitrogen will fail in the case of :

- (a) NH_2CONH_2 (b) $\text{NH}_2\text{CONHNH}_2 \cdot \text{HCl}$
(c) $\text{NH}_2\text{NH}_2 \cdot \text{HCl}$ (d) $\text{C}_6\text{H}_5\text{NHNH}_2 \cdot 2\text{HCl}$

556**ORGANIC Chemistry for IIT-JEE**

18. Sodium nitroprusside when added to an alkaline solution of sulphide ions produces a colouration which is :
 (a) red (b) blue
 (c) brown (d) purple
19. In Kjeldahl's method, nitrogen present is estimated as :
 (a) N_2 (b) NH_3
 (c) NO_2 (d) none of these
20. In Kjeldahl's method of estimation of nitrogen, K_2SO_4 acts as :
 (a) an oxidising agent (b) catalytic agent
 (c) hydrolysing agent (d) boiling point elevator
21. The prussian blue colour obtained during the test of nitrogen by Lassaigne's test is due to the formation of :
 (a) $Fe[Fe(CN)_6]_3$ (b) $Na_3[Fe(CN)_6]$
 (c) $Fe(CN)_3$ (d) $Na_4[Fe(CN)_5NOS]$
22. A compound which does not give a positive test in Lassaigne's test for nitrogen is:
 (a) urea (b) hydrazine (c) azobenzene (d) phenyl hydrazine
23. *p*-nitrophenol and *o*-nitrophenol are separated by :
 (a) distillation (b) steam distillation
 (c) crystallization (d) fractional crystallization
24. Which of the following reagent is used for the separation of acetaldehyde from acetophenone ?
 (a) NH_2OH (b) $NaOH$
 (c) Tollen's reagent (d) $C_6H_5NHNH_2$
25. The formula of gas is $[CO]_x$. If its vapour density is 70, the value of x will be :
 (a) 2.5 (b) 3.0
 (c) 5.0 (d) 6.0

ANSWERS — LEVEL 1

1.	(d)	2.	(d)	3.	(c)	4.	(b)	5.	(b)	6.	(c)	7.	(a)	8.	(d)
9.	(d)	10.	(c)	11.	(d)	12.	(d)	13.	(b)	14.	(d)	15.	(d)	16.	(c)
17.	(c)	18.	(b)	19.	(b)	20.	(d)	21.	(d)	22.	(b)	23.	(a)	24.	(c)
25.	(c)														

PRACTICAL ORGANIC CHEMISTRY


557




LEVEL-2

1. Comprehension


Given are the isomers of $C_8H_8O_2$.




(a)




(b)




(c)



(d)




(e)



(f)


- A. Which isomer gives positive iodoform test ?
 (a) a (b) b
 (c) d (d) c
- B. Which isomer gives +ve Tollens' test, also reacts with $FeCl_3$?
 (a) b (b) f
 (c) c (d) d
- C. Which isomer reacts with $NaHCO_3$?
 (a) c (b) d
 (c) e (d) f
- D. Which isomer on hydrolysis gives 1, 4-di hydroxybenzene ?
 (a) a (b) d
 (c) e (d) f



Sum of molecular mass of gas ($A+B=?$)

ANSWERS — LEVEL 2

1. A - d; B - b; C - a; D - b
 2. 48



14 BIOMOLECULES





LEVEL - 1

1. Which statement correctly completes the statement ?


Except for glycine, which is achiral, all the amino acids present in proteins

- (a) are chiral, but racemic
- (b) have the L configuration at their α carbon
- (c) have the R configuration at their α carbon
- (d) have the S configuration at their α carbon

2. Assume that a particular amino acid has an isoelectric point of 6.0. In a solution at pH 1.0, which of the following species will predominate ?



3. The pK_a values for the three ionizable groups X, Y and Z of glutamic acid are 4.3, 9.7 and 2.2 respectively






The isoelectric point for the amino acid is :

- (a) 7.00
- (b) 3.25
- (c) 4.95
- (d) 5.95


BIOMOLECULES

559

4. An amino acid may be represented by general formula $\text{H}_2\text{N}-\text{CH}(\text{R})-\text{COOH}$. If $\text{R} = -\text{CH}_2\text{C}_6\text{H}_5$ then it is phenylalanine (Phe) and if $\text{R} = \text{CH}_3$ then it is alanine (Ala). Find the sequence of reagents from those given below to synthesize Phe-Ala.




- (a) 1 and 2 (b) 1 and 4 (c) 2 and 3 (d) 3 and 4
5. Iso-electric point of alanine is ($\text{pH} = 6$). At which pH , maximum concentration of zwitter ion of alanine will be present ?
- (a) $\text{pH} > 6$ (b) $\text{pH} < 6$ (c) $\text{pH} = 6$ (d) $\text{pH} = 7$



Identify the amino acid obtained by hydrolysis of the above compound:


- (a) Glycine (b) Alanine (c) Both (a) and (b) (d) None of these
7. At iso-electric point
- (a) conc. of cation is equal to conc of anion
- (b) Net charge is zero.
- (c) Maximum conc. of di-polar ion (Zwitter ion) will be present
- (d) All of the above
8. Which of following amino acid has lowest iso-electric point ?
- (a) Glycine (b) Alanine (c) Aspartic acid (d) Lysine
9. Find iso-electric point of given amino acid



$$\text{p}K_b = 4.4$$

- (a) 3.3 (b) 5.9 (c) 9.6 (d) 11.8

10. Find iso-electric point of the given amino acid



- (a) 5.5 (b) 6.5 (c) 3 (d) 5

560

ORGANIC Chemistry for IIT-JEE

11. $\text{H}-\text{C}\equiv\text{C}-\text{H} \xrightarrow[\text{H}_2\text{SO}_4]{\text{HgSO}_4} (\text{A}) \xrightarrow[(2) \text{H}_3\text{O}^+]{(1) \text{NH}_3+\text{HCN}} (\text{B})$; Product (B) of given reaction is :

- (a) Glycine (b) Alanine
(c) valine (d) Leucine

12. Which amino acid does not contain chiral centre ?

- (a) Valine (b) Leucine (c) Glycine (d) Iso-leucine


13. Which of the following is Sanger reagent ?

- (a) 2,4-Di-nitro fluorebenzene (b) Phenyl isocyanate
(c) 2, 4-Di-nitro chlorobenzene (d) 2, 4-Di-nitro iodobenzene

14. A D-carbohydrate is :

- (a) Always dextrorotatory
(b) Always laevorotatory
(c) Always the mirror of the corresponding L-carbohydrate
(d) None of these

15. Which L-sugar on oxidation gives an optically active dibasic acid (2 COOH groups) ?




16. $\begin{array}{c} \text{CH}=\text{N}-\text{NH}-\text{Ph} \\ | \\ \text{C}=\text{N}-\text{NH}-\text{Ph} \\ | \\ \text{HO}-\text{C}-\text{H} \\ | \\ \text{H}-\text{C}-\text{OH} \\ | \\ \text{H}-\text{C}-\text{OH} \\ | \\ \text{CH}_2\text{OH} \end{array}$ The given osazone can be obtained by :

- (a) D-glucose (b) D-mannose (c) D-Idose (d) Both (a) & (b)

17. Which of the following pair gives same phenyl osazone ?

- (a) D-Glucose and D-Allose (b) D-Glucose and D-Alfrose
(c) D-Glucose and D-Mannose (d) D-Glucose and D-Talose


18. Which of the following is the Fischer projection of L-threonine (also known as (2S, 3R)-2-amino-3-hydroxybutanoic acid) ?



BIOMOLECULES

561



19. Among the three compounds shown below, two yield the same product on reaction with warm HNO_3 . The exception is :




20. The optical rotation of the α -form of a pyranose is $+150.7^\circ$, that of the β -form is $+52.8^\circ$. In solution an equilibrium mixture of these anomers has an optical rotation of $+80.2^\circ$. The percentage of the α -form in equilibrium mixture is :

(a) 28% (b) 32% (c) 68% (d) 72%

21. Which of the following represents the anomer of the compound shown ?



22. Which set of terms correctly identifies the carbohydrate shown ?



- (1) Pentose (2) Pentulose (3) Hexulose (4) Hexose
(5) Aldose (6) Ketose (7) Pyranose (8) Furanose
(a) 2, 6, 8 (b) 2, 6, 7
(c) 1, 5, 8 (d) A set of terms other than these

23. For the complex conversion of D-glucose into the corresponding osazone, the minimum number of equivalents of phenyl hydrazine required is :

(a) two (b) three (c) four (d) five


24. Which one of the following compounds will form an osazone derivative ?

- (a) $\text{CH}_3\text{CH}_2\text{COCH}_2\text{OH}$ (b) $\text{CH}_3\text{COCH}_2\text{CH}_2\text{OH}$
(c) $\text{CH}_3\text{CH}_2\text{CHOHCH}_2\text{OH}$ (d) $\text{CH}_3\text{CH}_2\text{COCH}_2\text{OCH}_3$

562

ORGANIC Chemistry for IIT-JEE

25. Which of the following structure is L-arabinose?




26. Which one of the statements concerning the equilibrium shown is true ?



- (a) The two structures are enantiomers of each other. They have equal but opposite optical rotations and racemize slowly at room temperature
- (b) The two structures are enantiomers of each other. They racemize too rapidly at room temperature for their optical rotations to be measured
- (c) The two structures are diastereomers of each other. Their interconversion is called mutarotation
- (d) The two structures are diastereomers of each other. Their interconversion does not require breaking and making bonds, only a change in conformation

27. The configurations of the chiral centres in D-threose (shown), are :




- (a) 2R, 3R (b) 2R, 3S (c) 2S, 3R (d) 2S, 3S

28. Rapid interconversion of α -D-glucose and β -D-glucose in solution is known as :

- (a) racemization (b) asymmetric induction
- (c) fluxional isomerization (d) mutarotation

29. Identify the correct set of stereochemical relationships amongst the following monosaccharides I-IV.




- (a) I and II are anomers ; III and IV are epimers
- (b) I and III are epimers ; II and IV are anomers
- (c) I and II are epimers ; III and IV are anomers
- (d) I and III are anomers ; I and II are epimers


BIOMOLECULES

563

30. What is the structure of L-glucose ?



31. What is the structure of L-glyceraldehyde?



32.
$$\begin{array}{c} \text{HC} - \text{OH} \\ || \\ \text{C} - \text{OH} \\ | \\ \text{HO} - \text{C} - \text{H} \\ | \\ \text{H} - \text{C} - \text{OH} \\ | \\ \text{H} - \text{C} - \text{OH} \\ | \\ \text{CH}_2\text{OH} \end{array}$$
, the given is an isomer of.

- (a) D-glucose (b) D-mannose (c) D-fructose (d) All of these

33. D-glucose $\xrightarrow{\text{HO}^-}$ A + B; A and B are :

- (a) D-mannose & D-mannitol (b) D-mannose & D-Fructose
(c) D-Allose & D-Altrose (d) D-Glucose & D-Idose

34. Stereoisomers of aldohexose is (a) and stereoisomers of ketohexose is (b).

Ratio of a/b is :

- (a) $\frac{1}{2}$ (b) $\frac{2}{1}$ (c) $\frac{4}{1}$ (d) $\frac{1}{4}$

35. D-Glucose $\xrightarrow[\Delta]{\text{HNO}_3}$ (A); Product (A) is :

- (a) D-Gluconic acid (b) D-Glucitol (c) D-Fructose (d) D-Glucaric acid

36. D-glucose & D-fructose can be differentiated by :


- (a) Fehling solution (b) Tollens reagent
(c) Benedict test (d) $\text{Br}_2/\text{H}_2\text{O}$

37. D-Glucose exist in x different forms. The value of x (stereoisomer) is :

- (a) 2 (b) 3
(c) 4 (d) 5

564







ORGANIC Chemistry for IIT-JEE



Product (A) of above reaction is :

- (a) D-glucose (b) D-fructose
(c) D-talose (d) D-idose

39. Which of the molecules below will react with Ag^+ ?




- (a) (i), (iii) and (v) (b) (ii) and (iv)
(c) (iv) and (vi) (d) (i), (ii), (iii) and (vi)


BIOMOLECULES

565


40. A.




B.



C.



D.



Which of the compounds (A-D) depicted above is NOT a hemiacetal linkage ?

- (a) Compound A (b) Compound B
(c) Compound C (d) Compound D
(e) None of the above (they are all hemiacetals)

ANSWERS — LEVEL 1

1.	(b)	2.	(a)	3.	(b)	4.	(c)	5.	(c)	6.	(c)	7.	(d)	8.	(c)
9.	(b)	10.	(c)	11.	(b)	12.	(c)	13.	(a)	14.	(d)	15.	(a)	16.	(d)
17.	(c)	18.	(b)	19.	(b)	20.	(a)	21.	(b)	22.	(a)	23.	(b)	24.	(a)
25.	(c)	26.	(c)	27.	(c)	28.	(d)	29.	(d)	30.	(a)	31.	(d)	32.	(d)
33.	(b)	34.	(a)	35.	(d)	36.	(d)	37.	(b)	38.	(b)	39.	(c)	40.	(e)

566

ORGANIC Chemistry for IIT-JEE




1. Match the Column (I) and Column (II). (Matrix)

Column (I)		Column (II)	
	Molecule		Configuration
(a)	$ \begin{array}{c} \text{CHO} \\ \\ \text{H} - \text{C} - \text{OH} \\ \\ \text{CH}_2\text{OH} \end{array} $	(p)	R- (Rectus)
(b)	$ \begin{array}{c} \text{CHO} \\ \\ \text{HO} - \text{C} - \text{H} \\ \\ \text{CH}_2\text{OH} \end{array} $	(q)	S- (Sinister)
(c)	$ \begin{array}{c} \text{NH}_2 \\ \\ \text{H} - \text{C} - \text{CO}_2\text{H} \\ \\ \text{CH}_3 \end{array} $	(r)	D
(d)	$ \begin{array}{c} \text{NH}_2 \\ \\ \text{H} - \text{C} - \text{CH}_3 \\ \\ \text{CO}_2\text{H} \end{array} $	(s)	L

BIOMOLECULES


567

2. Comprehension




One cyclic acetal form of D-galactose is shown above.

- A.** Which atom is the anomeric carbon ?
 (a) Atom A (b) Atom B (c) Atom C (d) Atom D
 (e) Atom E (f) Atom F
- B.** Which name most completely describes this cyclic acetal form ?
 (a) α -D-Galactofuranose (b) β -D-Galactofuranose
 (c) α -D-Galactopyranose (d) β -D-Galactopyranose
- 3.** How many compound which is given below is isomer of D-Glucose ?
 D-Mannose, D-Fructose, D-Cellobiose, D-Lactose, D-Galactose, D-Ribose, D-Ribose.
- 4.** How many acidic group is present in given amino acid ?




ANSWERS — LEVEL 2

1. a - p, r; b - q, s; c - q, s; d - p, r
2. A - f; B - c
3. 5
4. 2




15 IUPAC NAME




Total number of substituent present in the above compound :

- (a) 1 (b) 2
(c) 3 (d) 4



- (a) Hex-5-en-1-yne (b) Hex-1-en-5-yne
(c) Hex-6-en-1-yne (d) Hex-1-en-6-yne




- (a) 1-Ethoxy-2, 2-dimethylcyclohexane (b) 2-ethoxy-1, 1-dimethyl cyclohexane
(c) 1, 1-Dimethyl-2-ethoxycyclohexane (d) 2-methyl-1, 1-ethoxy cyclohexane


IUPAC NAME

569


4. How many secondary carbon and hydrogen atoms are present in the molecule given below respectively ?




- (a) 2, 3 (b) 2, 2 (c) 3, 3 (d) 2, 0
5. Which IUPAC name is correct for the given compound ?




- (a) 3, 7-dimethylocta-2, 6-dienal (b) 2, 6-dimethyloct-2, 6-dienal-8
(c) 7-formyl-2, 6-dimethylhept-2, 6-diene (d) 7-aldo-2,6-dimethylhept-2, 6-diene
6. Write the IUPAC name of the following compound :




- (a) ethyl-2-(chlorocarbonyl) benzoate (b) ethyl-2-(chlorocarbonyl) hexanoate
(c) 2-(thoxycarbonyl) benzoyl chloride (d) None of these
7. The IUPAC name of the compound is :



- (a) *trans*-2-chloro-3-iodo-2-pentene (b) *cis*-2-chloro-3-iodo-2-pentene
(c) *trans*-3-iodo-4-chloro-3-pentene (d) *cis*-3-iodo-4-chloro-3-pentene
8. The IUPAC name of the compound is :



- (a) 2-methyl-6-oxohex-3-enamide (b) 6-keto-2-methyl hexanamide
(c) 2-carbamoylhexanal (d) 2-carbamoylhex-3-enal

9. The IUPAC name of  is:

- (a) 1-Bromo-2-chloro-3-fluoro-6-iodo benzene
(b) 2-Bromo-1-chloro-5-fluoro-3-iodo benzene
(c) 4-Bromo-2-chloro-5-iodo-1-fluoro benzene
(d) 2-Bromo-3-chloro-1-iodo-5-fluoro benzene

570

ORGANIC Chemistry for IIT-JEE

10. The IUPAC name of $\text{CH}_3\text{—CH}_2\text{—CH}(\text{C}_2\text{H}_5)\text{—C}(=\text{O})\text{—OCH}_3$ is:

- (a) Methyl 2-ethylbutanoate (b) 1-methoxy-2-ethylbutanone
(c) 3-Methoxycarbonylpentane (d) 1-methoxy-2-ethylbutanal

11. The IUPAC name of $\text{CH}_3\text{—CH}=\text{CH—C}(=\text{O})\text{—OH}$ is:

- (a) But-1-en-4-oic acid (b) 1-hydroxybut-2-en-1-one
(c) But-2-en-1-oic acid (d) But-2-en-4-oic acid

12. The IUPAC name of $\text{CH}_3\text{—CH}=\text{C}(\text{OCH}_3)\text{—CH}_3$ is:

- (a) 1-Methoxy-1-methylpropene (b) 2-Methoxybut-2-ene
(c) dimethylpropeneether (d) none of these

13. The IUPAC name of $\text{CH}_3\text{—C}(\text{Me})(\text{O})\text{—C}(=\text{O})\text{—OC}_2\text{H}_5$ is:

- (a) Ethyl 2-methylprop-2-enoate (b) Ethyl 2-methylprop-1-enoate
(c) 1-Ethoxy-2-methylprop-2-enone (d) 1-Ethoxy-2-methylprop-2-enal

14. The IUPAC name of $\text{CH}_3\text{—CH}(\text{CH}_3)\text{—CH}(\text{CH}_3)\text{—CH}_2\text{NH}_2$ is:

- (a) 2, 3-Dimethylbutan-4-amine (b) 2, 4-Dimethylbutan-1-amine
(c) 2,4-Dimethylbutan-4-amine (d) 2, 3-Dimethylbutan-1-amine

15. The IUPAC name of $\text{CH}_3\text{—C}(=\text{O})\text{—CH}(\text{C}_2\text{H}_5)\text{—CH}(\text{CH}_3)\text{—CH}_3$ is:

- (a) 3-(Methylethyl) pentan-2-one (b) 3-(Methylethyl)pentan-4-one
(c) 3-Ethyl-4-methylpentan-2-one (d) 3-Ethyl-2-methylpentan-4-one

16. The IUPAC name of $\text{CH}_3\text{—CH}(\text{CH}_3)\text{—CH}_2\text{—C}(=\text{O})\text{—Br}$ is:

- (a) 2-Methylbutanoyl bromide (b) 2-Methylbutan-4-oyl bromide
(c) 1-Bromo-3-Methylbutanone (d) 3-Methylbutanoyl bromide

IUPAC NAME

571

17. The IUPAC name of $\text{CH}_3 - \underset{\substack{| \\ \text{C}_6\text{H}_5}}{\text{CH}} - \text{CH}_2 - \text{OH}$ is:

- (a) 2-Phenylpropan-1-ol (b) 2-Phenylpropan-3-ol
(c) 1-(2-Hydroxy-1-methylethyl) benzene (d) 1-((Hydroxymethyl)ethyl) benzene

18. The IUPAC name of $\text{CH}_3 - \underset{\substack{| \\ \text{CH}_3}}{\overset{\substack{| \\ \text{CH}_3}}{\text{C}}} - \underset{\substack{| \\ \text{CH}_3}}{\text{CH}} - \underset{\substack{| \\ \text{I}}}{\text{CH}} - \text{CH}_2 - \text{CH}_3$ is:

- (a) 3-Iodo-4,5,5-trimethylhexane (b) 4-Iodo-1, 1, 3-trimethylhexane
(c) 4-Iodo-2, 2-dimethylheptane (d) 4-Iodo-2, 2, 3-trimethylhexane

19. The IUPAC name of $\text{CH}_3 - \text{CH}_2 - \underset{\substack{| \\ \text{Cl}}}{\text{CH}} - \underset{\substack{| \\ \text{CH}_3}}{\text{CH}} - \underset{\substack{| \\ \text{OH}}}{\overset{\substack{| \\ \text{CH}_3}}{\text{C}}} - \text{CH}_3$ is:

- (a) 4-Chloro-2, 3-dimethylhexane-2-ol (b) 4-Chloro-2-hydroxy-2, 3-dimethylhexane
(c) 4-Chloro-1, 1, 2-trimethylpentan-2-ol (d) 3-Chloro-2, 3-dimethylhexane-2-ol

20. The IUPAC name of $\text{C}_6\text{H}_5 - \underset{\substack{| \\ \text{H}}}{\text{CH}} - \text{CHO}$ is

(a) 2-Phenylpropan-3-al (b) Formylethylbenzene
(c) 2-Phenylpropanal (d) Ethylformylbenzene

21. The IUPAC name of $\text{CH}_3 - \overset{\text{O}}{\parallel}{\text{C}} - \text{CH} \begin{matrix} \nearrow \text{CH}_3 \\ \searrow \text{CH}_3 \end{matrix}$ is:

(a) 2-Methylbutan-3-one (b) 3-Methylbutan-2-al
(c) 2-Methylbutan-3-al (d) 3-Methylbutan-2-one

22. The IUPAC name of $\begin{matrix} \text{CH}_3 - \text{CO} \\ \text{CH}_3 - \text{CH}_2 - \text{CO} \end{matrix} \text{O}$ is:

(a) Ethanoic propanoic anhydride (b) Propanoic ethanoic anhydride
(c) 1-Ethanoyloxypropanone (d) 3-Ethanoyloxypropan-3-one

23. The IUPAC name of $\text{CH}_3 - \underset{\substack{| \\ \text{OH}}}{\text{CH}} - \underset{\substack{| \\ \text{C}_2\text{H}_5}}{\text{CH}} - \text{CH}_2 - \text{OH}$ is:

- (a) 3-Ethylbutane-2, 4-diol (b) 2-Ethylbutane-1, 3-diol
(c) 3-Ethylbutane-1, 3-diol (d) 2-Ethyl-1-methylpropane-1, 3-diol

572

ORGANIC Chemistry for IIT-JEE

24. The IUPAC name of $\text{CH}_3 - \underset{\text{O}}{\underset{\parallel}{\text{C}}} - \underset{\text{O}}{\underset{\parallel}{\text{C}}} - \text{CH}_3$ is:

- (a) Butane-2, 3-dial (b) Butane-1, 3-dione
(c) Butane-2, 3-dione (d) 1, 2-dimethylethanedione

25. The IUPAC name of $\text{CH}_2 = \text{CH} - \text{CH} = \text{CH}_2$ is:

- (a) Butane (b) Buta-1, 3-diene (c) Butane-1, 3-diene (d) none of these

26. The IUPAC name of $\text{CH}_2 - \text{CH}_2 - \text{CH}_2$ is:
 $\begin{array}{c} | \qquad \qquad | \\ \text{COOH} \qquad \text{COOH} \end{array}$

- (a) Pentane-1, 5-dioic acid (b) Pentane-1, 5-dicarboxylic acid
(c) Propane-1, 3-dioic acid (d) none of these

27. The IUPAC name of $\text{CH}_2 - \text{CH} = \text{CH} - \text{CHO}$ is:

- $\begin{array}{c} | \\ \text{CHO} \end{array}$
 (a) propene-1, 3-dial (b) Propene-1, 3-dicarbaldehyde
(c) Pent-3-ene-1, 5-dial (d) Pent-2-ene-1, 5-dial

28. The IUPAC name of $\text{CH}_2 - \text{CN}$ is:

- $\begin{array}{c} | \\ \text{CH}_2 - \text{CN} \end{array}$
 (a) Butane-1, 4-dicarbonitrile (b) Ethane-1, 2-dicarbonitrile
(c) Ethane-1, 2-dinitrile (d) Butane-1, 4-dinitrile

29. The IUPAC name of $\text{CH}_2 - \text{CH} - \text{CH}_2 - \text{CH}_2$ is:

- $\begin{array}{c} | \qquad | \qquad | \\ \text{NH}_2 \quad \text{CH}_3 \quad \text{NH}_2 \end{array}$
 (a) 2-Methylbutane-1, 4-diamine
(b) 3-Methylbutane-1, 4-diamine
(c) 3-(Aminomethyl)butanamine
(d) 2-(Aminomethyl)butan-4-amine

30. The IUPAC name of $\text{CH}_2 - \text{CH} - \text{CH}_2$ is:
 $\begin{array}{c} \text{CH}_2\text{Cl} \\ | \\ \text{Cl} \qquad \qquad \text{Cl} \end{array}$

- (a) Tris(chloromethyl) methane
(b) 1, 3-Dichloro-2 (chloromethyl) propane
(c) 1-Chlorobis(chloromethyl) ethane
(d) none of these

IUPAC NAME

573

31. The IUPAC name of $\text{CH}_3 - \underset{\text{OH}}{\text{CH}} - \underset{\text{CH}_3}{\text{CH}} - \underset{\text{OH}}{\text{CH}} - \overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}} - \text{CH}_3$ is:
- (a) 3, 5, 5-Trimethylhexane-2, 4-diol (b) 2, 2, 4-Trimethylhexane-3, 5-diol
(c) 1, 2, 4, 4-Tetramethylpentane-1, 3-diol (d) 2, 2, 4, 5-Tetramethylpentane-3, 5-diol
32. The IUPAC name of $\text{HOOC} - \text{CH} = \text{CH} - \text{COOH}$ is:
- (a) But-2-ene-1, 4-dicarboxylic acid (b) But-2-ene-1, 4-dioic acid
(c) Ethene dicarboxylic acid (d) Ethene dioic acid
33. The IUPAC name of $\text{CH}_3 - \overset{\text{OCH}_3}{\underset{|}{\text{CH}}} - \text{CHO}$ is:
- (a) 1-Formyl-1-methoxyethane (b) 2-Methoxypropan-3-one
(c) 2-Methoxypropanal (d) 2-Methoxypropan-3-al
34. The IUPAC name of $\text{CH}_2 = \overset{\text{CH}_3}{\underset{|}{\text{C}}} - \text{COOCH}_3$ is:
- (a) Methyl-2-methoxyprop-1-en-1-oate (b) 2-Methoxyprop-1-enylpropene
(c) 2-Methoxycarbonylprop-2-ene (d) Methyl-2-methoxyprop-2-enoate
35. The IUPAC name of $\text{CH}_3 - \text{CH} = \text{CH} - \text{COOH}$ is:
- (a) But-2-ene-1-oic acid (b) But-1-ene-1-oic acid
(c) But-2-ene-1-carboxylic acid (d) Propene-1-carboxylic acid
36. The IUPAC name of $\text{CH}_3 - \overset{\text{OH}}{\underset{|}{\text{CH}}} - \text{COOH}$ is:
- (a) 2-Hydroxypropanoic acid (b) 1-Hydroxypropanoic acid
(c) 1-Hydroxyethane carboxylic acid (d) 1-Hydroxyethanoic acid
37. The IUPAC name of $\text{HO} - \overset{\text{OH}}{\underset{|}{\text{CH}}} - \text{COOH}$ is:
- (a) 2, 3-Dihydroxybutane-1, 4-carboxylic acid
(b) 2, 3-Dihydroxybutane-1, 4-dioic acid
(c) 1, 2-Dihydroxyethane dicarboxylic acid
(d) none of these

574

ORGANIC Chemistry for IIT-JEE

38. The IUPAC name of $\text{CH}_3-\text{CH}-\overset{\text{O}}{\underset{\text{CH}_3}{\underset{\text{||}}{\text{C}}}}-\text{COOH}$ is:

- (a) 3-Methyl-2-oxobutanecarboxylic acid (b) 2-Methyl-3-oxobutan-4-oic acid
(c) 3-Methyl-2-oxobutanoic acid (d) 3-Methyl-1,2-dioxobutanoic acid

39. The IUPAC name of $\text{NC}-\text{CH}_2-\text{CH}_2-\text{COOH}$ is:

- (a) 3-Carboxy propanenitrile (b) 4-Cyanobutanoic acid
(c) 2-Cyanoethane Carboxylic acid (d) 3-Cyanopropanoic acid

40. The IUPAC name of $\begin{array}{c} \text{CH}_2\text{COOH} \\ | \\ \text{C}-\text{OH} \\ | \\ \text{COOH} \\ | \\ \text{CH}_2\text{COOH} \end{array}$ is:

- (a) 3-Carboxy-3-hydroxypentanedicarboxylic acid
(b) 2-Hydroxypropane-1, 2, 3-tricarboxylic acid
(c) 2-Hydroxypropane-1, 2, 3-trioic acid
(d) 3-Hydroxypropane-1, 2, 3-tricarboxylic acid

41. The IUPAC name of $\text{CH}_3-\overset{\text{O}}{\underset{\text{COOCCH}_3}{\underset{||}{\text{C}}}}-\text{CH}=\text{CH}_2-\overset{\text{O}}{\underset{\text{H}}{\underset{||}{\text{C}}}}$ is:

- (a) 4-ethoxycarbonylpent-3-enoic acid
(b) 4-ethanoyloxypent-3-enoic acid
(c) 3-ethoxycarbonylbut-2-enecarboxylic acid
(d) 3-ethoxycarbonylpent-3-enoic acid

42. The IUPAC name of $\text{CH}_3-\overset{\text{CH}_3}{\underset{|}{\text{CH}}}-\overset{\text{O}}{\underset{||}{\text{C}}}-\overset{\text{O}}{\underset{||}{\text{C}}}-\overset{\text{Br}}{\underset{|}{\text{NH}}}$ is:


- (a) (N-Bromo)-2-keto-3-methylbutanamide
(b) (N-Bromo)-2-keto-4-methylbutanamide
(c) (N-Bromo)-1, 2-diketo-3-methylbutanamine carboxamide
(d) (N-Bromo)-1-keto-2-methylpropane

43. The IUPAC name of $\begin{array}{c} \text{CH}_2-\text{C}=\text{CH}-\text{CH}_2\text{OH} \\ | \quad | \\ \text{Cl} \quad \text{CH}_3 \end{array}$ is:


- (a) 4-Chloro-3-methylbut-2-en-1-ol (b) 1-Chloro-2-methylbut-2-en-4-ol
(c) 4-Chloro-1-hydroxy-3-methylbut-2-ene (d) 1-Chloro-4-hydroxy-2-methylbut-2-ene

IUPAC NAME


575




- (a) 2-(Bromomethyl)-3-oxopentane carboxamide
- (b) 1-Bromo-2-carbamoylpentan-3-one
- (c) 5-Bromo-4-carbamoylpentan-3-one
- (d) 2-(Bromomethyl)-3-oxopentanamide




- (a) 2, 2-Dimethyl-4-chloro butane
- (b) 1-Chloro-3, 3-dimethylbutane
- (c) 4-Chloro-2, 2-dimethyl butane
- (d) none of these



- (a) 2, 3-Dihydroxy-4-methylpentanal
- (b) 1-oxo-2, 3-Dihydroxy-4-methylpentanal
- (c) 2,3-Dihydroxy-4-methylpentanone
- (d) 1, 2-Dihydroxy-3-methylbutanecarbaldehyde




- (a) 1-Chloro-3-methylpentan-4-one
- (b) 1-Chloro-2-(oxoethyl)butane
- (c) 5-Chloro-3-methylpentan-2-one
- (d) 3-(2-Chloroethyl)butan-2-one




- (a) 2-Hydroxypentan-4-one
- (b) 4-Hydroxypentan-2-one
- (c) 4-oxopentan-2-ol
- (d) 2-oxopentan-4-ol

576


ORGANIC Chemistry for IIT-JEE




- (a) 3-Bromo-4-chloropentan-2-ol
 (b) 3-Bromo-2-chloro-4-hydroxypentane
 (c) 3-Bromo-2-chloropentane-4-ol
 (d) none of these




- (a) 3-Bromo-4, 5-dichloropentan-3-ol
 (b) 3-Bromo-1, 2-dichloro-3-hydroxypentane
 (c) 3-Bromo-1, 2-dichloropentan-3-ol
 (d) 3-Bromo-4, 5-dichloro-3-hydroxypentane



- (a) 1-Ethoxypropan-2-ol
 (b) 3-Ethoxypropan-2-ol
 (c) 1-Ethoxy-2-hydroxypropane
 (d) none of these




- (a) 4-Bromo benzenamine
 (b) 4-Amino-1-bromobenzene
 (c) 4-Bromo benzenamide
 (d) 1-Bromo benzencarboxamide




- (a) N, N-Dimethyl aminobenzene
 (b) N, N-Dimethyl benzenamine
 (c) (a) and (b) both are correct
 (d) none of these

IUPAC NAME


577




- (a) 2, 6-Dimethylhepta-2, 5-dienoic acid
- (b) 3, 7-Dimethylhepta-2, 5-dienoic acid
- (c) 1-Hydroxy-2, 6-dimethylhepta-2, 5-dienone
- (d) none of these




- (a) 3-Methylpent-1-en-4-al
- (b) 3-Methylpent-4-enal
- (c) 3-Methylpent-4-carbaldehyde
- (d) 3-Methyl-5-oxopent-1-ene




- (a) 2-Phenyl ethanone
- (b) 1-Phenyl ethanone
- (c) 1-(Oxoethyl)benzene
- (d) 1-(Ethylaloxo)-benzene



- (a) 2-Hydroxybenzenol
- (b) 1, 2-Dihydroxybenzene
- (c) Benzene-1, 2-diol
- (d) 2-Hydroxyphenol




- (a) 2-Carboxyphenol
- (b) 2-Hydroxybenzoic acid
- (c) 1-Carboxy-2-hydroxybenzene
- (d) 2-Carboxy-1-hydroxybenzene



- (a) 1,3-Dimethyl phenol
- (b) 1-Hydroxy-2,6-dimethyl benzene
- (c) 2, 6-Dimethyl benzenol
- (d) 2-Hydroxy-1-3-dimethylbenzene


578

ORGANIC Chemistry for IIT-JEE




The IUPAC name is :

- (a) 3-phenyl prop-2-enoic acid (b) 3-phenol prop-1-enoic acid
(c) 3-carboxy-prop-1-ene benzene (d) but-2-enoic acid



- (a) Chloromethylbenzene (b) Chlorophenylmethane
(c) (a) and (b) both (d) none of these

62. The IUPAC name of the compound having the formula is :



- (a) 3, 3, 3-Trimethylprop-1-ene (b) 1, 1, 1-Trimethylprop-2-ene
(c) 3, 3-Dimethyl prop-1-ene (d) 2, 2-Dimethylbut-3-ene

63. The IUPAC name of the compound $\text{CH}_2 = \text{CH} - (\text{CH}_2\text{CH}_3)_2$ is :

- (a) 1, 1-Dimethylprop-2-ene (b) 3-Methylbut-1-ene
(c) 2-Vinyl propane (d) none of these


64. The number of sigma and pi-bonds in 1-butene 3-yne are :

- (a) 5 sigma and 5 pi (b) 7 sigma and 3 pi
(c) 8 sigma and 2 pi (d) 6 sigma and 4 pi

65. The IUPAC name of $\text{C}_6\text{H}_5\text{COCl}$ is :

- (a) Benzoyl Chloride (b) Benzene chloro ketone
(c) Benzene carbonyl chloride (d) Chloro phenyl ketone

66. The IUPAC name of the following compound is :



- (a) 4-Bromo-3-cyanophenol (b) 2-Bromo-5-hydroxybenzonitrile
(c) 2-Cyano-4-hydroxybromobenzene (d) 6-Bromo-3-hydroxybenzonitrile

IUPAC NAME

579

ANSWERS — LEVEL 1

1.	(c)	2.	(b)	3.	(b)	4.	(b)	5.	(a)	6.	(a)	7.	(a)	8.	(a)
9.	(b)	10.	(a)	11.	(c)	12.	(b)	13.	(a)	14.	(d)	15.	(c)	16.	(d)
17.	(a)	18.	(d)	19.	(a)	20.	(c)	21.	(d)	22.	(a)	23.	(b)	24.	(c)
25.	(b)	26.	(a)	27.	(d)	28.	(d)	29.	(a)	30.	(b)	31.	(a)	32.	(b)
33.	(c)	34.	(d)	35.	(a)	36.	(a)	37.	(b)	38.	(c)	39.	(d)	40.	(b)
41.	(a)	42.	(a)	43.	(a)	44.	(d)	45.	(b)	46.	(a)	47.	(c)	48.	(b)
49.	(a)	50.	(c)	51.	(a)	52.	(a)	53.	(b)	54.	(a)	55.	(b)	56.	(b)
57.	(c)	58.	(b)	59.	(c)	60.	(a)	61.	(b)	62.	(c)	63.	(b)	64.	(b)
65.	(c)	66.	(b)												

ATDB.uno

580

ORGANIC Chemistry for IIT-JEE



Give the IUPAC name of the following compounds

- 1.
- 2.
- 3.
- 4.
- 5.
- 6.
- 7.
- 8.
- 9.
- 10.
- 11.
- 12.
- 13.
- 14.
- 15.
- 16.
- 17.
- 18.

IUPAC NAME


50 51 52 53 54

581


19.



20.




21.



22.




23.




24.




25.




26.




27.




28.




29.




30.




31.




32.




33.




34.




35.




36.




37.




38.




39.




40.



41.




42.



582

ORGANIC Chemistry for IIT-JEE

43.




44.




45.




46.




47.




48.




49.




50.



51.




52.




53.




54.




55.




56.




57.




58.




59.




60.



61.



62.




IUPAC NAME


01 SEP 07 0000Z

583

63.



65.



67.




69.




71.




64.




66.




68.



70.




72.




584

ORGANIC Chemistry for IIT-JEE


73.




74.




75.




76.




77.



78.




79.




80.




81.




82.




83.




84.




85.



86.




87.




IUPAC NAME

585


88.




89.




90.




91.




92.




93.




94.




95.




96.




97.




98.




99.




100.




101.




102.



103.




104.



586

ORGANIC Chemistry for IIT-JEE


105.




106.




107.




108.




109.




110.




111.




112.




113.




114.




115.




116.




117.



118.




119.




IUPAC NAME

587


120.




121.




122.




123.




124.



125.




126.




127.



128.



129.




588



ORGANIC Chemistry for IIT-JEE

130. Give the IUPAC name of the following compounds :














ATDB.uno

IUPAC NAME

589

131. Give the IUPAC name of the following :







590

ORGANIC CHEMISTRY

ORGANIC Chemistry for IIT-JEE

132. Match the Column :

Column (I)		Column (II)	
Compound		IUPAC Name	
(a)		(p)	1, 3-dibromo-2-chlorocyclopropane
(b)		(q)	1, 2-dibromo-3-chlorocyclopropene
(c)		(r)	3-bromo-1-chlorocyclopropene
(d)		(s)	1-bromo-3-chlorocyclopropene

IUPAC NAME

591

ANSWERS – IUPAC Name

1.	pent-4-yn-1-ol	2.	pent-4-en-1-ol
3.	(5E)-2-bromo-3-chlorohept-5-en-1-ol	4.	3-bromo-2-chloro-6-ethenylcyclohexanol
5.	4-bromo-3-chloro-5-cyclopropylcyclohexanol		
6.	3-bromo-2-chloro-5-methylidenecyclohexanol		
7.	3-bromo-2-chloro-5-methylidenecyclohexane-1,4-diol		
8.	2-bromo-4-chloro-5-ethynylcyclohexanol	9.	cyclopenta-1,3-dien-1-ol
10.	2-bromo-5-chlorocyclopenta-2,4-dien-1-ol		
11.	cyclopenta-2,4-dien-1-ol	12.	dodec-10-en-4-yne-3,8-diol
13.	1-chloro-3-methylbut-3-en-2-ol	14.	4-bromo-2-chlorooct-6-ene-3,5-diol
15.	3-chloro-5-(1-methylethenyl)cyclopenta-1,3-dien-1-ol		
16.	2-bromo-5-chloro-3-methylidenecyclopentanol		
17.	cyclohexane-1,3,5-triol	18.	cyclohexane-1,3-diol
19.	cyclopenta-1,3-dien	20.	(6E)-oct-6-en-1-yne
21.	but-1-en-3-yne	22.	(3Z)-pent-3-en-1-yne
23.	buta-1,3-diyne	24.	(2E)-dec-2-en-5-yne
25.	(6Z)-dodec-6-en-2-yne	26.	pent-1-en-4-yne
27.	4-ethynyl-1-methylcyclohexene	28.	cyclooctyne
29.	(2E)-hex-2-en-4-yne	30.	5-ethynylcyclopenta-1,3-diene
31.	5-(prop-2-enyl) cyclopenta-1,3-diene		
32.	3,5-dibromo-2-chloro-4-ethenylcyclohexanol		
33.	cyclopropylmethyl cyclobutanecarboxylate		
34.	cyclobutylmethyl cyclopropanecarboxylate		
35.	ethyl ethanoate	36.	2-methylpropyl butanoate
37.	prop-2-enyl 3-ethylpent-4-enoate	38.	4-methylpentan-2-one
39.	2,5-dimethylheptan-3-one	40.	octa-1,7-dien-4-one
41.	3,4-dimethylheptane-2,6-dione	42.	9-ethyl-3-methylundecane-2,5,10-trione

592		ORGANIC Chemistry for IIT-JEE	
43.	5-methyloctanal	44.	but-3-enal
45.	3-ethylheptanal	46.	N,3-dibromobutanamide
47.	4-chloro-N-(3-chlorobutyl)butanamide		
48.	3-bromo-N-(2-bromoethyl)-4-chloropentanamide		
49.	N,3-diethyl-N-propylpentanamide		
50.	3-cyclopropyl-N-(3-cyclopropylbutyl)pentanamide		
51.	4-bromo-N-(3-bromo-2-chloropropyl)-3-chloropentanamide		
52.	3-bromo-4,5-dichloro-N-(2-chloroethyl)hexanamide		
53.	(5E)-N-(prop-2-en-yl)oct-5-enamide		
54.	3-chloro-N-(2-cyclopentylpropyl)-4-cyclopropylpentanamide		
55.	5,6-dibromo-3,4-dichloro-N-(2-chloroethyl)heptanamide		
56.	3-bromo-N-(3-bromobutyl)butanamide		
57.	3,4-dibromo-N-(3,4-dibromobutyl)pentanamide		
58.	4-bromo-N-(3-chloro-2-cyclopropyl)oct-3-(cyclopentyl)-4-yl)butanamide		
59.	2-oxopropanal	60.	4-oxopentanal
61.	3-oxobutanal	62.	3,4-dicyclopropyloct-7-yn-2-one
63.	(2E,6E)-4-bromo-5-chloro-8-cyclopropyl-10-oxodeca-2,6-dienoic acid		
64.	4,5,6,7,8,9-hexaethynyldodec-11-ynoic acid		
65.	(2E,6E)-5,9-dibromo-4-cyclopropyl-6-formylundeca-2,6-dienoic acid		
66.	3-bromo-5-chloro-6-(formylamino)hexanoic acid		
67.	ethyl 3-phenylpropanoate	68.	ethyl methyl benzene-1,3-dicarboxylate
69.	ethenylbenzene	70.	benzoyl chloride
71.	methyl 4-acetyl-2-nitrobenzoate	72.	2-formyl-4-sulphobenzoic acid
73.	1,2-dibromo-4-chlorobenzene	74.	octylbenzene
75.	2-bromo-1-chloro-4-nitrobenzene	76.	4-formyl-2-sulphobenzoic acid
77.	3-(trichloromethyl)benzoic acid	78.	4-(ethoxycarbonyl)benzoic acid

IUPAC NAME		593	
79.	4-ethenylbenzoic acid	80.	4-benzylbenzoic acid
81.	biphenyl	82.	1,2,5-tribromo-3,4,6-trichlorobenzene
83.	ethyl 4-acetylbenzoate	84.	2-bromo-6-chlorophenol
85.	2, 4, 6-tribromophenol	86.	ethyl 4-bromo-3-chlorobenzoate
87.	ethyl 3-chloro-4-mercapto/sulpho benzoate	88.	ethyl 4-(chlorocarbonyl)benzoate
89.	3-bromo-4-sulphobenzaldehyde	90.	benzene-1, 3-dicarbaldehyde
91.	4-bromo-2-chlorobenzaldehyde	92.	1-phenylpentan-1-one
93.	1-(3-bromophenyl)hexan-1-one		
94.	4-bromo-1-(3-bromo-4-chlorophenyl)pentan-1-one		
95.	1-(3-bromo-5-chlorophenyl)-2-chloroethanone		
96.	1-(3-nitrophenyl)ethanone	97.	2,4-dinitrobenzoic acid
98.	methylcyclohexane carboxylate	99.	2-methylcyclopentane carboxamide
100.	methyl-2-bromocyclohexane carboxylate	101.	2-(2-methylcyclobut-1-enyl)ethanal
102.	4-formyl-2-oxocyclohexane-1-carboxylic acid		
103.	5-cyclohexyl-6-hydroxy-5-methylhex-3-en-1-oic acid		
104.	2-oxocyclohexane-1-carboxylic acid	105.	ethyl-2-oxocyclohexane-1-carboxylate
106.	N-methylmethanamide	107.	2-bromo-4-chloro-5-ethynylcyclohexanol
108.	2-bromo-6-chloro-3-methylidene-cyclohexanol		
109.	N-phenylcyclohexane carboxamide	110.	cyclohexane carbonitrile
111.	cyclopentanecarbaldehyde	112.	cyclohexanecarboxamide
113.	2-bromo-6-methyl-cyclohexanecarbaldehyde		
114.	cyclopropane carboxylic acid		
115.	4-cyno-2-hydroxy-5-nitrocyclohexane carboxylic acid		
116.	6-chloro-5-cyno-4-nitrocyclohex-2-ene carboxylic acid		
117.	6-chloro-4-(3-oxo cyclobutyl)cyclohex-2-ene carboxylic acid		
118.	4-(2-amino-4-hydroxycyclopentyl cyclohex-2-enecarboxylic acid		

594		ORGANIC Chemistry for IIT-JEE	
119.	cyclohexylidenemethanone	120.	2-bromo-4-chloro-1-methylbenzene
121.	1,2-diethyl-3-methyl-4-propylbenzene	122.	1-ethyl-3-nitrobenzene
123.	2-methyl-1, 3, 5-trinitrobenzene	124.	1-methyl-3-nitrobenzene
125.	Diphenylmethane	126.	4-Methyl-3-nitrobenzene sulphonic acid
127.	1-ethenyl-4-methylbenzene	128.	heptan-3-amine
129.	2-methylpentan-3-amine		
130.	(a) 2,11-dimethyltridecane		
	(b) 12-methyl-tridecan-3-ol		
	(c) 4-ethyl-3-methyloctane		
	(d) 1,2,7-trimethylcyclopentadecane		
	(e) 1,1,2,5-tetramethylcyclopentane		
131.	(a) 5-isobutyl-2-methylnonane, 5-2-methylnonane (2-methylpropyl)		
	(b) 2,7-dimethyl-4-propyloctane		
	(c) 6-(2,3-dimethylbutyl)-2-methylnonane		
	(d) 1,1,2,6-tetramethylcyclohexane		
	(e) 2,2,3-trimethylcyclohexanol		
	(f) 1,2,4-trimethylcyclopentane		
	(g) 2-methylbicyclo[3.1.1] heptane		
	(h) 2-methylbicyclo[3.1.1] heptane		
	(i) 2,2,6,6,7-pentamethyloctane		
	(j) 4-(1methylpropyl)-2,3,5-trimethyl nonane Not 2-butyl		
	(k) 5-(1-ethyl-2-methyl-1-(1-methylethyl)propyl) nonane		
132.	a - r; b - s; c - q; d - p		

ATDB.uno

ATDB.uno

ATDB.uno

ATDB.uno

ATDB.uno

About the Author



M.S. Chouhan, the author, is B.Tech in Chemical Engineering from Mumbai University. Due to his keen interest in teaching, he opted his career in guiding the JEE aspirants. He is highly dedicated to Organic Chemistry and successfully shaping the dreams of IITians, who are the most honoured technocrats and earning the name for themselves and the country.



Shri Balaji Publications
Muzaffarnagar (U.P.)