

# PRAAYAS

## JEE 2026

ATDB.uno

Mathematics

# Basic Maths

Lecture - 13

By - Ashish Agarwal Sir  
(IIT Kanpur)




# Topics *To be covered*



- A** Introduction to Modulus
- B** Problem Practice

ATDB.uno





# Homework Discussion

ATDB.uno

# QUESTION [JEE Mains 2023]



The number of integral solutions x of  $\log_{\left(x+\frac{7}{2}\right)} \left(\frac{x-7}{2x-3}\right)^2 \geq 0$  is:

$\left(\frac{x-7}{2x-3}\right)^2 > 0$   
 $x \neq 7$  — (A)  
 $x \neq \frac{3}{2}$

- A** 8
- B** 7
- C** 5
- ~~**D** 6~~

Case (i) if  $x + \frac{7}{2} > 1 \Rightarrow x > -5/2$

$\left(\frac{x-7}{2x-3}\right)^2 \geq 1$

$\frac{(x-7)^2}{(2x-3)^2} - 1 \geq 0$

$\frac{(x-7)^2 - (2x-3)^2}{(2x-3)^2} \geq 0$

$\frac{(3x-10)(-x-4)}{(2x-3)^2} \geq 0$

Case (ii) if  $0 < x + \frac{7}{2} < 1$   
 $-\frac{7}{2} < x < -5/2$

$\frac{(x-7)^2}{(2x-3)^2} \leq 1$

$\frac{(3x-10)(-x-4)}{(2x-3)^2} \leq 0$

$(3x-10)(x+4) \geq 0 \quad x \neq 3/2$


$x \in (-\infty, -4] \cup [10/3, \infty)$



ATDB.uno



$$\frac{(3x-10)(x+4)}{(2x-3)^2} \leq 0, x \neq \frac{3}{2}$$



$$x \in [-4, \frac{10}{3}] - \{\frac{3}{2}\}$$

⇓

$$x \in (-\frac{5}{2}, \frac{10}{3}] - \{\frac{3}{2}\}$$

ATDB.uno

$x \in \phi$

$$(-\frac{5}{2}, \frac{10}{3}] - \{\frac{3}{2}\} \text{ --- (B)}$$

FINAL ANS :  $A \cap B = (-\frac{5}{2}, \frac{10}{3}] - \{\frac{3}{2}\}$

Integral values of  $x = -2, -1, 0, 1, 2, 3 \Rightarrow 6$  values.

## QUESTION



Solve the following inequalities:

$$(a) \log(x^2 - 2x - 2) \leq 0$$

$$(b) \log_5(x^2 - 11x + 43) < 2$$

$$(c) 2 - \log_2(x^2 + 3x) \geq 0$$

$$(d) \log_{1.5} \frac{2x-8}{x-2} < 0$$

$$(e) \log_3 \frac{1+2x}{1+x} < 1$$

$$(f) \log_4 \frac{3x+2}{x} \leq 0.5$$

$$(g) \log_2 \frac{x^2-4x+2}{x+1} \leq 1$$

ATDB.uno

$$(h) \log_2 \left( \frac{4x-3}{4-3x} \right) > -\frac{1}{2}$$

Answers:

$$(a) [-1, 1 - \sqrt{3}) \cup (1 + \sqrt{3}, 3];$$

$$(b) (2, 9);$$

$$(c) [-4, -3) \cup (0, 1];$$

$$(d) (4, 6);$$

$$(e) (-\infty, -2) \cup (-1/2, \infty);$$

$$(f) [-2, -2/3);$$

$$(g) [0, 2 - \sqrt{2}) \cup (2 + \sqrt{2}, 6];$$

$$(h) \left( \frac{3}{4}, \frac{4}{3} \right)$$



$$\textcircled{H} \left( \log_2 \left( \frac{4x-3}{4-3x} \right) \right)^2 \geq -\frac{1}{2}$$

∴  $\log_2 \left( \frac{4x-3}{4-3x} \right)$  should be defined

$$\frac{4x-3}{4-3x} > 0$$

ATDB.uno

$$\frac{4x-3}{3x-4} < 0$$

$$\begin{array}{c} + \quad - \quad + \\ | \quad | \quad | \\ 3/4 \quad 4/3 \end{array}$$

$$x \in (3/4, 4/3) \text{ Ans}$$



## Home Challenge-04

JEE MAINS 10 Apr Morning SHIFT

2023



Let  $a, b, c$  be three distinct positive real numbers such that

$(2a)^{\log_e a} = (bc)^{\log_e b}$  and  $b^{\log_e 2} = a^{\log_e c}$ . Then,  $6a + 5bc$  is equal to

$$\ln(2a)^{\ln a} = \ln(bc)^{\ln b}$$

$$\ln a \cdot (\ln 2 + \ln a) = \ln b \cdot (\ln b + \ln c)$$

$$\ln a (\ln 2 + \ln a) = \ln b (\ln b + \ln c)$$

$$\ln 2 \cdot \ln a + (\ln a)^2 = (\ln b)^2 + \frac{\ln b \cdot \ln 2 \cdot \ln b}{\ln a}$$

$$(\ln a)^2 - (\ln b)^2 = \frac{(\ln b)^2 \cdot \ln 2}{\ln a} - \ln 2 \cdot \ln a = \ln 2 \left( \frac{(\ln b)^2}{\ln a} - \ln a \right) = \frac{\ln 2 ((\ln b)^2 - (\ln a)^2)}{\ln a}$$

$$\ln b^{\ln 2} = \ln a^{\ln c}$$

$$\ln 2 \cdot \ln b = \ln c \cdot \ln a$$

$$\ln c = \frac{\ln 2 \cdot \ln b}{\ln a}$$

ATDB.uno



$$(\ln a)^2 - (\ln b)^2 = \ln 2 \left( \frac{(\ln b)^2 - (\ln a)^2}{\ln a} \right)$$

$$(\ln a)^2 - (\ln b)^2 - \ln 2 \cdot \frac{(\ln b)^2 - (\ln a)^2}{\ln a} = 0$$

$$[(\ln a)^2 - (\ln b)^2] \left( 1 + \frac{\ln 2}{\ln a} \right) = 0$$

$$(\ln a)^2 - (\ln b)^2 = 0 \quad \text{or} \quad \frac{\ln 2}{\ln a} + 1 = 0$$

$$(\ln a - \ln b)(\ln a + \ln b) = 0$$

$$\ln a = \ln b \quad \text{or} \quad \ln a + \ln b = 0$$

$$a = b \quad \text{or} \quad \ln a = -\ln b = \ln \frac{1}{b}$$

(N.P)  $a = \frac{1}{b}$

$$\ln 2 = -\ln a$$

$$\ln a = \ln \frac{1}{2}$$

$$a = \frac{1}{2} \Rightarrow \ln c = \frac{\ln 2 \cdot \ln b}{\ln a} = \frac{\ln 2 \cdot \ln b}{-\ln 2} = -\ln b \Rightarrow \ln b + \ln c = 0$$

$$\ln bc = 0 \Rightarrow bc = 1$$

~~$$x^2 = x$$~~

$$x = 1$$

Gadho/Gadhiyoo a isaa  
naa kano!!

Sahi

$$x^2 = x$$

$$x^2 - x = 0 \Rightarrow x(x-1) = 0$$

$$x = 0, 1$$



$$\text{If } a = \frac{1}{b}$$

$$\ln c = \frac{\ln 2 \cdot \ln b}{\ln a}$$

$$\ln c = \frac{\ln 2 \cdot \ln b}{-\ln b} = -\ln 2$$

$$c = \frac{1}{2}$$

Hence

$$6a + 5bc = 6 \cdot \frac{1}{2} + 5 = \underline{8 \text{ Ans}}$$

ATDB.uno

$6a + 5bc$  has  $\infty$  many possible values.

$$(a^2)^{\ln a} = (bc)^{\ln b}, \ln 2 = a^{\ln c} \rightarrow b^{\ln 2} = \left(\frac{1}{b}\right)^{\ln 1/2} = (b^{-1})^{-\ln 2} = b^{\ln 2}$$

$$\left(\frac{2}{b}\right)^{\ln \frac{1}{b}} = \left(\frac{b}{2}\right)^{\ln b}$$

$$\left(\frac{2}{b}\right)^{-\ln b} = (b/2)^{\ln b} \Rightarrow (b/2)^{\ln b} = (b/2)^{\ln b}$$



# Aao Machaay Dhamaal Deh Swaal pe Deh Swaal

ATDB.uno

**QUESTION [JEE Mains 2021]**

The number of solutions of the equation  $\log_4(x - 1) = \log_2(x - 3)$  is

Tahoi

ATDB.uno

## QUESTION



Find  $x$  for  $\frac{(\ln x)^2 - 3 \ln x + 3}{\ln x - 1} < 1.$

$$\ln x = t$$

$$\frac{t^2 - 3t + 3}{t - 1} - 1 < 0$$

$$\frac{t^2 - 4t + 4}{t - 1} < 0$$

$$\frac{(t - 2)^2}{(t - 1)} < 0$$

$$\frac{1}{t - 1} < 0 \quad t \neq 2$$

$$t - 1 < 0$$

$$t < 1$$

ATDB.uno

$$\ln x < 1 \text{ \& } x > 0, \ln x \neq 2$$

$$\log_e x < 1$$

$$x < e$$

$$x \neq e^2$$

$$x \in (0, e)$$

## QUESTION



$$\text{Solve : } \log_{\frac{1}{x}} \left( \frac{2(x-2)}{(x+1)(x-5)} \right) \geq 1$$

Tah02

ATDB.uno




# Modulus/Absolute value Function



$$|x| = \begin{cases} x & x \geq 0 \\ -x & x < 0 \end{cases}$$

★  $|x| \geq 0$

★  $|x| = 0 \iff x = 0$




ATDB.uno



# Geometrical Meaning of Modulus



$|x|$  = distance of  $x$  on number line from 0



Ex:  $|x| = 2$

$x = -2, 2$

Ex:  $|2x - 1| = 5$

$2x - 1 = -5, 5$

$2x = -4, 6$

$x = -2, 3$

Ex:  $|2x - 6| = -10$

→ (No soln)

$|x| = a, a > 0$   
 $\Downarrow$   
 $x = \pm a$



## Important Properties



$$P_1: |-x| = |x| \quad \star\star\star\star$$

$$P_2: \left| \frac{x}{y} \right| = \frac{|x|}{|y|}$$

Module distributes  
over multiplication  
& division

ATDB.uno

$$|x_1 \cdot x_2 \cdots x_n| = |x_1| \cdot |x_2| \cdots |x_n|$$

$$P_3: |xy| = |x||y|$$



## Important Properties



**P<sub>4</sub>**:  $\sqrt{x^2} = |x|, x \in \mathbb{R}$ , Also  $x^2 = |x|^2$

Yaad Rakhe (variable/Expression)<sup>2</sup>  
 ko  $|variable/Expression|^2$   
 likh sakte hai

**P<sub>5</sub>**:  $|x| \geq 0$  also  $|x| \geq x$

✧  $|x| > x \rightarrow x \in \mathbb{R}^-$

✧  $|x| = x \rightarrow x \in [0, \infty)$

ATDB.uno



$$(i) |f(x)| = a, a \geq 0$$

$$f(x) = \pm a$$

$$(ii) |f(x)| = g(x)$$

$$\text{Case ① } f(x) \geq 0$$

$$\downarrow \text{ solve } f(x) = g(x)$$

$$\text{Case ② } f(x) < 0$$

$$-f(x) = g(x)$$

$$|f(x)| = |g(x)|$$

$$\downarrow$$

$$(f(x))^2 = (g(x))^2$$

$$\text{Apply } a^2 - b^2$$

$$\text{||}$$

$$(a-b)(a+b)$$

UNION

(FINAL Ans)

## QUESTION

$$|x^2 + x - 20| = -(x^2 + x - 20)$$


$$\Downarrow$$

$$x^2 + x - 20 \leq 0$$

$$\Downarrow$$

$$(x+5)(x-4) \leq 0$$

$$x \in [-5, 4]$$



ATDB.uno



## QUESTION



$$\left| \left( \frac{x^2 - 6x + 8}{x^2 - 4x + 3} \right) \right| = - \left( \frac{x^2 - 6x + 8}{x^2 - 4x + 3} \right)$$

Tah03

ATDB.uno

## QUESTION



$$\text{Solve: } |2x - 1| = 7$$

$$2x - 1 = -7,7$$

$$2x = -6,8$$

$$x = -3,4$$

# ATDB.uno

## QUESTION



Solve:  $|3 - x| = 2$

$3 - x = -2, 2$

$-x = -5, -1$

$x = 5, 1$

ATDB.uno

## QUESTION



$|x + 2| = 2(3 - x)$  then  $x$  is equal to

~~A~~  $4/3$

B  $-8$

C  $8$

D  $-4/3$

Case ① if  $x + 2 > 0 \Rightarrow x > -2$

$$x + 2 = 6 - 2x$$

$$3x = 4$$

$$x = \frac{4}{3}$$

Case ② if  $x + 2 < 0 \Rightarrow x < -2$

$$-x - 2 = 6 - 2x$$

$$x = 8$$

$x \in \phi$

ATDB.uno

$$x = \frac{4}{3}$$

## QUESTION



If  $|3x - 2| + x = 11$  then  $x$  is

Tah 04

- A**  $13/4$
- B**  $9/2$
- C**  $-9/2$
- D**  $-13/4$

ATDB.uno

## QUESTION



$||x - 2| - 2| - 2| = 2$  then sum of all values of  $x$  satisfying the equation is

~~A~~ 8

B 12

C 0

D 4

$$||x - 2| - 2| - 2| = 2$$

$$|m - 2| = 2$$

$$m - 2 = -2, 2$$

$$m = 0, 4$$

ATDB.uno

$$m = 0$$

$$|x - 2| - 2 = 0$$

$$|x - 2| - 2 = 0$$

$$|x - 2| = 2$$

$$x - 2 = -2, 2$$

$$x = 0, 4$$

$$m = 4$$

$$|x - 2| - 2 = 4$$

$$|x - 2| - 2 = -4, 4$$

$$|x - 2| = -2, 6$$

$$|x - 2| = -2 \text{ or } |x - 2| = 6$$

Not possible

$$x - 2 = 6, -6$$

$$x = 8, -4$$



$$x = 0, 4, 8, -4$$

$$\text{Sum} = 0 + 4 + 8 - 4 = 8$$


# ATDB.uno

**QUESTION**



$F_1$        $F_2$

Solve:  $|x + 1| + |x - 2| = 5$



Case I if  $x > 2$

$$x + 1 + x - 2 = 5$$

$$2x = 6$$

$$x = 3$$

$x = 3$

Case II  $-1 < x < 2$

$$x + 1 - (x - 2) = 5$$

$$3 = 5$$

↓  
Not possible

Case III if  $x \leq -1$

$$-(x + 1) - (x - 2) = 5$$

$$-2x + 1 = 5$$

$$x = -2$$

$x = -2$

UNION

Ans  $x = -2, 3$

ATDB.uno



# QUESTION

$|x - 1| + |x - 2| + |x - 3| = 9$  then x can be

Tah 05

- A** -5
- B** 9
- C** -1
- D** 5

$T_1$	-ve	+	+	+
$T_2$	-ve	-ve	+	+
$T_3$	-ve	-ve	-ve	+

Case ① if  $x \leq 1$

$$-(x-1) - (x-2) - (x-3) = 9$$

$$-3x + 6 = 9$$

$$x = -1$$

x = -1

Case ② if  $1 < x \leq 2$

Case ③

ATDB.uno

## QUESTION [JEE Mains 2021]



The number of real solution of the equation,  $x^2 - |x| - 12 = 0$  is:

- A** 2  
**B** 3  
**C** 1  
**D** 4

M(1)

case (i) if  $x \geq 0$

$$x^2 - x - 12 = 0$$

$$(x-4)(x+3) = 0$$

$$x = -3, 4$$

case (ii) if  $x < 0$

$$x^2 + x - 12 = 0$$

$$(x+4)(x-3) = 0$$

$$x = -4, 3$$

ATDB.uno

$$x = -4, 4$$

M(2)

$$x^2 - |x| - 12 = 0$$

$$|x|^2 - |x| - 12 = 0$$

$$|x| = t \quad t^2 - t - 12 = 0$$

$$t = 4, -3$$

~~$$|x| = 4, -3$$~~

$$x = \pm 4$$

## QUESTION [JEE Mains 2025 (8 April)]



Tah 06

The sum of the squares of the roots of  $|x - 2|^2 + |x - 2| - 2 = 0$  and the squares of the roots of  $x^2 - 2|x - 3| - 5 = 0$ , is

- A** 24
- B** 26
- C** 36
- D** 30

$$|x - 2| = t$$

$$t^2 + t - 2 = 0$$

$$(t + 2)(t - 1) = 0$$

$$t = -2, 1$$

ATDB.uno

~~$$|x - 2| = -2, 1$$~~

Ans. C

## QUESTION [JEE Mains 2025 (7 April)]



The number of real roots of the equation  $x|x - 2| + 3|x - 3| + 1 = 0$  is:

Tah 07

**A** 4

**B** 3

**C** 2

**D** 1

$$\begin{array}{cccc}
 T_1 & -ve & | & +ve & | & +ve \\
 T_2 & -ve & 2 & -ve & 3 & +ve
 \end{array}$$

case 1

ATDB.uno

Ans. D

## QUESTION [JEE Mains 2024 (5 April)]



The number real solutions of the equations  $x|x + 5| + 2|x + 7| - 2 = 0$  is

Tahos

# ATDB.uno

Ans. 3

## QUESTION [JEE Mains 2024 (5 April)]



The number of distinct real roots of the equation  $|x| |x + 2| - 5|x + 1| - 1 = 0$  is

Tah 09

$T_1$	-ve	-ve	-ve	+
$T_2$	-ve - 2	+ve - 1	+ve 0	+
$T_3$	-ve	-ve	+ve	

Case ① if  $x \leq -2$

$$-x \cdot -(x+2) + 5(x+1) - 1 = 0$$

$$x^2 + 2x + 5x + 5 - 1 = 0$$

$$x^2 + 7x + 4 = 0$$

$$x = \frac{-7 \pm \sqrt{49 - 16}}{2} = \frac{-7 \pm \sqrt{33}}{2}$$

$$\frac{-7 + \sqrt{33}}{2} > -2$$

$$\frac{-7 + \sqrt{33}}{2} > -4$$

$$\sqrt{33} > 3$$


$$x = \frac{-7 - \sqrt{33}}{2}$$

Ans. 3

**QUESTION [JEE Mains 2024 (8 April)]**



The number of distinct real roots of the equation  $|x + 1| |x + 3| - 4|x + 2| + 5 = 0$  is



Case (i) if  $x \leq -3$

Case (ii) if  $-3 < x \leq -2$

Case (iii) if  $-2 < x < -1$


Case (iv) if  $x > -1$

ATDB.uno

Thumbs up  
Thumbs down

Yes understood — 96%

Abhi bhi nahi aaya — 4%



No: of Number  
 ke Tukde  
 ||  
 No: of cases

Ans. 2

## QUESTION [JEE Mains 2023]



The sum of all the roots of the equation  $|x^2 - 8x + 15| - 2x + 7 = 0$  is:

**A**  $11 + \sqrt{3}$

~~**B**  $9 + \sqrt{3}$~~

**C**  $9 - \sqrt{3}$

**D**  $11 - \sqrt{3}$

case ① If

$$x^2 - 8x + 15 > 0$$

$$(x-3)(x-5) > 0$$

$$\begin{array}{c} + \quad - \quad + \\ | \quad | \quad | \\ 3 \quad 5 \end{array}$$

$$x \in (-\infty, 3) \cup (5, \infty)$$

$$x^2 - 8x + 15 - 2x + 7 = 0$$

$$x^2 - 10x + 22 = 0$$

$$x = \frac{10 \pm \sqrt{12}}{2}$$

$$x = 5 \pm \sqrt{3}$$

case ②

If  $x^2 - 8x + 15 < 0$

$$(x-5)(x-3) \leq 0$$

$$x \in [3, 5]$$

$$-(x^2 - 8x + 15) - 2x + 7 = 0$$

$$-x^2 + 6x - 8 = 0$$

$$x^2 - 6x + 8 = 0$$

$$x = 2, 4$$

$$x = 5 + \sqrt{3}$$

UNION

$$x = 4, 5 + \sqrt{3}$$

$$\text{Sum} = 9 + \sqrt{3}$$

## QUESTION [JEE Mains 2024 (30 Jan)]



The number of real solutions of the equation  $x(x^2 + 3|x| + 5|x - 1| + 6|x - 2|) = 0$  is \_\_\_\_\_

$$x = 0 \quad \text{or} \quad \underbrace{x^2}_{\geq 0} + \underbrace{3|x|}_{\geq 0} + \underbrace{5|x-1|}_{\geq 0} + \underbrace{6|x-2|}_{\geq 0} = 0$$

$x = 0$  (only one soln)  
 $x^2 = 0 \Rightarrow x = 0$   
 $x - 1 = 0 \Rightarrow x = 1$   
 $x - 2 = 0 \Rightarrow x = 2$

$x = 0$   
 $x = 0$   
 $x = 1$   
 $x = 2$

$x \in \phi$   
 NO soln

Ans. 1

# QUESTION [JEE Mains 2018]



Let  $S = \{x \in \mathbb{R} : x \geq 0 \text{ and } 2|\sqrt{x} - 3| + \sqrt{x}(\sqrt{x} - 6) + 6 = 0\}$ . Then  $S$  :

- ~~A~~ contains exactly two elements
- B contains exactly four elements
- C is an empty set
- D contains exactly one element

$$\sqrt{x} = t$$

$$2|t-3| + t(t-6) + 6 = 0$$

Case ①  $t \geq 3$

$$2(t-6) + t^2 - 6t + 6 = 0$$

$$t^2 - 4t = 0$$

$$t = 0, 4$$

$$t = 4$$

Case ② if  $t < 3$

$$-2t + 6 + t^2 - 6t + 6 = 0$$

$$t^2 - 8t + 12 = 0$$

$$t = 2, 6$$

$$t = 2$$

ATDB.uno

$$t = 4, 2$$

$$\sqrt{x} = 4, 2$$

$$S = \{16, 4\}$$

$$x = 16, 4$$

**QUESTION [JEE Mains 2021]**

Tah09

The number of the real roots of the equation  $(x + 1)^2 + |x - 5| = \frac{27}{4}$  is

**ATDB.uno**



**Sabse Important Baat**



**Sabhi Class Illustrations Retry Karnay hai...**

ATDB.uno



# Today's BPP

## ATDB.uno



## Lo Karo Duvaadaar Practice!!



1.  $\log_5(x^2 - 3x + 3) > 0$

2.  $\log_7[\log_5(x^2 - 7x + 15)] > 0$

3.  $\log_{\left(\frac{1}{2}\right)}[\log_5(x^2 - 7x + 17)] > 0$

4.  $\log_{\left(\frac{1}{2}\right)}(\log_5(\log_2(x^2 - 6x + 40))) > 0$

5.  $\log_3[\log_5 \log_2(x^2 - 9x + 50)] > 0$

6.  $\log_6\left(\frac{x-2}{6-x}\right) > 0$

7.  $\log_{0.5}(x^2 - 5x + 6) > -1$

8.  $\log_8(x^2 - 4x + 3) < 1$

9.  $\log_{\left(\frac{1}{4}\right)}\left(\frac{35-x^2}{x}\right) \geq -\frac{1}{2}$



## Answers



1.  $(-\infty, 1) \cup (2, \infty)$

2.  $(-\infty, 2) \cup (5, \infty)$

3.  $(3, 4)$

4.  $(2, 4)$

5.  $(-\infty, 3) \cup (6, \infty)$

6.  $(4, 6)$

7.  $(1, 4)$

8.  $(-1, 5)$

9.  $(-1, 0) \cup (5, \infty)$

ATDB.uno



## Home Challenge-05



If the value of  $x$  which satisfies the equation  $2 \log_3 \sqrt{3^{1-x} + 2} = 1 + \log_3(4 \cdot 3^x - 1)$  is given by,  $1 - \log_3 k$ , then find the value of  $k$ . [Ans. 4]

ATDB.uno



## Homework From Module



**Prarambh (Topicwise) : Q1 to Q17**

**Prabal (JEE Main Level) : Q1 to Q7**

# ATDB.uno



# Solution to Previous TAH

## ATDB.uno

**QUESTION [JEE Mains 2023]**

The number of integral solutions  $x$  of  $\log_{\left(x+\frac{7}{2}\right)} \left(\frac{x-7}{2x-3}\right)^2 \geq 0$  is :

**A** 8

**B** 7

**C** 5

**D** 6

**ATDB.uno**



**Q-3 (TAH-1):** The number of integral solutions  $x$  of  $\log_{x+\frac{7}{2}} \left(\frac{x-7}{2x-3}\right)^2 \geq 0$  is:  
 (A) 8 (B) 7 (C) 5 (D) 6.

**TAH-1, By Reed from WB**

**Soln**

$\left(\frac{x-7}{2x-3}\right)^2 > 0$  for log to be defined

$\Rightarrow \frac{x-7}{2x-3} \neq 0 \Rightarrow x \neq 7, x \neq \frac{3}{2}$  — (A)

**Case-1:** If  $x + \frac{7}{2} > 1$   
 $\Rightarrow x > 1 - \frac{7}{2} \Rightarrow x > -\frac{5}{2}$  — (P.C.)

$\therefore \log_{x+\frac{7}{2}} \left(\frac{x-7}{2x-3}\right)^2 \geq 0$   
 $\Rightarrow \left(\frac{x-7}{2x-3}\right)^2 \geq 1$   
 $\Rightarrow \frac{(x-7)^2}{(2x-3)^2} - 1 \geq 0$   
 $\Rightarrow \frac{(x-7)^2 - (2x-3)^2}{(2x-3)^2} \geq 0$   
 $\Rightarrow \frac{(x-7+2x-3)(x-7-2x+3)}{(2x-3)^2} \geq 0$   
 $\Rightarrow \frac{(3x-10)(-x-4)}{(2x-3)^2} \geq 0$   
 $\Rightarrow (3x-10)(x+4) \leq 0, x \neq \frac{3}{2}$



$\therefore x \in [-4, \frac{10}{3}] - \{\frac{3}{2}\}$

**TAH 1(part-1) by Reed from WB**


Taking 'n' with P.C.  
 $x > -\frac{5}{2} \cap x \in [-4, \frac{10}{3}] - \{\frac{3}{2}\}$   
  
 $\therefore x \in (-\frac{5}{2}, \frac{3}{2}) \cup (\frac{3}{2}, \frac{10}{3}]$  — (B)

**Case-2:** If  $0 < x + \frac{7}{2} < 1$ .  
 or,  $-\frac{7}{2} < x < -\frac{5}{2}$  — (P.C.)

**TAH 1(part 2) by Reed from WB**

now,  $\log_{x+\frac{7}{2}} \left(\frac{x-7}{2x-3}\right)^2 \geq 0$   
 $\Rightarrow \left(\frac{x-7}{2x-3}\right)^2 \leq 1$   
 $\Rightarrow \frac{(3x-10)(x+4)}{(2x-3)^2} \geq 0$   
 $\Rightarrow (3x-10)(x+4) \geq 0, x \neq \frac{3}{2}$


$\therefore x \in (-\infty, -4] \cup [\frac{10}{3}, \infty)$



$x \in \phi$  — (C)

$(\text{Case-1}) \cup (\text{Case-2}) \Rightarrow \text{Case-0}$   
 $x \in (-\frac{5}{2}, \frac{3}{2}) \cup (\frac{3}{2}, \frac{10}{3}]$  — (B)

now Taking intersection of (A) & (B) (i.e.  $A \cap B$ )



$\therefore x \in (-\frac{5}{2}, \frac{10}{3}] - \{\frac{3}{2}\} \rightarrow$  final condition of 'x'

$\therefore$  Integral values in this region

$= -2, -1, 0, 1, 2, 3.$

6 integral values.  $\therefore$  Ans  $\Rightarrow$  (D) 6.

**QUESTION**

$$\log_{x+3}(x^2 - x) < 1$$

# ATDB.uno



Tah 2

$\log_{x+3} (x^2-x) < 1$

Also  $x+3 > 0$   
 $x > -3$   
No Need

Also  $x-x > 0$   
 $x(x-1) > 0$

$x \in (-\infty, 0) \cup (1, \infty)$   
 (1)

Case I: if  $x+3 > 1 \Rightarrow x > -2$

Case II:  $0 < x+3 < 1$

$x^2-x < x+3$

$x^2-2x-3 < 0$

$(x-3)(x+1) < 0$

$x \in (-1, 3)$

$x \in (-1, 3)$  - (2)

$x^2-x > x+3$

$x^2-2x-3 > 0$

$(x-3)(x+1) > 0$

$x \in (-\infty, -1) \cup (3, \infty)$


$-3 < x < -2$

$x \in (-3, -2)$  - (3)

**Sakshi sahu**

(2)  $\cup$  (3)

$x \in (-3, -2) \cup (-1, 3) \cap (1)$



$x \in (-3, -2) \cup (-1, 0) \cup (1, 3)$

# Kriti Mathur Raj.



TAH20

$$\log_{x+3} (x^2 - x) < 1$$

Case 1: If  $x+3 > 1$   
 $x > -2$   
 $x \in (-2, \infty)$  - (A)

$$x^2 - x < x + 3$$


$$x^2 - x - x - 3 < 0$$

$$x^2 - 2x - 3 < 0$$


$$x^2 - 3x + x - 3 < 0$$

$$x(x-3) + 1(x-3) < 0$$

$$(x+1)(x-3) < 0$$



(A)  $\cap$  (B)



Case 2: If  $0 < x+3 < 1$   
 $-3 < x < -2$

$x \in (-3, -2)$  - (D)


$$x^2 - x > x + 3$$

$$x^2 - x - x - 3 > 0$$

$$x^2 - 2x - 3 > 0$$

$$(x+1)(x-3) > 0$$

$x \in (-\infty, -1) \cup (3, \infty)$  - (E)



(D)  $\cap$  (E)

(E)  $\cup$  (F)

$x \in (-3, -2) \cup (-1, 3)$  - (G)


Now,

$$x^2 - x > 0$$

$$x(x-1) > 0$$

Number line with points 0 and 1. The intervals  $(-\infty, 0)$  and  $(1, \infty)$  are shaded, with '+' signs above and '-' signs below. This is labeled (H).

(G)  $\cap$  (H)



$x \in (-3, -2) \cup (-1, 0) \cup (1, 3)$  Ans.

ATDB.uno




**TAH-2:**  $\log_{x+3}(x^2-x) < 1$ .

Solve!

$$\log_{x+3}(x^2-x) < 1$$

for log to be defined,

$$\begin{aligned} x+3 > 0 & \quad | \quad x+3 \neq 1 & \quad | \quad x^2-x > 0 \\ \Rightarrow x > -3 & \quad | \quad \Rightarrow x \neq -2 & \quad | \quad \Rightarrow x(x-1) > 0 \\ & & & \quad \Rightarrow x \in (-\infty, 0) \cup (1, \infty) \end{aligned}$$



$\therefore x \in (-3, 0) \cup (1, \infty) - \{-2\}$

**TAH 2  
by Reed  
From WB**

**Case-1:** If  $x+3 > 1 \Rightarrow x > -2$  — (P.C)

So,  $\log_{x+3}(x^2-x) < 1$

$$\begin{aligned} \Rightarrow (x^2-x) &< x+3 \\ \Rightarrow x^2-2x-3 &< 0 \\ \Rightarrow (x-3)(x+1) &< 0 \end{aligned}$$



$\therefore x \in (-1, 3)$  — (P.C)


**Case-2:** If  $x+3 \in (0, 1)$ .

$$\begin{aligned} \Rightarrow 0 < x+3 < 1 \\ \Rightarrow -3 < x < -2 \end{aligned} \quad \therefore x \in (-3, -2) \text{ — (P.C)}$$

So,  $\log_{x+3}(x^2-x) < 1$

$$\begin{aligned} \Rightarrow (x^2-x) &> (x+3) \\ \Rightarrow x^2-2x-3 &> 0 \\ \Rightarrow (x-3)(x+1) &> 0 \end{aligned}$$

$\therefore x \in (-\infty, -1) \cup (3, \infty)$




$\therefore x \in (-3, -2)$  — (P.C)

Taking (1)  $\cup$  (2)  $\Rightarrow x \in (-3, -2) \cup (-1, 3)$  — (P.C)

Taking intersection of (1) & (2);

$A \cap B \Rightarrow$



$\therefore x \in (-3, -2) \cup (-1, 3)$



# Solution to Previous BPPs

## ATDB.uno

## QUESTION



Solve the following inequalities:

(a)  $\log(x^2 - 2x - 2) \leq 0$

(b)  $\log_5(x^2 - 11x + 43) < 2$

(c)  $2 - \log_2(x^2 + 3x) \geq 0$

(d)  $\log_{1.5} \frac{2x-8}{x-2} < 0$

(e)  $\log_3 \frac{1+2x}{1+x} < 1$

(f)  $\log_4 \frac{3x+2}{x} \leq 0.5$

(g)  $\log_2 \frac{x^2-4x+2}{x+1} \leq 1$

(h)  $\log_2 \left( \frac{4x-3}{4-3x} \right) > -\frac{1}{2}$

Answers:

(a)  $[-1, 1 - \sqrt{3}) \cup (1 + \sqrt{3}, 3]$

(b)  $(2, 9)$

(c)  $[-4, -3) \cup (0, 1]$

(d)  $(4, 6)$

(e)  $(-\infty, -2) \cup (-1/2, \infty)$

(f)  $[-2, -2/3)$

(g)  $[0, 2 - \sqrt{2}) \cup (2 + \sqrt{2}, 6]$

(h)  $\left( \frac{3}{4}, \frac{4}{3} \right)$



**BPPs:** Solve the following meq.s.

Sol<sup>n</sup> ⇒ **(a)**  $\log(x^2 - 2x - 2) \leq 0$

⇒  $x^2 - 2x - 2 \leq 1$

⇒  $x^2 - 2x - 3 \leq 1$

⇒  $(x-3)(x+1) \leq 0$

⇒  $x \in [-1, 3]$  — (i)

&  $x^2 - 2x - 2 \geq 0$


↳  $D = 4 + 8 = 12$

∴  $x = \frac{2 \pm 2\sqrt{3}}{2} = 1 \pm \sqrt{3}$

$(x - (1 + \sqrt{3}))(x - (1 - \sqrt{3})) \geq 0$

∴  $x \in (-\infty, 1 - \sqrt{3}] \cup [1 + \sqrt{3}, \infty)$

(i) ∩ (ii) ⇒



∴  $x \in [-1, 1 - \sqrt{3}, 1 + \sqrt{3}, 3]$

(Ans)

→ **(b)**  $\log_5(x^2 - 11x + 43) < 2$

⇒  $x^2 - 11x + 43 < 25$

⇒  $x^2 - 11x + 18 < 0$

⇒  $(x-9)(x-2) < 0$

⇒  $x \in (2, 9)$

&  $x^2 - 11x + 43 > 0$

↳  $D = 121 - 172 < 0$

always true

$x \in R$

∴  $x \in 2, 9$



$$2 - \log_2(x^2 + 3x) \geq 0$$

$$\Rightarrow \log_2(x^2 + 3x) \leq 2$$

$$\Rightarrow x^2 + 3x \leq 4$$

$$\Rightarrow x^2 + 3x - 4 \leq 0$$

$$\Rightarrow (x+4)(x-1) \leq 0$$

$$\therefore x \in [-4, 1] \quad \text{--- (1)}$$

$$x^2 + 3x > 0$$

$$\Rightarrow x(x+3) > 0$$

$$\Rightarrow x \in (-\infty, -3) \cup (0, \infty)$$

$$\text{Ans. } x \in [-4, -3) \cup (0, 1]$$

Ans.

$$\log_{1.5} \frac{2x-8}{x-2} < 0$$

$$\Rightarrow \frac{2x-8}{x-2} - 1 < 0$$

$$\Rightarrow \frac{x-6}{x-2} < 0$$

$$\Downarrow x \in (2, 6)$$

&

$$\frac{2x-8}{x-2} > 0$$

$$\Rightarrow \frac{x-4}{x-2} > 0 \Rightarrow x \in (-\infty, 2) \cup (4, \infty)$$

$$\text{Ans. } x \in (4, 6)$$



→ (e)  $\log_3 \frac{1+2x}{1+x} < 1$

⇒  $\log \frac{1+2x}{1+x} < 3$

⇒  $\frac{1+2x - 3 - 3x}{1+x} < 0$

⇒  $\frac{x+2}{1+x} > 0$

∴  $x \in (-\infty, -2) \cup (-1, \infty)$  — (i)

&

$\frac{1+2x}{1+x} > 0$

∴  $x \in (-\infty, -1) \cup (-\frac{1}{2}, \infty)$  — (ii)

∴

∴  $x \in (-\infty, -2) \cup (-\frac{1}{2}, \infty)$   
Ans.

→ (f)  $\log_4 \frac{3x+2}{x} \leq 0.5$

⇒  $\frac{3x+2}{x} \leq \sqrt{4} = 2$  &  $\frac{3x+2}{x} > 0$

⇒  $\frac{3x+2 - 2x}{x} \leq 0$

⇒  $\frac{x+2}{x} \leq 0$

∴  $x \in [-2, 0)$

}

∴  $x \in (-\infty, -\frac{2}{3}) \cup (0, \infty)$

∴  $x \in [-2, -\frac{2}{3})$  Ans.




→ Q1)  $\log_4 \frac{3x+2}{x} \cdot \log_2 \left( \frac{x^2-4x+2}{x+1} \right) \leq 1$

$$\Rightarrow \frac{x^2-4x+2}{x+1} \leq 2$$

$$\Rightarrow \frac{x^2-4x+2-2x-2}{x+1} \leq 0$$

$$\Rightarrow \frac{x^2-6x}{x+1} \leq 0$$



$$\Rightarrow \frac{x(x-6)}{x+1} \leq 0$$



&  $\frac{x^2-4x+2}{x+1} > 0$

$D = 16 - 8 = 8$   
 $\therefore x = 2 \pm \sqrt{2}$

$$\Rightarrow \frac{(x - (2 + \sqrt{2}))(x - (2 - \sqrt{2}))}{x+1} > 0$$



Ans  $\Rightarrow \therefore x \in [0, 2 - \sqrt{2}) \cup (2 + \sqrt{2}, 6]$



# Solution to Previous KTKs

## ATDB.uno

## QUESTION [MHT CET 2023 (9 May)]

(KTK 01)



If  $\log_2 x + \log_4 x + \log_8 x + \log_{16} x = \frac{25}{36}$  and  $x = 2^k$  then k is

**A** 1

**B**  $\frac{1}{2}$

**C**  $\frac{1}{3}$

**D**  $\frac{1}{8}$

ATDB.uno

Ans. C

KTK-1.

Q. If  $\log_2 x + \log_4 x + \log_8 x + \log_{16} x = \frac{25}{36}$  and

$x = 2^k$  then:

$$\Rightarrow \log_2 x + \frac{1}{2} \log_2 x + \frac{1}{3} \log_2 x + \frac{1}{4} \log_2 x = \frac{25}{36}$$

let,  $\log_2 x = t$

krish keshri  
jharkhand

$$\Rightarrow t + \frac{t}{2} + \frac{t}{3} + \frac{t}{4} = \frac{25}{36} \Rightarrow \frac{12t + 6t + 4t + 3t}{12} = \frac{25}{36}$$

$$\Rightarrow \frac{25t}{12} = \frac{25}{36} \Rightarrow t = \frac{12}{36} = \frac{1}{3}$$

$$\# \log_2 x = \frac{1}{3}$$

$$\Rightarrow x = (2)^{1/3} \Rightarrow x = 2^k = 2^{1/3}$$

\*  $k = 1/3$  Ans.

$$t = \frac{1}{3}$$

## QUESTION [BITSAT 2021]

(KTK 02)



If  $\log_7 5 = a$ ,  $\log_5 3 = b$  and  $\log_3 2 = c$ , then the logarithm of the number 70 to the base 225 is

**A**  $\frac{1 - a + abc}{2a(1 + b)}$

**B**  $\frac{1 - a - abc}{2a(1 + b)}$

**C**  $\frac{1 + a - abc}{2a(1 + b)}$

**D**  $\frac{1 + a + abc}{2a(1 + b)}$

ATDB.uno

Ans. D



the logarithm of the number 70 to the base

225 is :  $\neq \log_7 5$ ,  $\log_5 3$ ,  $\log_3 2 = \log_7 2$   
 $\hookrightarrow (abc)$

$$\Rightarrow \log_{225} 70 \Rightarrow \frac{\log_2 70}{\log_2 225} \quad \neq \log_5 3, \log_3 2 = \log_5 2$$

$\hookrightarrow (bc)$

$$\Rightarrow \frac{\log_2 (2 \times 5 \times 7)}{\log_2 (3 \times 3 \times 5 \times 5)} \quad \Rightarrow \frac{(bc)(abc) + (abc) + (bc)}{(bc)(abc)}$$

$$\Rightarrow \frac{\log_2 2 + \log_2 5 + \log_2 7}{\log_2 3^2 + \log_2 5^2} \quad \Rightarrow \frac{bc + 2c}{(c)(bc)}$$

$$\Rightarrow \frac{1 + \frac{1}{\log_5 2} + \frac{1}{\log_7 2}}{2 \left( \frac{1}{\log_3 2} + \frac{1}{\log_5 2} \right)} \quad \Rightarrow \frac{(bc) [(abc) + a + 1]}{(bc)(abc)} \times \frac{(c)(bc)}{2c(1+b)}$$

$$\Rightarrow \frac{1 + \frac{1}{bc} + \frac{1}{abc}}{2 \left( \frac{1}{c} + \frac{1}{bc} \right)} \quad \Rightarrow \frac{1 + a + abc}{2a(1+b)} \quad \underline{\text{Ans.}}$$

**krish keshri**  
**iharkhand**



KTK-2) If  $\log_7^5 = a$ ,  $\log_5^3 = b$  and  $\log_3^2 = c$ , then the logarithm of the number 70 to the base 225 is:

$$\log_7^5 \times \log_5^3 \times \log_3^2 = \log_7^2 = abc$$

$$\log_{225}^{70} = \frac{\log_7^{70}}{\log_7^{225}} = \frac{\log_7^7 + \log_7^{10}}{\log_7^{25} + \log_7^9}$$

$$\log_7^5 \times \log_5^3 = \log_7^3 = ab$$

$$= \frac{1 + \log_7^2 + \log_7^5}{2 \log_7^5 + 2 \log_7^3}$$

$$= \frac{1 + abc + a}{2(\log_7^5 + \log_7^3)}$$

**Kriti Mathur  
Raj.**

$$= \frac{1 + abc + a}{2(a + ab)}$$

$$\log_{225}^{70} = \frac{1 + a + abc}{2a(1 + b)} \quad \textcircled{D} \quad \underline{\underline{\text{Ans.}}}$$

## QUESTION [BITSAT 2020]

(KTK 03)



If  $\log_5 \frac{(a + b)}{3} = \frac{\log_5 a + \log_5 b}{2}$ , then  $\frac{a^4 + b^4}{a^2 b^2}$  is equal to

**A** 50

**B** 47

**C** 44

**D** 53

ATDB.uno

Ans. B



KTK-03.

Q If  $\log_5 \frac{(a+b)}{3} = \frac{\log_5 a + \log_5 b}{2}$ , then  $\frac{a^4 + b^4}{a^2 b^2}$  is equal to:

$\Rightarrow 2 \log_5 \frac{(a+b)}{3} = \log_5 (a \cdot b)$

$\Rightarrow \log_5 \left(\frac{(a+b)}{3}\right)^2 = \log_5 (a \cdot b)$

$\Rightarrow a^2 + b^2 + 2ab = 3ab$

$\Rightarrow \boxed{a^2 + b^2 = ab}$   
SBS.

$\Rightarrow a^4 + b^4 + 2a^2 b^2 = 49 a^2 b^2$

$\Rightarrow a^4 + b^4 = 47 a^2 b^2$

$\Rightarrow \boxed{\frac{a^4 + b^4}{a^2 b^2} = 47}$  Ans.

**krish keshri**  
**jharkhand**



KTK-3) If  $\log_5 \left( \frac{a+b}{3} \right) = \frac{\log_5 a + \log_5 b}{2}$ , then  $\frac{a^4 + b^4}{a^2 b^2}$  is equal

to

A.) 50

B.) 47

C.) 44

D.) 53

$$\log_5 \left( \frac{a+b}{3} \right) = \frac{\log_5 a + \log_5 b}{2}$$

$$\log_5 \left( \frac{a+b}{3} \right) = \frac{\log_5 ab}{2}$$

$$2 \log_5 \left( \frac{a+b}{3} \right) = \log_5 ab \Rightarrow \log_5 \left( \frac{a+b}{3} \right)^2 = \log_5 ab$$

**Kriti Mathur  
Raj.**

ATDB.uno

$$\Rightarrow \frac{(a+b)^2}{9} = ab$$

$$\Rightarrow a^2 + b^2 + 2ab - 9ab = 0$$

$$\Rightarrow a^2 + b^2 = 7ab$$

$$\text{Now, } \frac{a^4 + b^4}{a^2 b^2} = \frac{(a^2 + b^2)^2 - 2a^2 b^2}{a^2 b^2}$$

$$= \frac{49a^2 b^2 - 2a^2 b^2}{a^2 b^2} = \frac{47a^2 b^2}{a^2 b^2}$$

$$= 47 \text{ (B) Ans.}$$

## QUESTION [WB JEE 2024]

(KTK 04)



If  $(x^2 \log_x 27) \cdot \log_9 x = x + 4$  then the value of  $x$  is

**A** 2

**B**  $-\frac{4}{3}$

**C** -2

**D**  $\frac{4}{3}$

ATDB.uno

Ans. A



KTK-04.

If  $(x^2 \log_x 27) \cdot \log_9 x = x+4$  then the value of  $x$  is :

$$\Rightarrow x^2 \cdot \frac{\log_3 3^3}{\log_3 x} \cdot \frac{1}{2} \log_3 x = x+4$$

**krish keshri  
jharkhand**

$$\Rightarrow \frac{3x^2}{2} = x+4 \Rightarrow 3x^2 - 2x - 8 = 0$$

$$\Rightarrow 3x^2 - 6x + 4x - 8 = 0$$

$$\Rightarrow 3x(x-2) + 4(x-2) = 0$$

$$\Rightarrow (x-2)(3x+4) = 0$$

$$\Rightarrow x = 2, \frac{-4}{3} \Rightarrow \boxed{x=2} \text{ Ans.}$$

$\left. \begin{array}{l} \text{ } \\ \text{ } \end{array} \right\} \text{REJECT}$



KTK-40) If  $(x^2 \log_a 27) \cdot \log_a x = x+4$  then the value of  $x$  is  
 A.) 2      B.)  $-\frac{4}{3}$       C.) -2      D.)  $\frac{4}{3}$

$$\Rightarrow x^2 \log_a 27 \cdot \log_a x = x+4$$

$$\Rightarrow \frac{x^2 \log_a 27}{\log_a x} = x+4$$

$$\Rightarrow \frac{x^2 \log_a 3^3}{\log_a 3^2} = x+4$$

$$\Rightarrow \frac{x^2 \times 3 \log_a 3}{2 \log_a 3} = x+4$$

$$\Rightarrow 3x^2 = 2x + 8$$

$$3x^2 - 2x - 8 = 0$$

$$3x^2 - 6x + 4x - 8 = 0$$

$$\Rightarrow 3x(x-2) + 4(x-2) = 0$$

$$(x-2)(3x+4) = 0$$

$$\textcircled{A} [x=2] \quad x = -\frac{4}{3} \times$$

Ans.

**Kriti Mathur  
Raj.**

ATDB.uno

**QUESTION [WB JEE 2020]****(KTK 05)**

If  $2 \log(x + 1) - \log(x^2 - 1) = \log 2$ , then  $x =$

- A** only 3
- B** -1 and 3
- C** only -1
- D** 1 and 3

**ATDB.uno****Ans. A**



KTK-05.

If  $2 \log (x+1) - \log (x^2-1) = \log 2$ , then  $x =$  .

$$\Rightarrow 2 \log \frac{(x+1)}{(x^2-1)} = \log 2$$

$$\Rightarrow \frac{(x+1)^2}{(x+1)(x-1)} = \log 2$$

$$\Rightarrow x+1 = 2x-2$$

$$\Rightarrow \boxed{x=3} \text{ Ans.}$$

**krish keshri**  
**jharkhand**



KTK-5.) If  $2 \log(x+1) - \log(x^2-1) = \log 2$ , then  $x =$

$$\log(x+1)^2 - \log(x^2-1) = \log 2$$

$$\log\left(\frac{(x+1)^2}{x^2-1}\right) = \log 2$$

$$\frac{(x+1)^2}{x^2-1} = 2$$

$$\frac{(x+1)^2}{(x+1)(x-1)}$$

$$\frac{(x+1)}{(x-1)} = 2 ; x \neq -1$$

$$x+1 = 2x-2$$

$$0 = 2x - x - 2 - 1$$

$$0 = x - 3 \Rightarrow \underline{\underline{[x = 3]}} \underline{\underline{\text{Ans.}}}$$

**Kriti Mathur**  
**Raj.**

ATDB.uno

## QUESTION [WB JEE 2018]

(KTK 06)



If  $x + \log_{10}(1 + 2^x) = x \log_{10} 5 + \log_{10} 6$ , then the value of  $x$  is

**A**  $\frac{1}{2}$

**B**  $\frac{1}{3}$

**C** 1

**D** 2

ATDB.uno

Ans. C



KTK-OG

Q. If  $x + \log_{10}(1+2^x) = x \log_{10} 5 + \log_{10} 6$ ,  
then the value of  $x$  is.

$$\Rightarrow x \log_{10} 10 + \log_{10}(1+2^x) = \log_{10} 5^x + \log_{10} 6$$

$$\Rightarrow \log_{10} [10^x (1+2^x)] = \log_{10} (5^x \cdot 6)$$

$$\Rightarrow 10^x \cdot (1+2^x) = 5^x \cdot 6$$

$$\Rightarrow 2^x (1+2^x) = 6$$

ATDB.uno

let,  $2^x = t$

krish keshri  
jharkhand

$$\Rightarrow t(1+t) = 6$$

$$\Rightarrow t^2 + t - 6 = 0 \Rightarrow (t+3)(t-2) = 0$$

$$\Rightarrow t = -3, 2$$

$$\Rightarrow 2^x = -3 \quad | \quad \Rightarrow 2^x = 2$$

$$\underline{\text{NOT POSSIBLE}} \quad | \quad \Rightarrow \boxed{x=1} \quad \underline{\text{Ans}}$$



KTK-6) If  $x + \log_{10}(1+2^x) = x \log_{10} 5 + \log_{10} 6$ , then the value of  $x$  is

Ans.  $x + \log_{10}(1+2^x) = \log_{10} 5^x + \log_{10} 6$

A.) 1/2

B.) 1/3

C.) 1

D.) 2

$$x + \log_{10}(1+2^x) = \log_{10} 5^x \cdot 6$$

$$\Rightarrow x = \log_{10} \frac{5^x \cdot 6}{1+2^x}$$

ATDB.uno

**Kriti Mathur  
Raj.**

$$\Rightarrow \log_{10} \left( \frac{5^x \cdot 6}{2^x + 1} \right) = x$$

$$\frac{5^x \cdot 6}{2^x + 1} = 10^x$$

Now, by hit & trial

$$\text{at } x = 1 \Rightarrow \frac{5 \cdot 6}{2+1} = \frac{30}{3} \Rightarrow 10 = 10 \quad \text{C) } \underline{\underline{\text{Ans.}}}$$

[x=1]

## QUESTION [WB JEE 2019]

(KTK 07)



If  $\log_2 6 + \frac{1}{2x} = \log_2 (2^{\frac{1}{x}} + 8)$ , then the value of x are

**A**  $\frac{1}{4}, \frac{1}{3}$

**B**  $\frac{1}{4}, \frac{1}{2}$

**C**  $-\frac{1}{4}, \frac{1}{2}$

**D**  $\frac{1}{3}, -\frac{1}{2}$

ATDB.uno

Ans. B



KTK-07.

Q. If  $\log_2 6 + \frac{1}{2x} = \log_2 (2^{1/x} + 8)$ , then the

value of  $x$  are : write:  $\frac{1}{2x} = \log_2 (2^{1/2x})$

$$\Rightarrow \log_2 (6 \cdot 2^{1/2x}) = \log_2 (2^{1/x} + 8)$$

$$\Rightarrow 6 \cdot 2^{1/2x} = 2^{1/x} + 8 \quad \text{let; } 2^{1/x} = t$$

$$\Rightarrow 6 t^{1/2} = t + 8$$

SBS

ATDB.uno

krish keshri  
jharkhand

$$\Rightarrow 36 t \quad t^2 + 16 t + 64$$

$$\Rightarrow t^2 - 20 t + 64 = 0$$

$$\Rightarrow t^2 - 16 t - 4 t + 64 = 0$$

$$\Rightarrow t(t-16) - 4(t-16) = 0$$

$$\Rightarrow (t-16)(t-4) = 0$$

$$\Rightarrow t = 16, 4.$$

$$\Rightarrow 2^{1/x} = 16 \quad | \quad \Rightarrow 2^{1/x} = 4$$

$$\Rightarrow \frac{1}{x} = 4 \quad | \quad \Rightarrow \frac{1}{x} = 2$$

$$\Rightarrow \boxed{x = \frac{1}{4}} \quad | \quad \Rightarrow \boxed{x = \frac{1}{2}}$$

Ans.

KTK-7.) If  $\log_2 6 + \frac{1}{2x} = \log_2 (2^{\frac{1}{2}} + 8)$ , then the value of  $x$  are.

A.)  $\frac{1}{4}, \frac{1}{3}$

~~B.)  $\frac{1}{4}, \frac{1}{2}$~~

C.)  $-\frac{1}{4}, \frac{1}{2}$

D.)  $\frac{1}{3}, -\frac{1}{2}$

$$\log_2 6 + \frac{1}{2x} = \log_2 (2^{\frac{1}{2}} + 8)$$

$$\log_2 6 + \log_2 2^{\frac{1}{2x}} = \log_2 (2^{\frac{1}{2}} + 8)$$

$$\log_2 (6 \times 2^{\frac{1}{2x}}) = \log_2 (2^{\frac{1}{2}} + 8)$$

$$6 \times 2^{\frac{1}{2x}} = 2^{\frac{1}{2}} + 8$$

$$\text{Let } 2^{\frac{1}{2x}} = t, \quad t^2 = 2^{\frac{1}{x}}$$

$$6t = t^2 + 8$$

$$t^2 - 6t + 8 = 0$$

$$t^2 - 4t - 2t + 8 = 0$$

$$t(t-4) - 2(t-4) = 0$$

$$t = 2$$

$$2^{\frac{1}{2x}} = 2$$

$$\frac{1}{2x} = 1$$

$$2x = 1$$

$$\left[ x = \frac{1}{2} \right] \underline{\underline{\text{Ans.}}}$$

$$t = 4$$

$$2^{\frac{1}{2x}} = 2^2$$

$$\frac{1}{2x} = 2$$

$$\left[ x = \frac{1}{4} \right] \underline{\underline{\text{Ans.}}}$$

$$\textcircled{B} \underline{\underline{\text{Ans.}}}$$

**Kriti Mathur  
Raj.**

ATDB.uno



**QUESTION [WB JEE 2017]****(KTK 08)**

If  $(\log_5 x)(\log_x 3x)(\log_{3x} y) = \log_x x^3$ , then  $y$  equals

**A** 125

**B** 25

**C** 513

**D** 243

**ATDB.uno****Ans. A**



KTK-08.

If  $(\log_5 x) (\log_x 3x) (\log_{3x} y) = \log_x x^3$  then

$y$  equals :

$$\Rightarrow \log_5 x \cdot \log_x 3x \cdot \log_{3x} y = 3$$

$$\Rightarrow \log_5 y = 3$$

$$\Rightarrow y = 5^3 \Rightarrow \boxed{y = 125} \text{ Ans.}$$

krish keshri  
jharkhand

## QUESTION [COMEDK 2023]

(KTK 09)



The value of  $a^{\log_b c} - c^{\log_b a}$ , where  $a, b, c > 0$  but  $a, b, c \neq 1$ , is

**A** a

**B** b

**C** c

**D** 0

ATDB.uno

Ans. D



KTK-09.

The value of  $a^{\log_b c} - c^{\log_b a}$ , where  $a, b, c > 0$

but  $a, b, c \neq 1$ , is:

$\Rightarrow a^{\log_b c} - a^{\log_b c}$  interchange.

$\Rightarrow 0$  Ans.

**QUESTION [COMEDK 2022]****(KTK 10)**

The value of  $3^{\log_4 5} - 5^{\log_4 3}$  is

**A** 0

**B** 1

**C** 2

**D** 4

**ATDB.uno****Ans. A**



KTK-10, The value of  $3^{\log_4 5} - 5^{\log_4 3}$  is :

$$\Rightarrow 3^{\log_4 5} - 3^{\log_4 5} \leftarrow \text{interchange.}$$

$$\Rightarrow 0 \text{ Ans.}$$

## QUESTION [COMEDK 2021]

(KTK 11)



$8^3 \log_8 5$  is equal to

**A**  $\log_8 25$

**B** 120

**C** 125

**D**  $\log_8 15$

ATDB.uno

Ans. C



KTK-11.

$8^3 \log_8 5$  is equal to :

$$\Rightarrow 8^{\log_8 5^3}$$

$$\Rightarrow 5^3 = 125 \text{ Ans.}$$

krish keshri  
jharkhand

## QUESTION [COMEDK 2020]

(KTK 12)



$7^{2 \log_7 5}$  is equal to

- A** 5
- B**  $\log_7 35$
- C**  $\log_7 25$
- D** 25

ATDB.uno

Ans. D



KTK-12.

$7^2 \log_7 5$  is equal to:

$$\Rightarrow 7 \log_7 5^2$$

$$\Rightarrow 5^2 = 25 \text{ Ans.}$$

## QUESTION

(KTK 13)



Find the exhaustive solutions set of  $\frac{(x^2-9)^{101}(x^2+6)(x^2-4)^{100}}{(x^2-5x+6)^{13}(x^2-16)^{16}} > 0$ .

ATDB.uno

Ans.  $(-\infty, -3) \cup (2, \infty) - \{\pm 4, 3\}$



KTK-13) Find the exhaustive solutions set of  $\frac{(x^2+9)^{101} (x^2+6) (x^2-4)^{100}}{(x^2-5x+6)^{13} (x^2-16)^{16}} > 0$ .

$$\Rightarrow \frac{(x+3)^{101} (x-3)^{101} (x^2+6)}{(x^2-5x+6)^{13}} > 0, \quad x \neq \pm 2, \pm 4$$


$$\Rightarrow \frac{(x+3)^{101} (x-3)^{101} (x^2+6)}{(x-2)^3 (x-3)^3} > 0, \quad x \neq \pm 2, \pm 4$$

$$\Rightarrow \frac{(x+3)^{101} (x-3)^{89} (x^2+6)^{atongkatul}}{(x-2)^3} > 0, \quad x \neq \pm 2, \pm 4, 3$$

ATDB.uno

**Kriti Mathur**  
**Raj.**

$$\Rightarrow \frac{(x+3)^{101}}{(x-2)^3} > 0$$



$$x \in (-\infty, -3) \cup (2, \infty) - \{\pm 4, 3\} \underline{\underline{\text{Ans.}}}$$

## QUESTION

(KTK 14)



Find the exhaustive solutions set of  $\frac{(x-4)^{30}(x^2-9)^9(x^2-3x+2)^{17}(3x^2+10)^{10}}{(x^2-5x+6)^{52}(x^2-25)^{60}(x^2+10)^{11}} \leq 0$ .

ATDB.uno

Ans.  $[-3, 1] \cup (2, 3)$



KTK-14) Find the exhaustive solutions set of

$$\frac{(x-4)^{30} (x^2-9)^9 (x^2-3x+2)^{17} (3x^2+10)^{10}}{(x^2-5x+6)^{52} (x^2-25)^{60} (x^2+10)^{11}} \leq 0$$


$$\Rightarrow \frac{(x+3)^9 (x-3)^9 (x^2-3x+2)^{17}}{(x^2-5x+6)^{52} (x^2+10)^{11}} \leq 0; x \neq \pm 5, x=4$$

$$\Rightarrow \frac{(x+3)^9 (x-3)^9 (x-2)^{17} (x-1)^{17}}{(x-2)^{52} (x-3)^{42}} \leq 0; x \neq \pm 5$$

$$\Rightarrow \frac{(x+3)^9 (x-1)^{17}}{(x-2)^{35} (x-3)^{42}} \leq 0; x \neq \pm 5$$

**Kriti Mathur**

**Raj.**



$$x \in [-3, 1] \cup (2, 3) \cup \{4\}$$

## QUESTION

(KTK 15)



Solve in real numbers the equation  $\sqrt{x} + \sqrt{y} + 2\sqrt{z-2} + \sqrt{u} + \sqrt{v} = x + y + z + u + v$ .

$$0 = x - 2 \cdot \sqrt{x} \cdot \frac{1}{2} + \frac{1}{4} + y - 2\sqrt{y} \cdot \frac{1}{2} + \frac{1}{4} + u - 2\sqrt{u} \cdot \frac{1}{2} + \frac{1}{4} + v - 2\sqrt{v} \cdot \frac{1}{2} + \frac{1}{4} + \sqrt{z-2}^2 - 2\sqrt{z-2} + 1 + z - \frac{1}{4} - \frac{1}{4} - \frac{1}{4} - \frac{1}{4} - 1$$

$$0 = (\sqrt{x} - 1/2)^2 + (\sqrt{y} - 1/2)^2 + (\sqrt{u} - 1/2)^2 + (\sqrt{v} - 1/2)^2 + (\sqrt{z-2} - 1)^2 = 0$$

ATDB.uno

$$x = 1/4 = y = u = v$$

$$z = 3$$

Ans.  $x = y = u = v = 1/4, z = 3$

**KTK-15**

Solve in real no.s, the eqn

$$\sqrt{x} + \sqrt{y} + 2\sqrt{z-2} + \sqrt{u} + \sqrt{v} = x + y + z + u + v.$$

KTK 15

by Reed

from WB

Soln

$$x + y + z + u + v = \sqrt{x} + \sqrt{y} + 2\sqrt{z-2} + \sqrt{u} + \sqrt{v}$$

$$\Rightarrow x - \sqrt{x} + y - \sqrt{y} + z - 2\sqrt{z-2} + u - \sqrt{u} + v - \sqrt{v} = 0$$

$$\Rightarrow x - 2 \cdot \sqrt{x} \cdot \frac{1}{2} + \left(\frac{1}{2}\right)^2 + y - 2\sqrt{y} \cdot \frac{1}{2} + \left(\frac{1}{2}\right)^2 + (z-2) - 2\sqrt{z-2} \cdot 1 + 1^2$$

$$+ u - 2\sqrt{u} \cdot \frac{1}{2} + \left(\frac{1}{2}\right)^2 + v - 2\sqrt{v} \cdot \frac{1}{2} + \left(\frac{1}{2}\right)^2$$

$$= \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^2 + 1 + \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^2$$

$$\Rightarrow \left(\sqrt{x} - \frac{1}{2}\right)^2 + \left(\sqrt{y} - \frac{1}{2}\right)^2 + \left(\sqrt{z-2} - 1\right)^2 + \left(\sqrt{u} - \frac{1}{2}\right)^2 + \left(\sqrt{v} - \frac{1}{2}\right)^2$$

$$= \frac{1}{4} + \frac{1}{4} + 1 + \frac{1}{4} + \frac{1}{4} - 2 = 2 - 2 = 0.$$

$$\Rightarrow \left. \begin{array}{l} \sqrt{x} = \frac{1}{2} \\ x = \frac{1}{4} \end{array} \right| \left. \begin{array}{l} \sqrt{y} = \frac{1}{2} \\ y = \frac{1}{4} \end{array} \right| \left. \begin{array}{l} \sqrt{z-2} = 1 \\ z-2 = 1 \\ z = 3 \end{array} \right| \left. \begin{array}{l} \sqrt{u} = \frac{1}{2} \\ u = \frac{1}{4} \end{array} \right| \left. \begin{array}{l} \sqrt{v} = \frac{1}{2} \\ v = \frac{1}{4} \end{array} \right.$$

$$\therefore \boxed{x = y = u = v = \frac{1}{4}} \text{ \& \ } \boxed{z = 3}$$



**QUESTION****(KTK 16)**

Find all pair of positive integer  $(m, h)$  that satisfy  $mn + 3m - 8n = 59$ .

**ATDB.uno****Ans. 3**



KTK-16) Find all pair of positive integers  $(m, n)$  that satisfy

$$mn + 3m - 8n = 59$$

Ans.  $m(n+3) - 8n = 24 + 35$

$$m(n+3) - 8n - 24 = 35$$

$$m(n+3) - 8(n+3) = 35$$

$$(m-8)(n+3) = 35$$

$$59$$

$$\underline{24}$$

$$35$$

$(1 \times 35, 5 \times 7,$

$7 \times 5, 35 \times 1$

**Kriti Mathur**  
**Raj.**

ATDB.uno

$$m-8$$

$$n+3$$

$$(m, n)$$

$$1$$

$$35$$

$$(9, 32)$$

$$5$$

$$7$$

$$(13, 4)$$

$$7$$

$$5$$

$$(15, 2)$$

$$35$$

$$1$$

$$(43, -2) \times$$

= 3 pairs

==



**KTK-161** Find all pair of +ve integer  $(m, n)$  that  
 $mn + 3m - 8n = 59.$

Soln  $mn + 3m - 8n = 59$

a.  $m(n+3) - 8n - 24 = 59 - 24$

a.  $m(n+3) - 8(n+3) = 35$

a.  $(m-8)(n+3) = 35.$

now,  $35 = 5 \times 7$

$35 = 1 \times 35$

✓

$35 = (-5) \times (-7)$

$35 = (-1) \times (-35)$

~~ATDB.uno~~

no need since  $m, n \in \mathbb{I}^+$

$m-8 =$	5	7	1	35
$n+3 =$	7	5	35	1
$\Rightarrow m =$	13	15	9	43
$\& n =$	4	2	32	-2

✗  $(\because m, n \in \mathbb{I}^+)$

$\therefore$  all pair of positive integers  $(m, n) \in (13, 4), (15, 2), (9, 32)$   
 (Ans.)

$\therefore$  no. of pairs of  $(m, n)$  is = 3.

## QUESTION

(KTK 17)



The least value of the expression  $(x + y)(y + z)$  where given that  $x, y, z > 0$  and  $xyz(x + y + z) = 1$

ATDB.uno

Ans. 2

KTK-170) The least value of the expression  $(x+y)(y+z)$  where given that  $x, y, z > 0$  and  $xyz(x+y+z) = 1$

Ans.  $x+y+z = \frac{1}{xyz}$

**Kriti Mathur  
Raj.**

$$(x+y)(y+z)$$

$$xy + xz + y^2 + yz$$

$$= y(x+y+z) + xz$$

ATDB.uno

$$\Rightarrow y \left( \frac{1}{xyz} \right) + xz$$

$$\Rightarrow \frac{1}{xz} + xz$$

$$\text{As } \frac{1}{xz} + xz \geq 2$$

Least value = 2 Ans.



Date: / /  
Page: /



**KTK-17!** The least value of the expression  
 $(x+y)(y+z)$  where given that  $x, y, z > 0$ .

and  $xyz(x+y+z) = 1$ .

Soln

$$E = (x+y)(y+z)$$

$$\Rightarrow E = xy + xz + yz + yz$$

$$\Rightarrow E = y(x+y+z) + xz$$

$$\Rightarrow E = \frac{y}{xyz} + xz \quad \neq y \neq 0$$

$$\Rightarrow E = \frac{1}{xz} + xz$$

$$\therefore E_{\min} \Big|_{\text{at } x=z=1} = 2$$

$$\therefore \boxed{E_{\min} = 2} \text{ Ans.}$$

$$\left. \begin{aligned} xyz(x+y+z) &= 1 \\ \Rightarrow (x+y+z) &= \frac{1}{xyz} \end{aligned} \right\}$$

KTK 17  
 by Reed  
 from WB



\* Read class Theory

\* Retry class Questions In rough COPY

\* TAH

\* BPP

\* KTK

\* DPP / Module.

**ATDB.uno**



**THANK**  
ATDB.uno

**YOU**