

PRAVIAS

JEE 2026

Mathematics

Sequence and Series

Lecture - 3

By – Ashish Agarwal Sir
(IIT Kanpur)



Topics *to be covered*



A Arithmetic Mean

B Geometric Progression



Recap *of previous lecture*



1. K^{th} term from end in on A.P. = $(n - K + 1)^{\text{th}}$ term from beginning.

2. If $T_n = 2n + 3$ then the sequence is an A.P

3. $30 + \frac{89}{3} + \frac{88}{3} + \dots$ has largest possible sum then number of terms that should be taken is _____

Recap *of previous lecture*



4. ✓ If a sequence has 40 terms then number of even numbered terms is 20
- 41 terms then no: of even numbered terms = 20
odd numbered terms = 21
5. ✓ If a sequence has 49 terms then number of even numbered terms is 24 &
number of odd numbered terms is 25.

Recap

of previous lecture



6. Three numbers in A.P. $a-d, a, a+d$ — $c.d = d$

Four numbers in A.P. $a-3d, a-d, a+d, a+3d$ — $c.d = 2d$

Five numbers in A.P. $a-2d, a-d, a, a+d, a+2d$ — $c.d = d$

Six numbers in A.P. $a-5d, a-3d, a-d, a+d, a+3d, a+5d$ — $c.d = 2d$

FIRST
TERM
IS NOT
 a

7. If we pick terms in an A.P. at an interval of 5 then the picked terms will form an A.P with common difference $5d$

Recap *of previous lecture*



8. If $a_1, a_2, a_3, a_4, a_5, \dots$ be an A.P. with common difference 'd' then $a_1, a_5, a_9, a_{13}, \dots$ will be in A.P. with common difference $4d$. Similarly $a_2, a_4, a_6, \dots$ will also be in A.P. with common difference $2d$.
9. If $m, n, p, q \in \mathbb{N}$ & $m + n = p + q$ then in an A.P. we have $T_m + T_n = T_p + T_q$.
10. If $a_1 + a_{12} = 10$ in an A.P. then value of $a_3 + a_7 + a_9 + a_4 + a_6 + a_{10}$ is $3(a_1 + a_{12}) = 30$.

Homework Discussion

Suppose $a_1, a_2, \dots, a_n, \dots$ be an arithmetic progression of natural numbers. If the ratio of the sum of first five terms to the sum of first nine terms of the progression is $5 : 17$ and, $110 < a_{15} < 120$, then the sum of the first ten terms of the progression is equal to

- A** 290
- B** 380
- C** 460
- D** 510

Consider an A.P. of positive integers, whose sum of the first three terms is 54 and the sum of the first twenty terms lies between 1600 and 1800. Then its 11th term is :

- A** 108
- B** 90
- C** 122
- D** 84

QUESTION [JEE Advanced 2018]

Tah03



Let X be the set consisting of the first 2018 terms of the arithmetic progression 1, 6, 11,, and Y be the set consisting of the first 2018 terms of the arithmetic progression 9, 16, 23, Then, the number of elements in the set $X \cup Y$ is

(A) 3857

QUESTION [JEE Mains 2024 (27 Jan)]



Tan04

The number of common terms in the progressions

4, 9, 14, 19,, up to 25th term and 3, 6, 9, 12,, up to 37th term is:

- A** 9
- B** 8
- C** 5
- D** 7

Ans. D

The sum of the common terms of the following three arithmetic progressions.

3, 7, 11, 15, ..., 399,
2, 5, 8, 11, ..., 359 and
2, 7, 12, 17, ..., 197,
is equal to _____

(ADIBST)

QUESTION [JEE Mains 2025 (4 April)]

Tah06



Let $A = \{1, 6, 11, 16, \dots\}$ and $B = \{9, 16, 23, 30, \dots\}$ be the sets consisting of the first 2025 terms of two arithmetic progressions. Then $n(A \cup B)$ is

- A** 3814
- B** 4003
- C** 4027
- D** 3761

Ans. D

QUESTION [JEE Mains 2021 (31 Aug)]

Tan07



Let $a_1, a_2, a_3, \dots$ be an A.P. If $\frac{a_1 + a_2 + \dots + a_{10}}{a_1 + a_2 + \dots + a_p} = \frac{100}{p^2}$, $p \neq 10$, then $\frac{a_{11}}{a_{10}}$ is equal to :

- ☐ A $\frac{19}{21}$
- ☐ B $\frac{100}{121}$
- ☒ C $\frac{21}{19}$
- ☐ D $\frac{121}{100}$

$$\frac{\frac{10}{2}(2a+9d)}{\frac{p}{2}(2a+(p-1)d)} = \frac{100}{p^2}$$

$$\frac{2a+9d}{2a+(p-1)d} = \frac{10}{p}$$

$$2ap + 9dp = 20a + 10dp - 10d$$

$$2a(p-10) = dp - 10d$$

$$2a(p-10) = d(p-10)$$

$$2a = d$$

$$\frac{a+10d}{a+9d} = \frac{a+20a}{a+18a} = \frac{21}{19}$$

Ans. C

For $a, b \in \mathbb{R} - \{0\}$, let $f(x) = ax^2 + bx + a$ satisfies $f\left(x + \frac{7}{4}\right) = f\left(\frac{7}{4} - x\right) \forall x \in \mathbb{R}$. Also the equation $f(x) = 7x + a$ has only one real and distinct solution.

The value of $(a + b)$ is equal to

- ☐ A 4
- ☒ B 5
- ☐ C 6
- ☐ D 7

$$\star f\left(\frac{7}{4} + x\right) = f\left(\frac{7}{4} - x\right)$$

f is symmt about $x = \frac{7}{4}$

$$f(x) = ax^2 + bx + a$$

symmt abt $x = -\frac{b}{2a}$

$$-\frac{b}{2a} = \frac{7}{4}$$

$f(x) = 7x + a$ has only one real & distinct soln.

$$-2b = 7a$$

$$ax^2 + bx + a = 7x + a$$

$$ax^2 + (b-7)x = 0$$

$$D = (b-7)^2 - 4a \cdot 0 = 0$$

$$b = 7$$

$$-14 = 7a$$

$$a = -2$$

$$a + b = 5$$

Ans. B

For $a, b \in \mathbb{R} - \{0\}$, let $f(x) = ax^2 + bx + a$ satisfies $f\left(x + \frac{7}{4}\right) = f\left(\frac{7}{4} - x\right) \forall x \in \mathbb{R}$. Also the equation $f(x) = 7x + a$ has only one real and distinct solution.

The minimum value $f(x)$ in $\left[0, \frac{3}{2}\right]$ is equal to

A $-\frac{33}{8}$

B 0

C 4

D -2

$$\begin{aligned} f(x) &= -2x^2 + 7x - 2 \\ &= -2\left(x^2 - \frac{7}{2}x + \frac{49}{16} - \frac{49}{16}\right) - 2 \\ &= -2\left(x - \frac{7}{4}\right)^2 + \frac{49}{8} - 2 \\ &= \frac{33}{8} - 2\left(x - \frac{7}{4}\right)^2 \end{aligned}$$

**Aao Machaay Dhamaal
Deh Swaal pe Deh Swaal**

Arithmetic Mean

* If a, b, c are in A.P then 'b' is called single A.M b/w a & c

$$b - a = c - b$$

$$2b = a + c \rightarrow b = \frac{a + c}{2}$$

$$a, b, c \text{ are in A.P} \iff 2b = a + c$$

* If $a, A_1, A_2, A_3, \dots, A_n, b$ are in A.P then $A_1, A_2, \dots, A_n$ are called n A.Ms b/w a & b .

$$T_1 = a$$

$$T_{n+2} = b \Rightarrow b = a + (n+1)d$$

$$d = \frac{b - a}{n+1}$$

$$A_1 = a + d$$

$$A_2 = a + 2d$$

$$A_3 = a + 3d$$

$$\vdots$$

$$A_n = a + nd$$

$$A_1 + A_2 + \dots + A_n = na + d(1 + 2 + 3 + \dots + n)$$

$$\sum_{i=1}^n A_i = na + d \cdot \frac{n(n+1)}{2} = na + \frac{b-a}{n+1} \cdot \frac{n(n+1)}{2}$$

$$\sum_{i=1}^n A_i = na + \frac{n(b-a)}{2} = n \left(\frac{2a+b-a}{2} \right) = n \left(\frac{a+b}{2} \right)$$

$a, A_1, A_2, \dots, A_n, b$ are in A.P

n A.Ms b/w a & b .

$$\sum_{i=1}^n A_i = n \left(\frac{a+b}{2} \right)$$

Sum of n A.Ms b/w 2 No:s = $n \cdot (\text{single A.M b/w the two no:s})$

Ex: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20

8 A.Ms b/w 2 & 20

$$\text{Sum of 8 A.Ms} = 4 + 6 + 8 + 10 + 12 + 14 + 16 + 18 = 88 = 8 \cdot \left(\frac{2+20}{2} \right)$$

QUESTION



Insert 20 AM's between 4 and 67.

$\xrightarrow{d}$
 $4, A_1, A_2, \dots, A_{20}, 67$ are in A.P. $\xleftarrow{-d}$

$$T_{22} = 67 = 4 + 21d$$

$$d = \frac{63}{21} = 3$$

20 A.Ms: 7, 10, 13, -----, 61, 64

QUESTION



If p arithmetic mean are inserted between 5 and 41 so that ratio $\frac{A_3}{A_{p-1}} = \frac{2}{5}$, then find the value of p .

M ① 5, $A_1, A_2, \dots, A_p, 41$ are in A.P

$$T_4 = A_3 = 5 + 3d \quad T_{p+2} = 41$$

$$T_p = A_{p-1} = 5 + (p-1)d \quad 5 + (p+1)d = 41$$

$$(p+1)d = 36.$$

$$\frac{A_3}{A_{p-1}} = \frac{5+3d}{5+(p-1)d} = \frac{2}{5}.$$

$$25 + 15d = 10 + 2(p-1)d$$

$$25 + 15 \cdot \frac{36}{p+1} = 10 + 2(p-1) \cdot \frac{36}{p+1}$$

$$15 + \frac{15 \cdot 36}{p+1} = \frac{72(p-1)}{p+1} \Rightarrow 15p + 15 + 540 = 72p - 72$$

$$\begin{aligned} 57p &= 627 \\ p &= 11 \end{aligned}$$

If p arithmetic mean are inserted between 5 and 41 so that ratio $\frac{A_3}{A_{p-1}} = \frac{2}{5}$, then find the value of p .

M② $\xrightarrow{d}$ $5, A_1, A_2, \dots, A_p, 41$ are in A.P. $\xleftarrow{-d}$

$$\frac{A_3}{A_{p-1}} = \frac{2}{5}$$

$$T_{p+2} = 41$$

$$5 + (p+1)d = 41$$

$$(p+1)d = 36$$

$$\frac{5+3d}{41-2d} = \frac{2}{5}$$

$$25 + 15d = 82 - 4d$$

$$19d = 57$$

$$d = 3$$

$$p+1 = 12$$

$$p = 11$$

QUESTION



If $A_1, A_2, A_3, \dots, A_{51}$ are arithmetic means inserted between the numbers a and b , then

find the value of $\left(\frac{b+A_{51}}{b-A_{51}}\right) - \left(\frac{A_1+a}{A_1-a}\right)$.

$$\underbrace{a, A_1, A_2, \dots, A_{51}, b}_{T_{53} = b = a + 52d} \quad b - a = 52d$$

$$\begin{aligned} \text{Let } E &= \frac{b+A_{51}}{b-A_{51}} - \frac{(A_1+a)}{A_1-a} \\ &= \frac{b+A_{51}}{d} - \frac{A_1+a}{d} \\ &= \frac{b+A_{51}-A_1-a}{d} \\ &= \frac{b+b-d-(a+d)-a}{d} \\ &= \frac{2b-2a-2d}{d} = \frac{104d-2d}{d} = 102 \text{ Ans} \end{aligned}$$

QUESTION



If $a_1 a_2 a_3 \dots a_n$ are n arithmetic means inserted between 7 and 2015 whose sum is 56616 then

$$7, a_1, a_2, \dots, a_n, 2015.$$

$$a_1 + a_2 + \dots + a_n = 56616.$$

- A** n is 56
- B** n is 28
- C** $a_{19} = 2029/3$
- D** $a_{19} = 36543/57$

QUESTION



If $S_1, S_2, S_3, \dots, S_p$ are the sums of n terms of ' p ' arithmetic series whose first term and 1, 2, 3, 4, and whose common difference are 1, 3, 5, 7,

Prove that $S_1 + S_2 + S_3 + \dots + S_p = \frac{np}{2} (np + 1)$


$$A.P_1: T_1=1, d=1 \rightarrow S_1 = \frac{n}{2} (2 \cdot 1 + (n-1) \cdot 1)$$

$$A.P_2: T_1=2, d=3 \rightarrow S_2 = \frac{n}{2} (2 \cdot 2 + (n-1) \cdot 3)$$

$$A.P_3: T_1=3, d=5 \rightarrow S_3 = \frac{n}{2} (2 \cdot 3 + (n-1) \cdot 5)$$

$$A.P_p: T_1=p, d=2p-1 \rightarrow S_p = \frac{n}{2} (2p + (n-1)(2p-1))$$

$$\begin{aligned} S_1 + S_2 + S_3 + \dots + S_p &= \frac{n}{2} (2(1+2+3+\dots+p) + (n-1)(1+3+5+\dots+(2p-1))) \\ &= \frac{n}{2} (2 \cdot \frac{p(p+1)}{2} + (n-1)p^2) \end{aligned}$$





$$s_1 + s_2 + \dots + s_p = \frac{np}{2} (p+1 + p(n-1))$$

$$s_1 + s_2 + \dots + s_p = \frac{np}{2} (np+1)$$

QUESTION [JEE Mains 2023 (Jan)]



Number of 4 digit integers less than 2800 which are either divisible by 3 or by 11 is equal to

$$n(D_3 \cup D_{11}) = n(D_3) + n(D_{11}) - n(D_3 \cap D_{11})$$

$$D_3 = \{1002, 1005, \dots, 2799\} \quad \text{---} \quad 2799 = 1002 + (n-1) \cdot 3 \quad \text{---} \quad n = 600$$

$$D_{11} = \{1001, 1012, 1023, \dots, 2794\} \quad \text{---} \quad 2794 = 1001 + (n-1) \cdot 11 \quad \text{---} \quad n = 164$$

$$D_{33} = \{1023, 1056, \dots, 2772\} \quad \text{---} \quad 2772 = 1023 + (n-1) \cdot 33 \quad \text{---} \quad n = 54$$


$$\begin{aligned} n(D_3 \cup D_{11}) &= 600 + 164 - 54 \\ &= 600 + 110 = 710 \text{ Ans} \end{aligned}$$

The sides of a right angled triangle are in arithmetic progression. If the triangle has area 24, then what is the length of its smallest side?

$$\frac{1}{2} \cdot (a-d) \cdot a = 24$$

$$a(a-d) = 48$$

$$a^2 - ad = 48$$



Consider two sets A and B, each containing three numbers in A.P. Let the sum and the product of the elements of A be 36 and p respectively and the sum and the product of the elements of B be 36 and q respectively. Let d and D be the common differences of A.P.'s in A and B respectively such that $D = d + 3, d > 0$. If $\frac{p+q}{p-q} = \frac{19}{5}$, then $p - q$ is equal to

- A** 540
- B** 450
- C** 600
- D** 630

QUESTION [JEE Mains 2023 (31 Jan)]



Let $a_1, a_2, \dots, a_n$ be in A.P. If $a_5 = 2a_7$ and $a_{11} = 18$, then

$12 \left(\frac{1}{\sqrt{a_{10}} + \sqrt{a_{11}}} + \frac{1}{\sqrt{a_{11}} + \sqrt{a_{12}}} + \dots + \frac{1}{\sqrt{a_{17}} + \sqrt{a_{18}}} \right)$ is equal to

$$\begin{aligned} a_{11} - 6d &= 2(a_{11} - 4d) \\ 18 - 6d &= 2(18 - 4d) \\ 9 - 3d &= 18 - 4d \\ d &= 9 \end{aligned}$$

$$12 \left(\frac{\sqrt{a_{11}} - \sqrt{a_{10}}}{a_{11} - a_{10}} + \frac{\sqrt{a_{12}} - \sqrt{a_{11}}}{a_{12} - a_{11}} + \dots + \frac{\sqrt{a_{18}} - \sqrt{a_{17}}}{a_{18} - a_{17}} \right)$$

$$\frac{12}{d} \left((\sqrt{a_{11}} - \sqrt{a_{10}}) + (\sqrt{a_{12}} - \sqrt{a_{11}}) + (\sqrt{a_{13}} - \sqrt{a_{12}}) + \dots + (\sqrt{a_{18}} - \sqrt{a_{17}}) \right)$$

$$\frac{12}{d} (\sqrt{a_{18}} - \sqrt{a_{10}})$$

$$\frac{12}{9} (\sqrt{a_{11} + 7d} - \sqrt{a_{11} - d})$$

$$\frac{4}{3} (9 - 3) = 4 \cdot 2 = 8 \text{ Ans}$$

QUESTION



Given $a_1, a_2, a_3, \dots, a_n$ in A.P. prove that

$$\frac{1}{a_1 a_n} + \frac{1}{a_2 a_{n-1}} + \frac{1}{a_3 a_{n-2}} + \dots + \frac{1}{a_n a_1} = \frac{2}{a_1 + a_n} \left[\frac{1}{a_1} + \frac{1}{a_2} + \dots + \frac{1}{a_n} \right]$$

$m+n = p+q$
 $a_m + a_n = a_p + a_q$
 $\underbrace{a_1 + a_n}_{n+1} = \underbrace{a_2 + a_{n-1}}_{n+1}$

LHS

$$\frac{a_1 + a_n}{a_1 + a_n} \left(\frac{1}{a_1 a_n} + \frac{1}{a_2 a_{n-1}} + \frac{1}{a_3 a_{n-2}} + \dots + \frac{1}{a_n a_1} \right) = \frac{1}{a_1 + a_n} \left(\frac{a_1 + a_n}{a_1 a_n} + \frac{a_1 + a_n}{a_2 a_{n-1}} + \frac{a_1 + a_n}{a_3 a_{n-2}} + \dots + \frac{a_1 + a_n}{a_n a_1} \right)$$

$$\frac{1}{a_1 + a_n} \left(\frac{a_1 + a_n}{a_1 a_n} + \frac{a_2 + a_{n-1}}{a_2 a_{n-1}} + \frac{a_3 + a_{n-2}}{a_3 a_{n-2}} + \dots + \frac{a_1 + a_n}{a_1 a_n} \right)$$

$$\frac{1}{a_1 + a_n} \left(\left(\frac{1}{a_n} + \frac{1}{a_1} \right) + \left(\frac{1}{a_{n-1}} + \frac{1}{a_2} \right) + \left(\frac{1}{a_{n-2}} + \frac{1}{a_3} \right) + \dots + \left(\frac{1}{a_1} + \frac{1}{a_n} \right) \right)$$

$$\frac{1}{a_1 + a_n} \cdot 2 \cdot \left(\frac{1}{a_1} + \frac{1}{a_2} + \frac{1}{a_3} + \dots + \frac{1}{a_n} \right)$$

QUESTION [JEE Mains 2021 (August)]



Tan 04

Let $a_1, a_2, \dots, a_{21}$ be an A.P. such that $\sum_{n=1}^{20} \frac{1}{a_n a_{n+1}} = \frac{4}{9}$.

If the sum of this A.P. is 189, then $a_6 \cdot a_{16}$ is equal to:

- A** 57 $S = \frac{1}{a_1 a_2} + \frac{1}{a_2 a_3} + \frac{1}{a_3 a_4} + \dots + \frac{1}{a_{20} a_{21}} = \frac{4}{9}$
- B** 72
- C** 48
- D** 36

QUESTION [JEE Mains 2024 (29 Jan)]



If $\log_e a, \log_e b, \log_e c$ are in an A.P. and $\log_e a - \log_e 2b, \log_e 2b - \log_e 3c, \log_e 3c - \log_e a$ are also in an A.P., then $a : b : c$ is equal to

- A** 6 : 3 : 2
- B** 9 : 6 : 4
- C** 25 : 10 : 4
- D** 16 : 4 : 1

$$2 \log_e b = \log_e a + \log_e c$$

$$\log_e b^2 = \log_e ac$$

$$b^2 = ac$$

$$9\lambda^2 = a \cdot 2\lambda$$

$$a = \frac{9}{2}\lambda$$

$$a : b : c = \frac{9}{2}\lambda : 3\lambda : 2\lambda = 9 : 6 : 4$$

$$2 \log_e \frac{2b}{3c} = \log_e \frac{a}{2b} + \log_e \frac{3c}{a}$$

$$\left(\frac{2b}{3c}\right)^2 = \frac{a}{2b} \cdot \frac{3c}{a}$$

$$8b^3 = 27c^3$$

$$2b = 3c = 6\lambda$$

$$b = 3\lambda$$

$$c = 2\lambda$$

Ans. B

If $\log_3 2, \log_3(2^x - 5), \log_3\left(2^x - \frac{7}{2}\right)$ are in an arithmetic progression, then the value of x is equal to

QUESTION



How many terms of the series $20 + 19\frac{1}{3} + 18\frac{2}{3} + \dots$ must be taken so that sum is 300.
Explain the reason of double answer.

$$S_n = 300.$$

$$\frac{n}{2} \left(40 + (n-1) \cdot \frac{2}{3} \right) = 300.$$

$$n \left(20 - \frac{n-1}{3} \right) = 300.$$

$$n(61-n) = 900.$$

$$-n^2 + 61n = 900$$

$$n^2 - 61n + 900$$

$$n = 25, 36.$$

$$d = -\frac{2}{3}$$

$$\underbrace{T_1, T_2, \dots, T_{25}}_{S_{25} = 300} \quad \underbrace{T_{26}, T_{27}, \dots, T_{31}}_{+ve} \quad \underbrace{T_{32}, T_{33}}_{-ve}$$

$\downarrow$
 $S_{36} = 300.$

$$T_n \leq 0$$

$$20 + (n-1) \left(-\frac{2}{3} \right) \leq 0$$

$$10 \leq \frac{n-1}{3}$$

$$n \geq 31$$

$$(b) S_{MAX} = S_{30} = S_{31}$$


$$S_{30} = \frac{30}{2} \left(40 + 29 \cdot \left(-\frac{2}{3}\right) \right)$$

$$= \frac{15 (120 - 58)}{3}$$

$$= 5 \cdot 62$$

$$S_{MAX} = 310$$

Sum inc till terms are +ve



G.P



G.P



Geometric Progression



Definition :

G.P. is the collection of non-zero terms in which each bears the same constant ratio with its immediately preceding term the series is called a G.P. and the constant ratio is called the common ratio.

Standard appearance of a G.P, where a is first term and r is common ratio is

$$\text{Agli Term} \div \text{Peechi Term} = \text{constant} = \text{common ratio}$$

$$\text{G.P: } a, ar, ar^2, ar^3, ar^4, \dots$$

$$T_1 = a \quad \text{common ratio} = r$$

$$T_n = ar^{n-1}$$

None of the terms of G.P
can be zero

Sum of n terms of G.P

$$S_n = a + ar + ar^2 + ar^3 + \dots + ar^{n-1}$$

$$rS_n = ar + ar^2 + ar^3 + \dots + ar^{n-1} + ar^n$$

$$(1-r)S_n = a - ar^n$$

$$S_n = \frac{a(1-r^n)}{1-r} = \frac{a(r^n-1)}{r-1}, \quad r \neq 1$$

$$S_n = \begin{cases} \frac{a(r^n-1)}{r-1} & r \neq 1 \\ na & r = 1 \end{cases}$$

If $r=1$

$$S = a + a + \dots + a = na$$

n terms.

* A G.P of +ve terms is inc if $r > 1$

* A G.P of +ve terms is dec if $0 < r < 1$

* A G.P of -ve terms is inc if $0 < r < 1$

* A G.P of -ve terms is dec if $r > 1$

* If $r < 0$ then G.P is alternatingly composed of
+ve & -ve terms Ex: $1, -2, 4, -8, 16, -32, \dots$

* If $r = 1$ then it is a constant G.P.


Sum of Infinite Terms



$$S_{\infty} = a + ar + ar^2 + ar^3 + \dots - \infty \quad 0 < |r| < 1$$

$$S_{\infty} = \lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \frac{a(1-r^{n+1})}{1-r} = \frac{a}{1-r} \quad -1 < r < 1, r \neq 0$$

$$S_{\infty} = \frac{a}{1-r} \quad -1 < r < 1, r \neq 0$$





* k^{th} from End in G.P of n terms = $(n-k+1)^{\text{th}}$ term from Beginning

* T_k from End = $l \cdot \left(\frac{1}{r}\right)^{k-1}$

$$\begin{array}{c} \xrightarrow{r} \qquad \qquad \qquad \xleftarrow{1/r} \\ a, ar, ar^2, ar^3, \dots, l \end{array}$$



Properties of G.P.



Property 1:

In an G.P. product of k^{th} term from beginning and k^{th} term from the last is always constant which equal to product of first term and last term.

$$T_k \cdot T_{n-k+1} = \text{constant} = a \cdot l$$



$$T_k = a \cdot r^{k-1}$$

$$T_k \text{ from End} = l \cdot (1/r)^{k-1}$$

$$T_k \cdot (T_k \text{ from End}) = a l.$$

If $m, n, p, q \in \mathbb{N}$ & $m+n = p+q$ then $T_m \cdot T_n = T_p \cdot T_q$ for a G.P

(prove yourself)

Property 2:

Three numbers in G.P.

$$: \quad a/r, a, ar \quad \text{---} \quad CR=r$$

Five numbers in G.P.

$$: \quad a/r^2, a/r, a, ar, ar^2 \quad \text{---} \quad CR=r$$

Four numbers in G.P.

$$: \quad a/r^3, a/r, ar, ar^3 \quad \text{---} \quad CR=r^2$$

Six numbers in G.P.

$$: \quad a/r^5, a/r^3, a/r, ar, ar^3, ar^5 \quad \text{---} \quad CR=r^2$$

If product of first few terms is given

first term is not a .



Sabse Important Baat



Sabhi Class Illustrations Retry Karnay hai...

Solution to Previous TAH

Suppose that all the terms of an arithmetic progression (A.P.) are natural numbers. If the ratio of the sum of the first seven terms to the sum of the first eleven terms is $6 : 11$ and the seventh term lies in between 130 and 140, then the common difference of this A.P. is

Lecture -02 (Sequence & Series)

JEE Adv. -2015

Tab-01 Suppose that all the terms of an arithmetic progression (A.P.) are natural numbers. If the ratio of the sum of the first seven terms to the sum of the first 11 terms is 6:11 and the seventh term lies between 130 and 140, then the common difference of this A.P. is:-

$$\Rightarrow \frac{\frac{7}{2}(2a+6d)}{\frac{11}{2}(2a+10d)} = \frac{6}{11} \quad \& \quad \boxed{130 < a+6d < 140}$$

$$\Rightarrow \frac{7(2a+6d)}{2a+10d} = 6$$

$$\Rightarrow \frac{7(a+3d)}{a+5d} = 6 \Rightarrow \boxed{7a+21d = 6a+30d}$$

$$\boxed{a = 9d}$$

$$\text{as, } 130 < a+6d < 140$$

$$130 < 15d < 140$$

$$\frac{130}{15} < d < \frac{140}{15}$$

$$\frac{26}{3} < d < \frac{28}{3}$$

as the terms of the A.P. are natural numbers

$$\boxed{d = \frac{27}{3} = 9} \quad (\text{Ans})$$

Kritisha

Page No.

Date 23/MAY

Tab-01

Q) Suppose that all the terms of an A.P. are natural numbers. If the ratio of the sum of the first seven terms to the first eleven terms is 6:11 and the seventh term lies between 130 and 140, then the common difference of the A.P. is:-

Solⁿ:

$$\begin{aligned} \text{all terms of AP} & \quad \frac{7}{2}(2a+6d) = 6 \\ \text{are natural given} & \quad \frac{11}{2}(2a+10d) = 11 \end{aligned}$$

$$\frac{2a+6d}{2a+10d} = \frac{6}{11} \quad 130 < T_7 = a+6d < 140$$

$$\frac{a+3d}{a+5d} = \frac{6}{11}$$

$$7a+21d = 6a+30d$$

$$a = 9d$$

$$T_7 = a+6d = 15d$$

factor 15's number lying between 130 and 140 is 135
so the value of 15d is 135 $\Rightarrow d = 9$

$$a = 9d \Rightarrow a = 81 \text{ and } d = 9$$

Suppose $a_1, a_2, \dots, a_n, \dots$ be an arithmetic progression of natural numbers. If the ratio of the sum of first five terms to the sum of first nine terms of the progression is $5 : 17$ and, $110 < a_{15} < 120$, then the sum of the first ten terms of the progression is equal to

- A** 290
- B** 380
- C** 460
- D** 510

Tah2

$a_1, a_2, \dots, a_n \rightarrow A.P.$

$$\frac{S_5}{S_9} = \frac{5}{17}$$

$$110 < a_{15} < 120$$

$$110 < a + 14d < 120$$

$$\frac{5a + 10d}{9a + 36d} = \frac{5}{17}$$

$$110 < a + 56d < 120$$

$$\frac{110}{57} < a < \frac{120}{57}$$

$$\frac{a + 2d}{a + 4d} = \frac{9}{17}$$

$$1.9 < a < 2.1$$

$$17a - 9a = 36d - 34d$$

$$a = 2, d = 8$$

$$4a = d$$

SAKSHI

Now we find Sum of first ten terms $= S_{10} = a_1 + a_2 + a_3 + \dots + a_{10}$

$$S_{10} = \frac{10}{2} [2 \times 2 + 9 \times 8]$$

$$S_{10} = 5(4 + 72) = 380 \text{ Ans (B)}$$

Tah-02 : Suppose $a_1, a_2, \dots, a_n$ be an arithmetic progression of natural numbers. If the ratio of the sum of first five numbers to the sum of first nine terms of the progression is $5:17$ and $110 < a_{15} < 120$, then the sum of the first ten terms of the progression is equal to:

$$\Rightarrow \frac{S_5}{S_9} = \frac{\frac{5}{2}(2a + 4d)}{\frac{9}{2}(2a + 8d)} = \frac{5}{17}$$

$$\Rightarrow \frac{a + 2d}{9(a + 4d)} = \frac{1}{17}$$

$$\Rightarrow 17a + 34d = 9a + 36d$$

$$\Rightarrow 8a = 2d$$

$$a_{15} = (a + 14d) \rightarrow \text{acc. to the question,}$$

$$110 < (a + 14d) < 120$$

$$110 < 57a < 120$$

$$\frac{110}{57} < a < \frac{120}{57}$$

Kritisha

as $a \in \mathbb{N}$

$$a = \frac{114}{57} = 2; d = 8$$

$$S_{10} = 5(4 + 9(8)) = 5(4 + 72) = 20 + 360 = 380 \text{ (B)}$$

Ans.

Consider an A.P. of positive integers, whose sum of the first three terms is 54 and the sum of the first twenty terms lies between 1600 and 1800. Then its 11th term is :

- A** 108
- B** 90
- C** 122
- D** 84

TAM-03

$$S_3 = 54$$

$$\frac{3}{2}(2a+2d) = 54$$

$$3(a+d) = 54$$

$$a+d = 18$$

$$\boxed{a = 18-d}$$

$$1600 < S_{20} < 1800 \quad \text{--- ①}$$

$$\therefore S_{20} = \frac{20}{2}(2a+19d)$$

$$10(2(18-d)+19d)$$

$$10(36-2d+19d)$$

$$360 + 170d$$

put in ①

$$1600 < 360 + 170d < 1800$$

$$1240 < 170d < 1440$$

$$7.2 < d < 8.4$$

$$\boxed{d = 8}$$

$$\boxed{a = 10}$$

$$a_{11} = a + 10d$$

$$= 10 + 10(8)$$

$$\boxed{\text{Ans} = 90}$$

Tah3

$$a_1 + a_2 + a_3 = 54$$

$$1600 < S_{20} < 1800$$

$$a_{11} = ?$$

$$a + a + d + a + 2d = 54$$

$$1600 < \frac{20}{2}(2(18-d) + 19d) < 1800$$

$$3a + 3d = 54$$

$$1600 < 360 + 170d < 1800$$

$$a + d = 18$$

$$a = 18 - d$$

$$\frac{1240}{170} < d < \frac{1440}{170}$$

$$\boxed{a = 10}$$

$$7.2 < d < 8.4$$

$$\text{Now, } a_{11} = a + 10d$$

$$a_{11} = 10 + 80$$

$$\boxed{d = 8}$$

$$\boxed{a_{11} = 90}$$

SAKSHI

Tah-03 Consider an A.P. of ~~positive~~ positive integers, whose sum of the first three terms is 54 and the sum of the first twenty terms lies between 1600 and 1800. Then its 11th term is :-

$$\Rightarrow S_3 = \frac{3}{2}(2a + 2d) = 54 \quad ; \quad a + d = 18$$

$$S_{20} = \frac{20}{2}(2a + 19d)$$

$$1600 < 10(2a + 19d) < 1800$$

$$160 < (2a + 19d) < 180$$

$$160 < 2(18 - d) + 19d < 180$$

$$160 < 36 + 17d < 180$$

$$\frac{124}{17} < d < \frac{144}{17}$$

$$\begin{matrix} d = 8 \\ a = 10 \end{matrix} \left[\begin{matrix} \text{as terms of A.P. are} \\ \text{+ve integers.} \end{matrix} \right]$$

$$t_{11} = a + 10d = 10 + 80 = \underline{\underline{90}} \quad (\text{B}) \quad (\text{Ans})$$

Let A.P. of positive integers

$$\text{Given, } S_3 = \frac{3}{2}(2a + 2d) = 54 \quad | \quad 18$$

$$a + d = 18$$

$$\text{and, } 1600 < S_{20} < 1800$$

Hint

$$1600 < \frac{20}{2}(2a + 19d) < 1800$$

$$2a + 19d$$

$$(a + d) + (a + d) + 17d$$

$$18 + 18 + 17d$$

$$36 + 17d$$

$$1600 < 10(36 + 17d) < 1800$$

$$160 - 36 < 17d < 180 - 36$$

$$\frac{124}{17} < d < \frac{144}{17}$$

tah-3
by vandana
from bihar

$$7.2 < d < 8.9$$

$$d = 8 \in \mathbb{I}, \quad a = 18 - 8 = 10$$

$$\begin{aligned} \text{Now, } t_{11} &= a + 10d \\ &= 10 + 10 \times 8 \\ &= 90 \quad \underline{\underline{\text{Ans}}} \end{aligned}$$

QUESTION [JEE Advanced 2018]



Let X be the set consisting of the first 2018 terms of the arithmetic progression 1, 6, 11,, and Y be the set consisting of the first 2018 terms of the arithmetic progression 9, 16, 23, Then, the number of elements in the set $X \cup Y$ is

TAH-04

$$\begin{aligned} X &= 1, 6, 11, 16, \dots & \text{upto 2018 terms} & d_1 = 5 \\ Y &= 9, 16, 23, \dots & & d_2 = 7 \end{aligned} \quad \} 35 = D$$

$$A = 16$$

$$D = 35$$

$$\begin{aligned} a_{n_1} &= a_1 + (n-1)d_1 \\ &= 1 + 2017(5) \\ &= 10086 \end{aligned}$$

$$\begin{aligned} a_{n_2} &= a_2 + (n-1)d_2 \\ &= 9 + 2017 \times 7 \\ &= 14128 \end{aligned}$$

$$16 + (n-1)35 \leq 10086$$

$$(n-1)35 \leq 10070$$

$$(n-1) \leq 287.7$$

$$n \leq 288.7$$

$$n \leq 288$$

$$\hookrightarrow (X \cap Y)$$

$$n(X) = 2018$$

$$n(Y) = 2018$$

$$n(X \cap Y) = 288$$

$$n(X \cup Y) = 2018 + 2018 - 288$$

$$n(X \cup Y) = 3748$$

Ans

TAH 04

Soln:

$$X: 1, 6, 11, 16, 17, \dots \text{ upto 2018 terms}$$

$$Y: 9, 16, 23, \dots \text{ upto 2018 terms}$$

$$\begin{aligned} d_x &= 5 \\ d_y &= 7 \end{aligned} \quad \} \text{ LCM} = 35$$

$$\therefore (X \cap Y) = 16, 54, \dots$$

$$X_{2018} = 1 + (2017 \times 5) = 10086$$

$$Y_{2018} = 9 + (2017 \times 7) = 14128$$

Let $(X \cap Y)$ has n terms.

$$(X \cap Y)_n = a + (n-1)d \leq 10086$$

$$16 + (n-1)(35) \leq 10086$$

$$(n-1) \leq \frac{10070}{35}$$

$$n-1 \leq 287.7 \dots$$

$$n-1 = 287$$

$$n = 288$$

$$\therefore n(X \cup Y) = n(X) + n(Y) - n(X \cap Y)$$

$$= 2018 + 2018 - 288$$

$$= 3748$$

$$\text{Ans } 3748$$

RASIDUL

Tab 04 :- Let X be the set consisting of the first 2018 terms of the A.P. $1, 6, 11, \dots$ and Y be the set consisting of the first 2018 terms of the A.P. $9, 16, 23, \dots$. Then the no. of elements in the set $X \cup Y$ is

$\Rightarrow$ We have to find the common terms in X & Y

$$X = \{1, 6, 11, 16, \dots, 2018^{\text{th}} \text{ term}\}$$

$Y = \{9, 16, 23, \dots, 2018^{\text{th}} \text{ term}\}$ $\xrightarrow{\text{LCM of Common diff}}$ $\Rightarrow 35$

LCM of
Common diff
 $\Rightarrow 35$

hence, A.P, consisting of the common terms of X & Y would have common diff = 35 & 1st term = 16

Now, 2018th term of X $\rightarrow 1 + (2017)(5) = 10086$

$$n \quad n \quad n \quad \gamma \longrightarrow g + 2017(\gamma) = 14128$$

$$(\text{Last term of new AP}) < 10086 \quad \left[\begin{array}{l} 10086 > 16 + 35(n-1) \\ 10070 > n-1 \end{array} \right]$$

$$10086 > 16 + 35(n-1)$$

$$\frac{10070}{100} > n-1$$

$$m \leq 288.71$$

putting $n=288$

$$16 + 35(287) = 10061$$

$$(X \cap Y) = \{6, 51, 86, \dots, 1006\} \quad \& \quad n(X \cap Y) = 288 \quad \checkmark$$

Thus, $n(X \cup Y) = n(X) + n(Y) - 288$
 $= 2018 + 2018 - 288$

$$= 2018 + 2018 - 288$$

$$n(X \cup Y) = 3748 \text{ (Ans)}$$

Kritisha

Tahap 4

$$n(x \cup y) = n(x) + n(y) - n(x \cap y) \quad \text{--- (1)}$$

$$n(x) = 2018$$

$$n(y) = 2018$$

1, 6, 11, 16, ... $T_n = 10086$ and 9, 16, 23, 30, ... $T_n = 14128$

$d=5$

$$-d = 3$$

LCM(35)

SAKSHI

common $\Rightarrow 16, 51, 86, \dots, d=35$
A.p. $T_n < 10086$ an

A.p.

$$T_n \leq 10086 \text{ and } T_n \leq 14128$$

$$T_m \leq 10086$$

$$a + (n-1)d \leq 10086$$

$$16 + 35n - 35 \leq 10086$$

$$35n \leq 10105$$

$$n \leq \frac{10105}{35}$$

$$n \leq 288.71 \quad n \approx 288$$

from Eqn ①

$$\eta(X \cup Y) = 2 \times 2018 - 288$$

$$= 40.36 - 288$$

$$= 3748 \text{ dm}$$

The number of common terms in the progressions

4, 9, 14, 19,, up to 25th term and 3, 6, 9, 12,, up to 37th term is:

- A** 9
- B** 8
- C** 5
- D** 7

Ex 5

$$4, 9, 14, 19, \dots T_n = 124$$

$$\hookrightarrow \text{upto } 25^{\text{th}} \quad d=5$$

$$\text{and } 3, 6, 9, 12, \dots T_n = 111$$

$$\hookrightarrow \text{upto } 37^{\text{th}} \quad d=3$$

common diff. of ^{A.P.} consecutive terms = 15

$$\text{common A.P.} \rightarrow 9, 24, 39, \dots d=15$$

$$T_n \leq 111 \text{ \& } T_n < 124$$

$\downarrow$

$$9 + (n-1)15 \leq 111$$

SAKSHI

$$9 + 15n - 15 \leq 111$$

$$15n \leq 117$$

$$n \leq 7.8$$

almost $n=7$ is

$$\rightarrow T_n = 371$$

Tah-05 The no. of common terms in the progressions:
 $4, 9, 14, 19 \dots$ upto 25^{th} term
 $3, 6, 9, 12 \dots$ " 37^{th} term is

$$\Rightarrow (A.P)_1 = 4, 9, 14, 19 \dots 25^{\text{th}} \text{ term}$$

$$(A.P)_2 = 3, 6, 9, 12 \dots 37^{\text{th}} \text{ term}$$

Last term of $(A.P)_1 \Rightarrow 4 + 24 \times 5$
 $= 4 + 24(5) = 124$

Last term of $(A.P)_2 = 3 + 3(36) = 111$

Common diff. of new A.P. $\Rightarrow 15$

1st term " " " $\Rightarrow 9$

Last " " " " $\Rightarrow < 111$

Now,

$$111 > 9 + 15(n-1)$$

$$\frac{102 + 15}{15} > n$$

$$(n < 7.8) \Rightarrow \boxed{n=7} \text{ (Ans)}$$

Kritisha

[Tah-05]

Sol For 1st AP : $T_n = 4 + (n-1)5$
 $= 4 + 24 \times 5$
 $= 4 + 120$
 $= 124$

For 2nd AP : $T_n = 3 + (n-1)3$
 $= 3 + 36 \times 3$
 $= 3 + 108$
 $= 111$

AP : $4, 9, 14, 19, \dots, 124, d=5$ $>$ $(\text{com} = 15)$

AP : $3, 6, 9, 12, \dots, 111, d=3$

New AP : $9, 24, 39, \dots, T_n$

$$\Rightarrow T_n \leq 124 \quad \text{and} \quad T_n \leq 111$$

$$\Rightarrow 9 + (n-1)15 \leq 111$$

$$\Rightarrow (n-1)15 \leq 102$$

$$\Rightarrow (n-1) \leq \frac{102}{15}$$

$$\Rightarrow n-1 \leq 6.8$$

$$\Rightarrow n \leq 7.8$$

$$\boxed{n=7} \quad \underline{\underline{=}}$$

Question - [JEE Mains 2024] Tah - OS



4, 9, 14, 19 - - - - -

$$n = 25$$

$$a = 4$$

$$d = 5$$

$$\begin{aligned} a_n &= a + (n-1)d \\ &= 4 + 24 \times 5 \\ &= 124 \end{aligned}$$

3, 6, 9, 12 - - - - -

$$n = 37$$

$$a = 3$$

$$d = 3$$

$$\begin{aligned} a_n &= 3 + 36 \times 3 \\ &= 111 \end{aligned}$$

4, 9, 14, 19 - - - - - 124

$$d = 5$$

3, 6, 9, 12 - - - - - 111

$$d = 3$$

$$L.C.M = 15$$

$$a = 9, d = 15 \text{ and } T_n \leq 111$$

$$a + (n-1)d \leq 111$$

$$9 + 15(n-1) \leq 111$$

$$n \leq \frac{111 - 9 + 15}{15}$$

$$n \leq \frac{117}{15}$$

$$n \leq 7$$

Option D

manik bihar

The sum of the common terms of the following three arithmetic progressions.

3, 7, 11, 15, ..., 399,

2, 5, 8, 11, ..., 359 and

2, 7, 12, 17, ..., 197,

is equal to _____

Qn-06 The sum of the common terms of the following three arithmetic progressions

$$3, 7, 11, 15, \dots, 399$$

$$2, 5, 8, 11, \dots, 359 \text{ and}$$

$$2, 7, 12, 17, \dots, 197 \text{ is equal to}$$

$$\begin{aligned} &\Rightarrow \begin{array}{l} 3, 7, 11, 15, \dots, 399 \\ 2, 5, 8, 11, \dots, 359 \\ 2, 7, 12, 17, \dots, 197 \end{array} \end{aligned}$$

$\text{LCM}(4, 3) = 12$
 com. terms $\rightarrow 11, 23, 35, \dots, 47, \dots$
 $\text{LCM}(3, 5) = 15$
 com. terms $\rightarrow 2, 17, 32, 47, \dots$

common terms in all three A.P

$$\Rightarrow \text{LCM}(12, 15)$$

$$\Rightarrow 60$$

New A.P $\Rightarrow (47, 107, 167)$ LAST TERM CAN'T EXCEED 197

Thus, sum of all the terms $\rightarrow 47 + 107 + 167 = 321$ (Ans)

Kritisha

Question JEE Mains 2023 (1 Feb)

$$(d=4) \leftarrow P = 3, 7, 11, 15, \dots, 399$$

$$(d=3) \leftarrow Q = 2, 5, 8, 11, \dots, 359$$

$$(d=5) \leftarrow R = 2, 7, 12, 17, \dots, 197$$

For Q & P common term = 11

common difference = $L.C.M = 4 \times 3 = 12$

$$(d=12) \leftarrow PQ \Rightarrow 11, 23, 35, 47, 59, 71, \dots, 359$$

Q or R & P, common term = 2

" difference = $L.C.M = 3 \times 5 = 15$

$$(d=15) \leftarrow QR \Rightarrow 2, 17, 32, 47, 62, \dots, 197$$

(a) common terms = 47

(d) " difference = $L.C.M = 12, 15 = 60$

$$T_n \leq 197$$

$$47 + (n-1)60 \leq 197$$

$$n \leq \frac{197 - 47 + 60}{60}$$

$$n \leq \frac{210}{60}$$

$$n \leq 3.$$

Sum of common term $S_3 = \frac{3}{2}(2a + (n-1)d)$

$$= \frac{3}{2}(2 \times 47 + 2 \times 60)$$

$$= 3(47 + 60)$$

$$= 321 \checkmark$$

manik bihar



QUESTION [JEE Mains 2025 (4 April)]



Let $A = \{1, 6, 11, 16, \dots\}$ and $B = \{9, 16, 23, 30, \dots\}$ be the sets consisting of the first 2025 terms of two arithmetic progressions. Then $n(A \cup B)$ is

- A** 3814
- B** 4003
- C** 4027
- D** 3761

Ans. D

Ques-07 Let $A = \{1, 6, 11, 16, \dots\}$ and $B = \{9, 16, 23, 30, \dots\}$ be the sets consisting of the first 2025 terms of two arithmetic progressions. Then $n(A \cup B)$ is

$$\Rightarrow A = \{1, 6, 11, 16, \dots, 2025^{\text{th}} \text{ term}\}$$

$$B = \{9, 16, 23, 30, \dots, 2025^{\text{th}} \text{ term}\}$$

$$n(A \cup B) = n(A) + n(B) - n(A \cap B)$$

$$\text{Last term of } A \Rightarrow 1 + 2024(5) = 10121$$

$$\text{" " " } B \Rightarrow 9 + 2024(7) = 14177$$

$$\text{common diff. of } (A \cap B) \Rightarrow 35$$

$$\text{1st term " } (A \cap B) \Rightarrow 16$$

$$\text{Last " " } (A \cap B) < 10121$$

$$35 \Rightarrow 16 + 35(n-1) < 10121$$

$$35(n-1) < 10105$$

$$(n-1) < 288.71$$

$$n < 289.71$$

$$n = 289$$

$$n(A \cap B) = 289$$

$$\text{hence, } n(A \cup B) = (2025 \times 2) - 289$$

$$n(A \cup B) = 3761$$

Kritisha

TAH-07
Question [JEE Mains 2025 (4 April)]

$$A = 1, 6, 11, 16, \dots$$

$$a = 1, d = 5, n = 2025$$

$$a_n = 1 + (2024) \times 5$$

$$= 1 + 10110$$

$$= 10111$$

Now

$$1, 6, 11, 16, \dots, 10111$$

$$B = 9, 16, 23, 30, \dots$$

$$a = 9, d = 7, n = 2025$$

$$a_n = 9 + (2024) \times 7$$

$$= 9 + 14168$$

$$= 14177$$

Now

$$9, 16, 23, 30, \dots, 14177$$

$$\text{common term } (a) = 16$$

$$\text{common difference } (d) = L.C.M. = 5 \times 7 = 35$$

$$T_n \leq 10111$$

$$a + (n-1)d \leq 10111$$

$$16 + (n-1)35 \leq 10111$$

$$n \leq \frac{10111 - 16 + 35}{35}$$

$$n \leq \frac{10130}{35}$$

$$\begin{array}{r} 35 \overline{) 10130} \quad (289) \\ \underline{70} \\ 313 \\ \underline{280} \\ 330 \end{array}$$

$$(A \cap B) \rightarrow n \leq 289$$

$$(A \cup B) = n(A) + n(B) - (A \cap B)$$

$$= 2025 + 2025 - 289$$

$$= 4050 - 289$$

$$= 3761$$

Tah-07

let $A = \{1, 6, 11, 16, \dots\}$ and $B = \{9, 16, 23, 30, \dots\}$ be the sets consisting of the first 2025 terms of two A.P.

Then $n(A \cup B)$ is

A. 3814

C. 4027

B. 4003

~~D. 3761~~

$$A = \{1, 6, 11, 16, \dots\} \rightarrow d_1 = 5$$

$$B = \{9, 16, 23, 30, \dots\} \rightarrow d_2 = 7$$

$$A \cap B = \{16, 51, 86, \dots\} \quad d = 35$$

$$a = 16$$

2025th term of A

$$d = 35$$

$$T_{2025} = 16 + (2024)5$$

2025th term of B

$$T_{2025} = 10121$$

$$T_{2025} = 9 + (2024)7$$

$$T_{2025} = 9 + 14168 \Rightarrow 14177$$

Let n^{th} term of $A \cap B = 10191$

$$a_n = a + (n-1)d$$

$$10191 = 16 + (n-1)35$$

$$n-1 = \frac{10191-16}{35}$$

tah07

$$n-1 = \frac{10105}{35}$$

Richathakur

$$n = 289$$

$$n(A \cup B) = n(A) + n(B) - n(A \cap B)$$

$$n(A \cup B) = 2025 + 2025 - 289$$

$$n(A \cup B) = 4050 - 289$$

$$n(A \cup B) = 3761 \quad (D)$$

Exams-2021

QUESTION [JEE Mains 2021 (31 Aug)]



Let $a_1, a_2, a_3, \dots$ be an A.P. If $\frac{a_1 + a_2 + \dots + a_{10}}{a_1 + a_2 + \dots + a_p} = \frac{100}{p^2}$, $p \neq 10$, then $\frac{a_{11}}{a_{10}}$ is equal to :

- A** $\frac{19}{21}$
- B** $\frac{100}{121}$
- C** $\frac{21}{19}$
- D** $\frac{121}{100}$

Ans. C

Tab-08

Let $a_1, a_2, a_3, \dots$ be an A.P. If $\frac{a_1 + a_2 + \dots + a_{10}}{a_1 + a_2 + \dots + a_p} = \frac{100}{p^2}$

$(p \neq 10)$ then $\frac{a_{11}}{a_{10}}$ is equal to:-

$$\Rightarrow \frac{a_1 + a_2 + a_3 + \dots + a_{10}}{a_1 + a_2 + \dots + a_p} = \frac{100}{p^2} \quad (p \neq 10)$$

$$\Rightarrow \frac{\frac{10}{2} (2a_1 + 9d)}{\frac{p}{2} (2a_1 + (p-1)d)} = \frac{100}{p^2}$$

$$\Rightarrow \frac{2a_1 + 9d}{2a_1 + (p-1)d} = \frac{10}{p}$$

$$2a_1 p + 9dp = 2a_1 (p-1) + 10d(p-1)$$

$$\Rightarrow 2a_1 (10-p) + dp - 10d = 0$$

$$\Rightarrow 2a_1 (10-p) - d(10-p) = 0$$

$$\Rightarrow (2a_1 - d)(10-p) = 0$$

$p \neq 10$ hence, $2a_1 = d \Rightarrow a_1 = \frac{d}{2}$

$$a_{10} = a_1 + 9d$$

$$a_{11} = a_1 + 10d$$

$$\frac{a_{11}}{a_{10}} = \frac{a_1 + 10d}{a_1 + 9d} = \frac{21a_1}{19a_1} = \frac{21}{19} \quad (\text{Ans})$$

Kritisha



Tab-08

Let $a_1, a_2, a_3, \dots$ be an A.P. If $\frac{a_1 + a_2 + \dots + a_{10}}{a_1 + a_2 + \dots + a_p} = \frac{100}{p^2}$, $p \neq 10$, then $\frac{a_{11}}{a_{10}}$ equal to

- (A) $19/21$ (B) $100/121$ (C) $21/19$ (D) $121/100$

$$\frac{a_1 + a_2 + \dots + a_{10}}{a_1 + a_2 + \dots + a_p} = \frac{100}{p^2}$$

$$\frac{\frac{10}{2} [2a_1 + 9d]}{\frac{p}{2} [2a_1 + (p-1)d]} = \frac{100}{p^2}$$

$$\Rightarrow \frac{10 [2a_1 + 9d]}{p [2a_1 + (p-1)d]} = \frac{100}{p^2}$$

$$2a_1 p + 9dp = 20a_1 + 10dp - 10d$$

$$2a_1 (10-p) + dp - 10d = 0$$

$$2a_1 (10-p) - d(10-p) = 0$$

$$(2a_1 - d)(10-p) = 0$$

$$a_1 = \frac{d}{2} \quad ; \quad p = 10$$

not possible $p \neq 10$

$$\frac{d}{a_1} = 2$$

$$\frac{1 + 10 \times 2}{1 + 9 \times 2} \Rightarrow \frac{21}{19} \quad (A)$$

$$\frac{a_{11}}{a_{10}} = \frac{21}{19}$$

Solution to Previous KTKs

For $a, b \in \mathbb{R} - \{0\}$, let $f(x) = ax^2 + bx + a$ satisfies $f\left(x + \frac{7}{4}\right) = f\left(\frac{7}{4} - x\right) \forall x \in \mathbb{R}$. Also the equation $f(x) = 7x + a$ has only one real and distinct solution.

The value of $(a + b)$ is equal to

- A** 4
- B** 5
- C** 6
- D** 7

For $a, b \in \mathbb{R} - \{0\}$, let $f(x) = ax^2 + bx + a$ satisfies $f\left(x + \frac{7}{4}\right) = f\left(\frac{7}{4} - x\right) \forall x \in \mathbb{R}$. Also the equation $f(x) = 7x + a$ has only one real and distinct solution.

The minimum value $f(x)$ in $\left[0, \frac{3}{2}\right]$ is equal to

A $-\frac{33}{8}$

B 0

C 4

D -2

KTK-1

For $a, b \in \mathbb{R} - \{0\}$, let $P(x) = ax^2 + bx + a$ satisfies $P\left(x + \frac{7}{4}\right) = P\left(\frac{7}{4} - x\right) \forall x \in \mathbb{R}$. Also the eqⁿ $P(x) = 7x + a$ has only one real root and distinct solⁿ. Then value of $(a+b) = ?$

(A) 4

(B) 5

(C) 6

(D) 7

$$P\left(x + \frac{7}{4}\right) = P\left(\frac{7}{4} - x\right)$$

$\frac{7}{4} \rightarrow$ Symmetry

$$P(x) = ax^2 + bx + a$$

$$\frac{-b}{2a} = \frac{7}{4 \cdot 2}$$

$$b = \frac{-7a}{2}$$

$$P(x) = 7x + a$$

$$ax^2 + bx + a = 7x + a$$

$$ax^2 + x(b-7) = 0$$

$$(b-7)^2 - 4(a)(0) = 0$$

$$b^2 - 14b + 49 = 0 \quad (b-7)^2 = 0$$

$$b = \frac{-7 \times a}{2}$$

$$b = 7$$

$$a = \frac{-7}{2}$$

$$a+b \Rightarrow -2+7$$

$$a+b = 5$$

$$\boxed{a = -2}$$

$$\boxed{b = 7}$$

KTK-2

For $a, b \in \mathbb{R} - \{0\}$ - Same question
The minimum value $P(x)$ in $\left[0, \frac{3}{2}\right]$ is equal to

(A) $-33/8$

(B) 0

(C) 4

(D) -2

$a = -2$ means downward parabola

$$P(x) = -2x^2 + 7x - 2$$

$$x = \frac{7}{4} \rightarrow \text{Vertex}$$

$$f(0) = -2$$

$$f\left(\frac{3}{2}\right) = -2 + 7 - 2 \Rightarrow 3$$

$$\text{min value} = -2$$

KTK-01

Q. For $a, b \in \mathbb{R} - \{0\}$, let $f(x) = ax^2 + bx + a$ satisfies;
 $f\left(x + \frac{7}{4}\right) = f\left(\frac{7}{4} - x\right) \forall x \in \mathbb{R}$. Also the equation
 $f(x) = 7x + a$ has only one real and distinct solⁿ.

* The value of $(a+b)$ is equal to:

$$\Rightarrow f(x) = ax^2 + bx + a, \text{ is satisfies: } f\left(x + \frac{7}{4}\right) = f\left(\frac{7}{4} - x\right)$$

$\Rightarrow$ To see this eqⁿ:

Clearly graph is sym about

$$\boxed{x = \frac{7}{4}}$$

$$\Rightarrow \frac{-b}{2a} = \frac{7}{4 \cdot 2}$$

$$\Rightarrow \boxed{7a = -2b}$$

$$\# f(x) = ax^2 + bx + a \text{ \& } f(x) = 7x + a$$

$$\Rightarrow 7x + a = ax^2 + bx + a$$

$$\Rightarrow ax^2 + (b-7)x = 0 \quad (\text{only one real \& distinct sol}^n)$$

$$\Rightarrow (b-7)^2 - 4 \cdot a \cdot 0 = 0$$

$$\Downarrow$$

$$\boxed{b=7}$$

$$\Rightarrow b-7 = 0$$

$$\Rightarrow \boxed{b=7}$$

$$\# 7a = (-2)(7)$$

$$7a = -14$$

$$\boxed{a = -2}$$

krish

$\Rightarrow$ find value of $(a+b)$.

$$a+b = -2+7$$

$$= \boxed{5} \text{ Ans.}$$

THANK
YOU